arXiv:1504.02775v1 [math.AP] 10 Apr 2015

Splash singularities for the free boundary Navier-Stokes equations

Angel Castro, Diego Córdoba, Charles Fefferman,
Francisco Gancedo and Javier Gómez-Serrano

April 14, 2015

Abstract

In this paper, we prove the existence of smooth initial data for the 2D free boundary incompressible Navier-Stokes equations, for which the smoothness of the interface breaks down in finite time into a splash singularity.

Keywords: singularities, splash, Navier-Stokes, free boundary, incompressible

1 Introduction

In this paper, we prove that an initially smooth solution of the 2D water wave equation with non-zero viscosity may break down in finite time by forming a splash singularity, see Figure 1.

The analogous result for inviscid water waves was proven in our previous paper [9]. The strategy there was to start with a splash configuration and solve backwards in time. To do so, we first made a conformal map (essentially a branch of the square root) $P(z)$ from physical space to the "tilde domain" and then adapted the proof of Ambrose-Masmoudi [3] (see also [10]) of short time existence of solutions of the inviscid water wave equation.

The strategy of [9] cannot work for the present case of nonzero viscosity, since the equations cannot be solved backwards in time. We will instead make use of the transformation to the tilde domain in a new way, which we explain below.

We refer the reader to the further historical discussion at the end of the introduction, including references to alternate proofs by Coutand-Shkoller of several of our results.

Let us first present the inviscid and viscous water wave equations, next define a splash singularity, and then state our main result.

The 2D water wave equations govern a system consisting of water, vacuum and the interface between them. At time $t \in \mathbb{R}$, the water occupies a region $\Omega(t) \subset \mathbb{R}^2$, and the vacuum occupies the complementary region $\mathbb{R}^2 \setminus \Omega(t)$. For points x in the water region $\Omega(t)$, the velocity of the water at position x and time t is $u(x,t) \in \mathbb{R}^2$, and the pressure is $p(x,t) \in \mathbb{R}$. Thus, $u(x,t), p(x,t)$ are defined only for $x \in \Omega(t)$; finding $\Omega(t)$ is part of the problem.



Figure 1: How a splash forms.

We assume here that the interface $\partial\Omega(t) \subset \mathbb{R}$ is a smooth simple closed curve, which we write in parametric form:

$$\partial\Omega(t) = \{z(\alpha, t) : \alpha \in \mathbb{R}/\mathbb{Z}\},$$

where $z : \mathbb{R}/\mathbb{Z} \to \mathbb{R}^2$ is smooth and satisfies the chord-arc condition

$$|z(\alpha, t) - z(\alpha', t)| \geq CA(t)\|\alpha - \alpha'\| \text{ for } \alpha, \alpha' \in \mathbb{R}/\mathbb{Z}.$$

Here, $CA(t) > 0$ is the "chord-arc constant", and $\|\alpha - \alpha'\|$ denotes the distance from α to α' in $\mathbb{R}/\mathbb{Z}$.

The parametrization of the interface has no physical meaning, and can be picked to simplify our analysis.

The inviscid water wave equations are as follows:

$$\begin{aligned}
(\partial_t + u \cdot \nabla_x)u(x, t) &= -\nabla_x p(x, t) \text{ for } x \in \Omega(t) \\
\text{div} u(x, t) &= 0 \text{ for } x \in \Omega(t) \\
\text{curl} u(x, t) &= 0 \text{ for } x \in \Omega(t) \\
p(x, t) &= 0 \text{ for } x \in \partial\Omega(t) \\
\partial_t z(\alpha, t) &= u(z(\alpha, t), t) + c(\alpha, t)\partial_\alpha z(\alpha, t) \text{ for } \alpha \in \mathbb{R}/\mathbb{Z}.
\end{aligned} \tag{1}$$

(The last equation asserts that the interface moves with the fluid. The function $c(\alpha, t)$ affects only the parametrization of the interface, and may be chosen arbitrarily).

The initial conditions for inviscid water waves are as follows:

- $\Omega(0) = \Omega_0$ (a given domain bounded by a smooth simple closed curve).
- $u(x, 0) = u_0$ (a given smooth divergence-free irrotational vector field) for $x \in \Omega_0$.

For water waves with nonzero viscosity, the relevant equations take the following form in suitable units:

$$\begin{aligned}
(\partial_t + u \cdot \nabla_x)u(x, t) &= \Delta_x u(x, t) - \nabla_x p(x, t) \text{ for } x \in \Omega(t) \\
\text{div} u(x, t) &= 0 \text{ for } x \in \Omega(t) \\
\left(p\,\mathbb{I} - \left(\nabla u + (\nabla u)^*\right)\right) n &= 0, \text{ for } x \in \partial\Omega(t) \\
\partial_t z(\alpha, t) &= u(z(\alpha, t), t) + c(\alpha, t)\partial_\alpha z(\alpha, t) \text{ for } \alpha \in \mathbb{R}/\mathbb{Z}.
\end{aligned} \tag{2}$$

Again, $c(\alpha, t)$ may be chosen arbitrarily.

The initial conditions are:

- $\Omega(0) = \Omega_0$ (as before).
- $u(x, 0) = u_0$ for $x \in \Omega_0$, where u_0 is a given smooth divergence-free vector field on Ω_0, satisfying the constraint
- $n_0^\perp \left(\left(\nabla u_0 + (\nabla u_0)^*\right)\right) n_0 = 0$ on $\partial\Omega_0$.

Next, we adapt from [9] the definition of a splash singularity for the compact case:

Note that the inviscid water wave equations (1) have a symmetry under time reversal, but the viscous equations (2) have no such symmetry. This reflects the presence of the Euler equation in (1) and the Navier-Stokes equation in (2).

Definition 1.1 *We say that* $z(\alpha) = (z_1(\alpha), z_2(\alpha))$ *is a* splash curve *if*

1. $z_1(\alpha), z_2(\alpha)$ *are smooth functions and* 2π*-periodic.*
2. $z(\alpha)$ *satisfies the arc-chord condition at every point except at* α_1 *and* α_2*, with* $\alpha_1 < \alpha_2$ *where* $z(\alpha_1) = z(\alpha_2)$ *and* $|z_\alpha(\alpha_1)|, |z_\alpha(\alpha_2)| > 0$*. This means* $z(\alpha_1) = z(\alpha_2)$*, but if we remove either a neighborhood of* α_1 *or a neighborhood of* α_2 *in parameter space, then the arc-chord condition holds.*

3. *The curve $z(\alpha)$ separates the complex plane into two regions; a connected water region and a vacuum region (not necessarily connected). We choose the parametrization such that the normal vector $n = \frac{(-\partial_\alpha z_2(\alpha), \partial_\alpha z_1(\alpha))}{|\partial_\alpha z(\alpha)|}$ points to the vacuum region. We regard the interface to be part of the water region.*

4. *We can choose a branch of the function P on the water region such that the curve $\tilde{z}(\alpha) = (\tilde{z}_1(\alpha), \tilde{z}_2(\alpha)) = P(z(\alpha))$ satisfies:*
    - *(a) $\tilde{z}_1(\alpha)$ and $\tilde{z}_2(\alpha)$ are smooth and 2π-periodic.*
    - *(b) $\tilde{z}$ is a closed contour.*
    - *(c) $\tilde{z}$ satisfies the arc-chord condition.*

Here are two examples of curves which are not splash curves (see Figure 2).



Figure 2: Two examples of non splash curves.

We can now state the main result of this paper:

Theorem 1.2 *There exists a solution of the viscous water wave equation that remains smooth for times $t \in [0, t_*)$ but forms a splash singularity at time t_*.*

Next, we recall from [9] how to produce the inviscid water waves [1] that end in a splash at time $t_* > 0$. We start with the splash $\Omega(t_*), u(\cdot, t_*)$ and solve the inviscid equations (1) backwards in time. It is well known that the inviscid equations (1) can be solved (forward or backwards in time) starting from smooth initial data (See S. Wu [38] and [25] for a comprehensive list of references) The difficulty here is that the initial $\Omega(t_*)$ is singular. To overcome this difficulty, we make a slit Γ in the complex plane as in Figure 4, and then make the conformal mapping $\tilde{z} = P(z)$ for $z \in \mathbb{C} \setminus \Gamma$; here $P(z)$ is a branch of $\sqrt{z}$. The inverse map is simply $P^{-1}(\tilde{z}) \equiv \tilde{z}^2$, which of course is well defined and smooth on the whole complex plane.

We want a solution of (1) for which $\Omega(t_*)$ is as in Figure 4, but for times $t < t_*$ (t close to t_*), $\Omega(t)$ avoids the slit Γ. The corresponding domains $\tilde{\Omega}(t)$ in the tilde domain (i.e. $\tilde{\Omega}(t) = P(\Omega(t))$) behave as in Figure 1.

---

[1] We treat here the case in which the water region $\Omega(t) \subset \mathbb{R}^2$ is a bounded region. In [9], we studied the case in which $\Omega(t)$ is periodic with respect to horizontal translation as in Figure 3. In this introduction we ignore the distinction between the compact and periodic cases.



Figure 3: Horizontally periodic setting.



(a) The z-plane. (b) The $\tilde{z}$-plane.

Figure 4: Desingularization of the splash domain.



(a) (b)

Figure 5: Evolution of a splash in the tilde world for $t = t_*$ and $t < t_*$.

In [9] we give simple transformation laws that allow us to pass from the velocity $u(x,t)$ and pressure $p(x,t)$ (defined for $x \in \Omega(t)$) to a transformed velocity and pressure $\tilde{u}(\tilde{x},t), \tilde{p}(\tilde{x},t)$, defined for $\tilde{x} \in \tilde{\Omega}(t)$.

We can then write down the analogue of the inviscid water wave equations (1) in the tilde domain. We call these equations $(\tilde{1})$; they govern the evolution of $\tilde{\Omega}(t), \tilde{u}(\tilde{x},t)$ backwards in time, with initial conditions at $t = t_*$.

Note that, whereas $\Omega(t_*)$ is singular (its chord-arc constant is zero), $\tilde{\Omega}(t_*)$ is bounded by a smooth simple closed curve. Moreover, the transformed water wave equation $(\tilde{1})$ behave much like the original equations (1). Adapting the energy estimates of Ambrose and Masmoudi [3], we prove existence of a smooth solution of $(\tilde{1})$ for times $t \in [t_* - \varepsilon, t_* + \varepsilon]$, with $\tilde{\Omega}(t_*)$ as in Figure 4. As long as $\tilde{\Omega}(t)$ avoids $P(\Gamma)$ for all $t \in [t_* - \varepsilon, t_*)$, we obtain a corresponding solution of (1) with $\Omega(t) = P^{-1}(\tilde{\Omega}(t))$ a smooth simple closed curve for $t \in [t_* - \varepsilon, t_*)$, We can guarantee that $\tilde{\Omega}(t)$ avoids $P(\Gamma)$ for $t \in [t_* - \varepsilon, t_*)$ by taking ε smaller and carefully picking the initial velocity $\tilde{u}$ at the two points $\tilde{z}_{splash,1}$ and $\tilde{z}_{splash,2}$ in Figure 1. Thus, we have produced an inviscid water wave that starts out smooth at time $t_* - \varepsilon$ and ends in a splash at time t_*.

This concludes our discussion of the inviscid water wave equations (1).

We pass to the viscous case, where we no longer have the luxury of solving backwards in time. Just as in the inviscid case, there is an analogue of the water wave equations (2) in the tilde domain, which we call equations $(\tilde{2})$ (we won't write them out in the introduction, see section 2 for a precise definition).

The unknowns in the tilde domain are a time-varying domain $\tilde{\Omega}(t)$, a velocity field $\tilde{u}(\tilde{x},t)$ and a pressure $\tilde{p}(\tilde{x},t)$ with $\tilde{u}, \tilde{p}$ defined for $\tilde{x} \in \tilde{\Omega}(t)$.

The relationship between equations (2) and equations $(\tilde{2})$ is as follows:

Every solution $\Omega(t), u(\cdot,t), p(\cdot,t)$ of (2) such that $\Omega(t)$ avoids the slit Γ gives rise to a solution $\tilde{\Omega}(t), \tilde{u}(\cdot,t), \tilde{p}(\cdot,t)$ of $(\tilde{2})$, with $\tilde{\Omega}(t) = P(\Omega(t))$. On the other hand, let $\tilde{\Omega}(t), \tilde{u}(\cdot,t), \tilde{p}(\cdot,t)$ be a solution of $(\tilde{2})$. We would like to define a solution $\Omega(t), u(\cdot,t), p(\cdot,t)$ of (2) such that $\Omega(t) = P^{-1}(\tilde{\Omega}(t))$. However, this may not be possible, because $P^{-1}(\partial\tilde{\Omega}(t))$ may self-intersect, as in Figures 6(b) and 6(c). In particular, $P^{-1}(\partial\tilde{\Omega}(t))$ in Figure 6(c) is clearly not the boundary of any physically meaningful water



(a) (b) (c)

Figure 6: Possibilities for $P^{-1}(\tilde{\Omega}(t))$.

region.

The good news is that Figures 6(b) and 6(c) are the only obstructions; as long as $P^{-1}(\partial\tilde{\Omega}(t))$ is as in Figure 6(a), we can easily pass from our solution $\tilde{\Omega}(t), \tilde{u}(\cdot,t), \tilde{p}(\cdot,t)$ of $\widetilde{(2)}$ to a solution $\Omega(t), u(\cdot,t), p(\cdot,t)$ of (2) with $\Omega(t) = P^{-1}(\tilde{\Omega}(t))$.

Let us now solve equations $\widetilde{(2)}$ for times $t > 0$, starting with smooth initial $\tilde{\Omega}(0)$ and $\tilde{u}(\cdot,0)$. Adapting the analysis of Beale [5] from (2) to the tilde domain, we prove that smooth solutions $\tilde{\Omega}(t), \tilde{u}(\cdot,t), \tilde{p}(\cdot,t)$ of $\widetilde{(2)}$ with the given initial conditions exist for short time, i.e. for $t \in [0,T]$ with small positive T depending on $\tilde{\Omega}(0), \tilde{u}(\cdot,0)$. Moreover, the solutions of $\widetilde{(2)}$ depend stably on the initial conditions.

For suitable one-parameter families of initial conditions $\tilde{\Omega}_\varepsilon(0), \tilde{u}_\varepsilon(\cdot,0)$ depending on a small parameter ε, there is a family of smooth solutions $\tilde{\Omega}_\varepsilon(t), \tilde{u}_\varepsilon(\cdot,t), \tilde{p}_\varepsilon(\cdot,t)$ solving $\widetilde{(1)}$ up to time T, with $\|\partial\tilde{\Omega}_\varepsilon(t) - \partial\tilde{\Omega}(t)\| = O(\varepsilon)$ in a suitable norm $\|\cdot\|$.

We are ready to combine the ingredients above. We start with smooth initial conditions $\tilde{\Omega}(0), \tilde{u}(\cdot,0)$ with $P^{-1}(\partial\tilde{\Omega}(0))$ as in Figure 6(b). Solving $\widetilde{(2)}$ for a short time, we obtain smooth solutions $\tilde{\Omega}(t), \tilde{u}(\cdot,t), \tilde{p}(\cdot,t)$ for times $t \in [0,T]$ (some $T > 0$). By making T smaller and picking the initial velocity $\tilde{u}(\cdot,0)$ so that $\tilde{u}(z_{splash,1},0)$ and $\tilde{u}(z_{splash,2},0)$ point in the right direction, we can guarantee that $P^{-1}(\partial\tilde{\Omega}(T))$ behaves as in Figure 6(c).

Next, we pick a one-parameter family of initial conditions $\tilde{\Omega}_\varepsilon(0)$ and $\tilde{u}_\varepsilon(\cdot,0)$ perturbing our $\tilde{\Omega}(0), \tilde{u}(\cdot,0)$. We can easily arrange that for small positive ε, $P^{-1}(\partial\tilde{\Omega}_\varepsilon(0))$ looks like Figure 6(a), even though $P^{-1}(\partial\tilde{\Omega}(0))$ is as in Figure 6(b). The perturbed solution $\tilde{\Omega}_\varepsilon(t), \tilde{u}_\varepsilon(\cdot,t), \tilde{p}_\varepsilon(\cdot,t)$ will satisfy $\|\partial\tilde{\Omega}_\varepsilon(T) - \partial\tilde{\Omega}(T)\| = O(\varepsilon)$. Hence, for $\varepsilon > 0$ small enough, $P^{-1}(\partial\tilde{\Omega}_\varepsilon(T))$ will be as in Figure 6(c), since the same holds for $P^{-1}(\partial\tilde{\Omega}(T))$. For such ε, $P^{-1}(\partial\tilde{\Omega}_\varepsilon(t))$ starts out as in Figure 6(a) for $t = 0$ and ends as in Figure 6(c) for $t = T$. Fix such an ε and let

$$t_* = \inf\Big\{t \in [0,T] : P^{-1}(\partial\tilde{\Omega}_\varepsilon(t)) \text{ is as in Figure 6(b) or 6(c)}\Big\}$$

Then, $0 < t_* < T$, $P^{-1}(\partial\tilde{\Omega}_\varepsilon(t_*))$ is as in Figure 6(b), and $P^{-1}(\partial\tilde{\Omega}_\varepsilon(t))$ is as in Figure 6(a) for $0 \le t < t_*$. Consequently, $\tilde{\Omega}_\varepsilon(t), \tilde{u}_\varepsilon(\cdot,t), \tilde{p}_\varepsilon(\cdot,t)$ gives rise to a solution of (2), the viscous water wave equation, for $t \in [0, t_*)$, ending in a splash at time t_*.

We close the introduction by citing some of the relevant literature on viscous water waves.

Some of the earliest papers in this topic were written by Solonnikov. He studied the problem of a viscous fluid bounded by a free boundary in the vacuum (the fluid domain is bounded). He showed local existence of solutions with [32] using Hölder spaces in the frame of parabolic systems for bounded domains and without surface tension [33, 34]. Local existence and uniqueness of solutions was shown by Beale in [5] in Sobolev spaces with non-slip boundary condition at a regular bottom for the fluid and extended to L^q spaces by Abels [1]. A second theorem in the former paper showed that for any $T > 0$ there exist solutions of sufficiently small initial data on $[0,T]$. For the case with surface tension, see also Tani [36] and Coutand-Shkoller [12]. Masmoudi and Rousset [28] studied the case where the viscosity tends to zero for the free boundary problem (see [18] for the case with surface tension). The case of a viscous fluid lying above a bottom has been extensively studied. For the case without surface tension Sylvester, in [35], showed global well-posedness for small initial data. Hataya, in [23], showed the existence of solutions which decay algebraically in time for a periodic in the horizontal variable surface. Guo and Tice, in [22, 21, 20], have proved algebraic decay rate in time for asymptotically flat surfaces and almost exponential decay rate in time for periodic in the horizontal variable surfaces.

Global in time regularity was first given for small initial data in [6] by adding to the system surface tension effects (see also [4] for an alternative proof). In [35, 37] global existence is obtained without the help of surface tension. Decay rates have been also considered to understand the long time behavior of the solution. For the case with surface tension see [7, 30].

In the case of a two fluid problem there are some recent results where local well-posedness and global existence for small data is shown. The situations consider low regular initial velocities in critical spaces, in some cases within the chain of Besov spaces (see [16, 17, 24, 31, 26] and the references therein for more details).

Existence of splash singularities for inviscid water waves was proven in our paper [9], see also Coutand-Shkoller [13] for an alternate proof with applications to a 3D setting.

The inviscid splash is not prevented by taking into account gravity or surface tension (see [9, 8]), but it is prevented by replacing the vacuum in $\mathbb{R}^2 \setminus \Omega(t)$ by an incompressible opposing fluid; see [19] as well as an alternate proof by Coutand-Shkoller [14]. It has been shown in [11] that there exist “almost splash” stationary solutions in the two fluid case. We caution that the nonexistence of a splash does not rule out a breakdown in which the chord-arc constant degenerates and the solution loses smoothness; again, see [19].

Our strategy for the viscous splash, as outlined above, was announced at the *40èmes journées EDP (2013)* in Biarritz, at the *2013 Clay Research Conference* (Oxford) and more recently at the *Minerva Distinguished Visitor Lectures* in Princeton in 2014. Finally, in this paper we provide details.

We refer the reader to Coutand-Shkoller [15] for a different proof of the formation of splash singularities for viscous water waves.

2 Equations: transformation to a nonsplash domain

We have to solve the 2D-Navier-Stokes equations in the fluid domain $\Omega(t)$,

$$
\begin{aligned}
\partial_t u + (u\cdot\nabla)u - \Delta u &= -\nabla p, \quad \text{in } \Omega(t), && (3)\\
\nabla\cdot u &= 0, \quad \text{in } \Omega(t), && (4)\\
\left(p\,\mathbb{I} - \left(\nabla u + (\nabla u)^*\right)\right) n &= 0, \quad \text{on } \partial\Omega(t), && (5)\\
u(t)|_{t=0} &= u_0, \quad \text{in } \Omega(0) = \Omega_0, && (6)
\end{aligned}
$$

where

$$\Omega(t) = X(\Omega(0), t)$$

with $X(\alpha, t)$ solving

$$
\begin{aligned}
\frac{dX(\alpha,t)}{dt} =& u(X(\alpha,t),t)\\
X(\alpha,0) =& \alpha, \quad \alpha\in\Omega(0),
\end{aligned}
$$

and $n(t)$ is unit normal vector to $\partial\Omega(t)$ (pointing out). Here, Ω_0 and u_0 are given and satisfy the compatibility conditions

$$
\begin{aligned}
\nabla\cdot u_0 = 0, \quad \text{in } \Omega_0\\
n_0^\perp\left(\nabla u_0 + (\nabla u_0)^*\right) n_0 = 0 \quad \text{on } \partial\Omega_0,
\end{aligned}
$$

with $n_0 = n(0)$. The condition

$$\left(p\,\mathbb{I} - \left(\nabla u + (\nabla u)^*\right)\right)\big|_{\partial\Omega(t)} n = 0$$

states the continuity of the normal stress through the interface. We will use the symbol * to denote the transpose of a matrix.

We notice that the pressure, p, will be given by the following elliptic problem

$$
\begin{aligned}
-\Delta p =& \nabla\cdot((u\cdot\nabla)u) \quad \text{in } \Omega(t)\\
p =& n\left(\nabla u + (\nabla u)^*\right) n \quad \text{on } \partial\Omega(t).
\end{aligned}
$$

Let $P(z)$ be defined as in the introduction and let $\tilde{\Omega}(t) = P(\Omega(t))$. Next we will write the system (3-6) in the domain $\tilde{\Omega}(t)$ and after that, by using Lagrangian coordinates, we will fix the domain in order to work in the domain $\tilde{\Omega}(0)$ rather than in $\tilde{\Omega}(t)$.

Remark 2.1 *At this point one should notice that we are assuming that $\tilde{\Omega}(t)$ is the projection by P of $\Omega(t)$ and that $\tilde{\Omega}(t)$ is a simply connected bounded domain. Therefore P^{-1} is a well defined analytic function. Once we have written the N-S system in $\tilde{\Omega}(t)$ this assumption will not be needed anymore.*

We define $\tilde{u} = u\circ P^{-1}$. Then

$$\partial_j u^i = (\partial_k \tilde{u}^i\circ P)\partial_j P^k$$

and therefore

$$\partial_j u^i \circ P^{-1} = A_{kj}\partial_k \tilde{u}^i,$$

where

$$A_{kj} = \partial_j P^k \circ P^{-1}.$$

In addition, let's assume that we traverse (clockwise) the boundary of $\Omega(t)$ using the parametrization $z(\gamma, t)$, i.e.,

$$\partial\Omega(t) = \{z(\gamma,t)\in\mathbb{R}^2 \,:\, \gamma\in[-\pi,\pi)\times[0,T]\}.$$

and

$$n = (-z^2_\gamma(\gamma,t), z^1_\gamma(\gamma,t)).$$

Since the boundary of $\tilde{\Omega}(t)$ can be parameterized by $\tilde{z}(\gamma,t) = P(z(\gamma,t))$ we have that

$$\tilde{z}^i_\gamma = \partial_k P^i\circ P^{-1}(\tilde{z}(\gamma,t))z^k_\gamma(\gamma,t)$$

thus

$$\tilde{n} = -JA|_{\partial\tilde{\Omega}(t)} Jn,$$

where

$$J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Using the previous expression and the fact that P is a conformal map we can write the system (3-6) in the domain $\tilde{\Omega}(t)$ as follows:

$$\begin{aligned}
\partial_t \tilde{u} + (A\tilde{u}\cdot\nabla)\tilde{u} - Q^2\Delta\tilde{u} &= -\, A^*\nabla\tilde{p}, \quad \text{in } \tilde{\Omega}(t), && (7)\\
Tr\,(\nabla\tilde{u}A) &= 0, \quad \text{in } \tilde{\Omega}(t), && (8)\\
\left(\tilde{p}\mathbb{I} - \left(\nabla\tilde{u}A + (\nabla\tilde{u}A)^*\right)\right) A^{-1}\tilde{n} \quad &= 0, \quad \text{on } \partial\tilde{\Omega}(t), && (9)\\
\tilde{u}(t)|_{t=0} &= \tilde{u}_0, \quad \text{in } \tilde{\Omega}(0) = \tilde{\Omega}_0. && (10)
\end{aligned}$$

where $Tr(A)$ is the trace of the matrix A and

$$\tilde{\Omega}(t) = \tilde{X}(\tilde{\Omega}(0), t)$$

with $\tilde{X}(\alpha, t)$ solving

$$\begin{aligned}
\frac{d\tilde{X}(\alpha,t)}{dt} &= \left(A\circ\tilde{X}(\alpha,t)\right)\left(\tilde{u}\circ\tilde{X}(\alpha,t)\right)\\
\tilde{X}(\alpha,t) &= \alpha, \quad \alpha\in\tilde{\Omega}(0).
\end{aligned}$$

Here we have used that

$$Q^2 A^{-1} = -JAJ$$

and

$$Q^2 = \left|\frac{dP}{dz}\circ P^{-1}\right|^2.$$

Now we will fix the domain by working with the variables

$$\begin{aligned}
\tilde{v}(\alpha,t) &= \tilde{u}\circ\tilde{X}(\alpha,t) \quad \alpha\in\tilde{\Omega},\\
\tilde{q}(\alpha,t) &= \tilde{p}\circ\tilde{X}(\alpha,t) \quad \alpha\in\tilde{\Omega}.
\end{aligned}$$

The system (7-10) in terms of $(\tilde{v},\tilde{q})$ reads

$$\begin{aligned}
\partial_t\tilde{v}_i - Q^2\circ\tilde{X}\tilde{\zeta}_{kj}\partial_k\left(\tilde{\zeta}_{lj}\partial_l\tilde{v}_i\right) &= -A_{ki}\circ\tilde{X}\tilde{\zeta}_{jk}\partial_j\tilde{q} \quad \text{in } \tilde{\Omega}_0, && (11)\\
Tr\left(\nabla\tilde{v}\tilde{\zeta}A\circ\tilde{X}\right) &= 0 \quad \text{in } \tilde{\Omega}_0 && (12)\\
\left(\tilde{q}\mathbb{I} - \left(\left(\nabla\tilde{v}\tilde{\zeta}A\circ\tilde{X}\right) + \left(\nabla\tilde{v}\tilde{\zeta}A\circ\tilde{X}\right)^*\right)\right) A^{-1}\circ\tilde{X}\nabla_J\tilde{X}\tilde{n}_0 &= 0 \quad \text{on } \partial\tilde{\Omega}_0, && (13)\\
\tilde{v}|_{t=0} \equiv \tilde{v}_0 &= \tilde{u}_0, && (14)
\end{aligned}$$

where

$$\tilde{\zeta} = \left(\nabla\tilde{X}\right)^{-1} \quad \left(\tilde{\zeta}^{-1}\right)_{ij} = \partial_j\tilde{X}^i, \quad \nabla_J\tilde{X} = -J\nabla\tilde{X}J$$

and

$$\begin{aligned}
\frac{d\tilde{X}(\alpha,t)}{dt} &= A\circ\tilde{X}(\alpha,t)\tilde{v}(\alpha,t) && (15)\\
\tilde{X}(\alpha,0) &= \alpha, \quad \alpha\in\tilde{\Omega}_0. && (16)
\end{aligned}$$

We will solve the system (11-14) by iteration towards a fixed point. We show first the linear system that will be used for that purpose.

$$\partial_t \tilde v^{(n+1)} - Q^2 \Delta \tilde v^{(n+1)} = -A^* \nabla \tilde q^{(n+1)} + \tilde f^{(n)} \quad \text{in } \tilde\Omega_0, \tag{17}$$

$$Tr\left(\nabla \tilde v^{(n+1)} A\right) = \tilde g^{(n)} \quad \text{in } \tilde\Omega_0 \tag{18}$$

$$\left(\tilde q^{(n+1)}\mathbb{I} - \left(\left(\nabla \tilde v^{(n+1)} A\right) + \left(\nabla \tilde v^{(n+1)} A\right)^*\right)\right) A^{-1}\tilde n_0 = \tilde h^{(n)} \quad \text{on } \partial\tilde\Omega_0, \tag{19}$$

$$\tilde v|_{t=0} \equiv \tilde v_0 = \tilde u_0, \tag{20}$$

where

$$\tilde f_i^{(n)} = Q^2 \circ \tilde X^{(n)} \tilde\zeta_{kj}^{(n)} \partial_k \left(\tilde\zeta_{lj}^{(n)} \partial_l \tilde v_i^{(n)}\right) - Q^2 \Delta \tilde v_i^{(n)} - A_{ki} \circ \tilde X^{(n)} \tilde\zeta_{jk}^{(n)} \partial_j \tilde q^{(n)} + A_{ki} \partial_k \tilde q^{(n)} \tag{21}$$

$$\tilde g^{(n)} = - Tr\left(\nabla \tilde v^{(n)} \tilde\zeta^{(n)} A \circ \tilde X^{(n)}\right) + Tr\left(\nabla \tilde v^{(n)} A\right) \tag{22}$$

$$\begin{aligned}\tilde h^{(n)} = & \left(\left(\nabla \tilde v^{(n)} \tilde\zeta^{(n)} A \circ \tilde X^{(n)}\right) + \left(\nabla \tilde v^{(n)} \tilde\zeta^{(n)} A \circ \tilde X^{(n)}\right)^*\right) A^{-1} \circ \tilde X^{(n)} \nabla_J \tilde X^{(n)} \tilde n_0 \\ & - \left(\left(\nabla \tilde v^{(n)} A\right) + \left(\nabla \tilde v^{(n)} A\right)^*\right) A^{-1}\tilde n_0 + \tilde q^{(n)} A^{-1}\tilde n_0 - \tilde q^{(n)} A^{-1} \circ \tilde X^{(n)} \nabla_J \tilde X^{(n)} \tilde n_0,\end{aligned} \tag{23}$$

$\tilde X^{(n+1)}$ satisfies

$$\tilde X^{(n+1)}(\alpha, t) = \alpha + \int_0^t A \circ \tilde X^{(n)}(\alpha,\tau) \tilde v^{(n)}(\alpha,\tau) d\tau, \tag{24}$$

and

$$\tilde\zeta^{(n)} = \left(\nabla \tilde X^{(n)}\right)^{-1} \qquad\qquad \nabla_J \tilde X^{(n)} = -J \nabla \tilde X^{(n)} J.$$

Formally, assuming convergence as $n \to \infty$, it is easy to check that in the limit we find a solution of the nonlinear system. In what follows, we will either remove the tilde from the notation or it will become clear from the context.

3 Definitions of the spaces and auxiliary lemmas

This section is devoted to present the main tools used for both the linear and the nonlinear case. We will also define all the spaces used for the construction of the solutions and its norms:

$$H^{ht,s}\left([0,T];\Omega_0\right) = L^2\left([0,T];\, H^s(\Omega_0)\right) \cap H^{\frac{s}{2}}\left([0,T];\, L^2(\Omega_0)\right),$$

$$H^{ht,s}_{pr}\left([0,T];\Omega_0\right) = \left\{ q \in L^\infty([0,T]; \dot H^1(\Omega_0) \,:\, \nabla q \in H^{ht,s-1}\left([0,T];\Omega_0\right),\, q \in H^{ht,s-\frac{1}{2}}\left([0,T];\partial\Omega_0\right)\right\},$$

$$\overline{H}^{ht,s}([0,T];\Omega_0) = L^2([0,T]; H^s(\Omega_0)) \cap H^{\frac{s+1}{2}}([0,T]; H^{-1}(\Omega_0)),$$

and

$$F^{s+1}\left([0,T];\, \Omega_0\right) = L^\infty_{1/4}([0,T]; H^{s+1}(\Omega_0)) \cap H^2\left([0,T]; H^\gamma(\Omega_0)\right), \quad s-1-\varepsilon < \gamma < s-1 \quad (\varepsilon > 0,\ \text{suff. small}),$$

with

$$||f||_{L^\infty_{1/4}} = \sup_{t\in[0,T]} t^{-1/4} |f(t)|,$$

Also we will use the following notation:

$$\begin{array}{ll} ||\cdot||_{H^{ht,s}([0,T],\Omega_0)} = ||\cdot||_{H^{ht,s}} & ||\cdot||_{\overline{H}^{ht,s}([0,T],\Omega_0)} = ||\cdot||_{\overline{H}^{ht,s}} \\ ||\cdot||_{H^{ht,s}([0,T],\partial\Omega_0)} = |\cdot|_{H^{ht,s}} & ||\cdot||_{F^{s+1}([0,T],\Omega_0)} = ||\cdot||_{F^{s+1}} \\ ||\cdot||_{H^r([0,T];\, H^s(\Omega_0))} = ||\cdot||_{H^r H^s} & ||\cdot||_{H^r([0,T];\, H^s(\partial\Omega_0))} = |\cdot|_{H^r H^s} \\ ||\cdot||_{L^\infty([0,T];H^s(\Omega_0))} = ||\cdot||_{L^\infty H^s} & ||\cdot||_{L^\infty_{1/4}([0,T];H^s(\Omega_0))} = ||\cdot||_{L^\infty_{1/4} H^s} \\ ||\cdot||_{H^{ht,\,s}_{pr}([0,T],\Omega_0)} = ||\cdot||_{H^{ht,\,s}_{pr}} & . \end{array}$$

Lemma 3.1 (Parabolic Trace) *1. Suppose* $\frac{1}{2} < r \leq 5$*. The mapping* $v \to \partial_n^j v$ *extends to a bounded operator* $H^{ht,r}([0,T];\Omega_0) \to H^{ht,r-j-\frac{1}{2}}([0,T];\partial\Omega_0)$*, where* j *is an integer with* $0 \leq j < r - \frac{1}{2}$*. The mapping* $v \to \partial_t^k v(x,0)$ *also extends to a bounded operator* $H^{ht,r} \to H^{r-2k-1}(\Omega_0)$ *if* k *is an integer with* $0 \leq k < \frac{1}{2}(r-1)$*.*

2. Suppose $\frac{3}{2} < r < 5$*,* $r \neq 3$*, and* $r - \frac{1}{2}$ *is not an integer. Let*

$$W^r = \prod_{0 \leq j < r - \frac{1}{2}} H^{ht,r-j-\frac{1}{2}}([0,T];\partial\Omega_0) \times \prod_{0 \leq k < \frac{r-1}{2}} H^{ht,r-2k-1}([0,T];\Omega_0),$$

and let W_0^r *be the subspace consisting of* $\{a_j, w_j\}$ *so that*

$$\partial_t^k a(x,0) = \partial_n^j w_k(x) \quad x \in \partial\Omega,$$

for $j + 2k < r - \frac{3}{2}$*. Then the traces of 1 form a bounded operator* $H^{ht,r} \to W_0^r$*, and this operator has a bounded right inverse.*

Proof: See [5, Lemma 2.1]. □

Lemma 3.2 *Let* B *a Hilbert space.*

1. *For* $s \geq 0$*, there is a bounded extension operator* $H^s((0,T);B) \to H^s((0,\infty);B)$*.*
2. *For* $0 \leq s \leq 2$*,* $s - \frac{1}{2}$ *not an integer, there is an extension operator from*

   $$\left\{ v \in H^s((0,T);B)\,;\, \partial_t^k v|_{t=0} = 0, 0 \leq k < s - \frac{1}{2} \right\} \to H^s((-\infty,\infty);B)$$

   with norm bounded uniformly if T *is bounded above.*
3. *Similar statements apply to the extension of* $H^{ht,2s}$*.*

Proof: The proof can be found in [5, p.365, Lemma 2.3]. In this paper the second statement asserts that the operator extends to $H^s((0,\infty);B)$ rather than $H^s((-\infty,\infty);B)$. But one can easily adapt the proof to the case $H^s((-\infty,\infty);B)$. □

Lemma 3.3 *Suppose* $0 \leq r \leq 4$*.*

1. *The identity extends to a bounded operator*

   $$H^{ht,s} \to H^p H^{r-2p},$$

   $p \leq \frac{r}{2}$*.*
2. *If* r *is not an odd integer, the restriction of this operator to the subspace with* $\partial_t^k v|_{t=0}$*,* $0 \leq k < \frac{r-1}{2}$ *is bounded uniformly if* T *is bounded above.*

Proof: The proof can be found in [5, p.365, Lemma 2.3]. □

Lemma 3.4 *Let* $T_0 > 0$ *be arbitrary and* B *a Hilbert space, and choose* $T \leq T_0$*.*

1. *For* $v \in H^0((0,T);B)$*, we define* $V \in H^1((0,T);B)$ *by*

   $$V(t) = \int_0^t v(\tau)d\tau.$$

   Suppose $0 < s < \frac{1}{2}$ *and* $0 \leq \varepsilon < s$*. Then* $v \to V$ *is a bounded operator from* $H^s((0,T);B)$ *to* $H^{s+1-\varepsilon}((0,T);B)$*, and*

   $$||V||_{H^{s+1-\varepsilon}((0,T);B)} \leq C_0 T^\varepsilon ||v||_{H^s((0,T);B)}.$$

   where C_0 *is independent of* T *for* $0 < T \leq T_0$*.*
2. *For* $v \in H^s((0,T);B)$*, such that* $v(0) = 0$*,* $0 < s \leq 1$ *and* $0 \leq \varepsilon < s$*,*

   $$||V||_{H^{s+1-\varepsilon}((0,T);B)} \leq C_0 T^\varepsilon ||v||_{H^s((0,T);B)}.$$

   where C_0 *is independent of* T *for* $0 < T \leq T_0$*.*

Proof: The proof can be found in [5, p.365, Lemma 2.4]. The first statement is just the statement of Lemma 2.4 in [5]. The second one is an intermediate step in the proof of Lemma 2.4 in [5].

□

Lemma 3.5 *1. Suppose $r > 1$ and $r \geq s \geq 0$. If $v \in H^r(\Omega)$ and $w \in H^s(\Omega)$, then $vw \in H^s(\Omega)$, and*

$$||vw||_{H^s} \leq C||v||_{H^r}||w||_{H^s}.$$

2. If $v, w \in H^1(\Omega)$, then $vw \in H^0(\Omega)$, and

$$||vw||_{H^0} \leq C||v||_{H^1}||w||_{H^1}.$$

3. If $v \in H^r(\Omega)$, $r > 1$, and $w \in H^{-1}(\Omega)$, the dual space of $H^1(\Omega)$, then vw is defined in $H^{-1}(\Omega)$ and

$$||vw||_{H^{-1}} \leq ||v||_{H^r}||w||_{H^{-1}}.$$

4. If $v \in H^1(\Omega)$ and $v \in H^0(\Omega)$, then vw is defined in $H^{-1}(\Omega)$; and

$$||vw||_{H^{-1}} \leq C||v||_{H^1}||w||_{H^0}.$$

Proof: The proof can be found in [5, p. 366, Lemma 2.5]. Here we notice that we work in dimension 2 rather than in dimension 3 as in [5] and the second statement of lemma 3.5 can be improved in dimension 2.

□

Lemma 3.6 *If $v \in H^{\frac{1}{q}}$ and $w \in H^{\frac{1}{p}}$ with $\frac{1}{p} + \frac{1}{q} = 1$ and $1 < p < \infty$ then*

$$||vw||_{H^0} \leq C||v||_{H^{\frac{1}{q}}}||w||_{H^{\frac{1}{p}}}.$$

Proof: Applying Hölder inequality yields

$$||vw||_{H^0} \leq ||v||_{L^{2p}}||w||_{L^{2q}},$$

Now Gagliardo-Nirenberg inequality provides

$$||v||_{L^{2p}} \leq C||v||_{H^s}$$

for $\frac{1}{2p} = \frac{1}{2} - \frac{s}{2}$, that implies $s = \frac{1}{q}$. Proceeding similarly for w we have that

$$||vw||_{H^0} \leq C||v||_{H^{\frac{1}{q}}}||w||_{H^{\frac{1}{p}}}.$$

□

Lemma 3.7 *Suppose B, Y, Z are Hilbert spaces, and $M : B \times Y \to Z$ is a bounded, bilinear, multiplication operator. Suppose $v \in H^s((0,T); B)$ and $w \in H^s((0,T); Y)$, where $s > \frac{1}{2}$. If vw is defined by $M(v,w)$, then $vw \in H^s((0,T); Z)$ and*

1. $$||vw||_{H^s((0,T);Z)} \leq C||v||_{H^s((0,T);Y)}||w||_{H^s((0,T);B)}.$$

2. *Also, if $s \leq 2$ and v, w satisfy the additional condition $\partial_t^k v|_{t=0} = \partial_t^k w|_{t=0} = 0$, $0 \leq k < s - \frac{1}{2}$, and $s - \frac{1}{2}$ is not an integer, then the constant C in 1 can be chosen independent of T.*

3. *In addition, if $s \leq 2$ we have that, if w satisfies $\partial_t^k w|_{t=0} = 0$, $0 \leq k < s - \frac{1}{2}$, and $s - \frac{1}{2}$ is not an integer*

$$||vw||_{H^s((0,T);Z)} \leq C||v||_{H^s((0,T);Y)}||w||_{H^s((0,T);B)} + C||w||_{H^s((0,T);B)}$$

where the constants depend on $\partial_t^k v|_{t=0}$, but they do not depend on T.

Proof: The proof of 1 and 2 can be found in [5, p. 366, Lemma 2.6]. For part 3 we proceed as follows. Since k can take the values $k = 0, 1$ we have that

$$||vw||_{H^s((0,T);Z)} = \left\| \left(v - \sum_{j=0}^{k} t^j e^{-jt^2} \partial_t^j v|_{t=0} \right) w \right\|_{H^s((0,T);Z)} + \left\| \left(\sum_{j=0}^{k} t^j e^{-jt^2} \partial_t^j v|_{t=0} \right) w \right\|_{H^s((0,T);Z)},$$

and we can apply part 2 of the Lemma to yield

$$\begin{aligned} ||vw||_{H^s((0,T);Z)} \leq C & \left\| \left(v - \sum_{j=0}^{k} t^j e^{-jt^2} \partial_t^j v|_{t=0} \right) \right\|_{H^s((0,T);Y)} ||w||_{H^s((0,T);B)} \\ & + \left\| \left(\sum_{j=0}^{k} t^j e^{-jt^2} \partial_t^j v|_{t=0} \right) w \right\|_{H^s((0,T);Z)} \end{aligned}$$

□

Lemma 3.8 *For $2 < s < 2.5$ and ε, $\delta > 0$ small enough and $v \in F^{s+1}$ the following estimates hold:*

1. $||v||_{H^{\frac{s+1}{2}} H^{1-\varepsilon}} \leq C \, ||v||_{F^{s+1}}$.
2. $||v||_{H^{\frac{s+1}{2}+\varepsilon} H^{1+\delta}} \leq C \, ||v||_{F^{s+1}}$.
3. $||v||_{H^{\frac{s-1}{2}+\varepsilon} H^{2+\delta}} \leq C \, ||v||_{F^{s+1}}$.
4. $||v||_{H^{\frac{s}{2}-\frac{1}{4}+\varepsilon} H^{2+\delta}} \leq C \, ||v||_{F^{s+1}}$.
5. $||v||_{H^1 H^{s-1}} \leq C \, ||v||_{F^{s+1}}$.
6. $||v||_{H^{\frac{1}{2}+2\varepsilon} H^s} \leq C \, ||v||_{F^{s+1}}$.

For $v_{|t=0} = 0 = \partial_t v_{|t=0}$ the constants above are independent of T, if T is bounded above.

Proof: We will show the most singular cases, which are 2. and 4. The rest is proved in an analogous way.

Let us consider the space $L^2 H^{s+1} \cap H^2 H^\gamma \subset F^{s+1}$ and we will show the inequality on the first space. To do so, we extend the function to times in $(0, \infty)$ as in Lemma 3.2, part 1. It is easy to see that there also exists a linear and continuous extension for all t, taking

$$v(t) = 3v(-t) - 2v(-2t) \quad \forall t < 0,$$

and thus we can consider $t \in \mathbb{R}$. Since Ω_0 is a regular domain, we also consider the extension to the whole plane $\mathbb{R}^2$. This way, we can think of $v : \mathbb{R} \times \mathbb{R}^2 \to \mathbb{R}$.

Case 2: We first consider $\varepsilon = \delta = 0$. Using the Fourier transform in $\mathbb{R} \times \mathbb{R}^2$, we have

$$\|v\|^2_{H^{\frac{s+1}{2}} H^1} = \int_{\mathbb{R}\times\mathbb{R}^2} (1 + |\tau|^{s+1})(1 + |\xi|^2) |\hat{v}(\tau, \xi)|^2 d\tau d\xi = \int_{\mathbb{R}\times\mathbb{R}^2} (1 + |\xi|^2 + |\tau|^{s+1} + |\tau|^{s+1}|\xi|^2) ||^2 d\tau d\xi. \tag{25}$$

We only need to bound the previous integral by

$$\|v\|^2_{L^2 H^{s+1}} + \|v\|_{H^2 H^\gamma} = \int_{\mathbb{R}\times\mathbb{R}^2} (2 + |\xi|^{2(s+1)} + |\tau|^4 + |\xi|^{2\gamma} + |\tau|^4 |\xi|^{2\gamma}) |\hat{v}(\tau, \xi)|^2 d\tau d\xi.$$

We are only left to bound the integral with $|\tau|^{s+1}|\xi|^2$. Using Young's inequality:

$$|\tau|^{s+1} |\xi|^{2\lambda} |\xi|^{2(1-\lambda)} \leq C(|\xi|^{2\lambda p} + |\tau|^{(s+1)q} |\xi|^{2(1-\lambda)q}),$$

where $p^{-1} + q^{-1} = 1$. Taking $q = \frac{4}{s+1}, p = \frac{4}{3-s}, \lambda = \frac{(s+1)(3-s)}{4}$, we are only left to check that

$$(1-\lambda)q < \gamma < s-1,$$

but elementary formulas show it as long as $s > 1$. It is easy to see that in the previous calculation there is room for the case $\varepsilon, \delta > 0$ as long as ε and δ are sufficiently small.

Case 4:

Proceeding similarly to Case 2, it is enough to show that

$$|\tau|^{s-\frac{1}{2}}|\xi|^4 \leq C(1+|\tau|^4+|\xi|^{2(s+1)}+|\tau|^4|\xi|^{2\gamma}).$$

Using Young

$$|\tau|^{s-\frac{1}{2}}|\xi|^{4\lambda}|\xi|^{4(1-\lambda)} \leq C(|\xi|^{4\lambda p}+|\tau|^{(s-\frac{1}{2})q}|\xi|^{4(1-\lambda)q}).$$

Taking $q=\frac{4}{s-1/2}, p=\frac{4}{9/2-s}, \lambda=\frac{(s+1)(9-2s)}{16}$, we are only left to check that

$$2(1-\lambda)q < \gamma < s-1,$$

which is true for $s > \frac{3}{2}$. Similarly, there is room for the case $\varepsilon, \delta > 0$ and sufficiently small. Finally, the boundedness with respect to T is shown if one considers the extension given by Lemma 3.2, part 2. □

Lemma 3.9 *Let $G \in C^\infty(\overline{\Omega})$ and $v \in X$, where X is either H^pH^q or $L^\infty H^p$. Then*

$$\|G\circ v\|_X \leq C\|v\|_X, \quad \textit{and } \|G\circ u - G\circ v\|_X \leq C\|u-v\|_X.$$

Proof:

The proof follows easily since the most singular terms of $\partial_x^k\partial_t^l(G\circ v)$ are $G'(v)\partial_x^k\partial_t^l(v)$ and they can be bounded respectively in L^2 or L^∞ by $C\|\partial_x^k\partial_t^l v\|_{L^2}$ or $C\|\partial_x^k\partial_t^l v\|_{L^\infty}$. The rest of the (less singular) terms are bounded by interpolation.

The second inequality follows straightaway from the Fundamental Theorem of Calculus and interpolation. □

4 Solving the linear equation

4.1 Case $g = h = v_0 = 0$

We would like to solve the following system:

$$\begin{aligned} v_t - Q^2\Delta v + A^*\nabla q &= f && (26)\\ Tr(\nabla v A) &= 0 && (27)\\ (qI - ((\nabla v A) + (\nabla v A)^*)B_1 n &= 0 && (28)\\ v|_{t=0} &= 0, && (29) \end{aligned}$$

where

$$B_1 = -JA^{-1}J = \frac{1}{Q^2}A^*$$

and

$$A_{ij} = \partial_j P^i \circ P^{-1}.$$

Our first purpose will be to obtain a weak formulation of this system. In order to do it we will use the following identities

$$AA^* = Q^2 I$$

$$\Delta v = \operatorname{div}\left(AA^*\frac{1}{Q^2}\nabla v\right)$$

and also

$$Q^2\partial_l\left(\frac{1}{Q^2}A_{lj}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)\right) = Q^2\Delta v$$

Using this relation it is easy to arrive to the following identity:

$$\begin{aligned}
&\int_\Omega \partial_l\left(\frac{1}{Q^2}A_{lj}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)\right)\phi^i dx \\
&= -\int_\Omega \frac{1}{Q^2}A_{lj}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)\partial_l\phi^i dx + \int_{\partial\Omega} n^l A_{lj}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)\phi^i\frac{1}{Q^2}d\sigma \\
&= -\int_\Omega \frac{1}{Q^2}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)A_{lj}\partial_l\phi^i dx + \int_{\partial\Omega} n^l A_{lj}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)\phi^i\frac{1}{Q^2}d\sigma \\
&= -\frac{1}{2}\int_\Omega \frac{1}{Q^2}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)(A_{lj}\partial_l\phi^i + A_{li}\partial_l\phi^j)dx + \int_{\partial\Omega} n^l A_{lj}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)\phi^i\frac{1}{Q^2}d\sigma
\end{aligned}$$

We also have that

$$\int_\Omega A_{ki}\partial_k p\phi^i\frac{1}{Q^2}dx = -\int_\Omega p\partial_k\left(A_{ki}\phi^i\frac{1}{Q^2}\right)dx + \int_{\partial\Omega} pA_{ki}n_k\phi^i\frac{d\sigma}{Q^2}$$

The following identities hold:

$$\partial_k\left(A_{ki}\phi^i\frac{1}{Q^2}\right) = A_{ki}\partial_k\phi^i\frac{1}{Q^2}, \quad \partial_k\left(\frac{A_{ki}}{Q^2}\right) = 0$$

The energy identity reads:

$$\begin{aligned}
&\frac{1}{2}\int_\Omega \frac{1}{Q^2}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)(A_{lj}\partial_l\phi^i + A_{li}\partial_l\phi^j)dx - \int_\Omega p\partial_k\left(A_{ki}\frac{1}{Q^2}\phi^i\right)dx \\
&+ \int_{\partial\Omega} n^l A_{lj}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)\phi^i\frac{1}{Q^2}d\sigma + \int_{\partial\Omega}\phi^i pA_{ki}\tilde{n}_k\frac{d\sigma}{Q^2} = \int_\Omega f\phi\frac{1}{Q^2}dx
\end{aligned}$$

and therefore

$$\begin{aligned}
&\frac{1}{2}\int_\Omega \frac{1}{Q^2}(A_{kj}\partial_k v^i + A_{ki}\partial_k v^j)(A_{lj}\partial_l\phi^i + A_{li}\partial_l\phi^j)dx - \int_\Omega p\partial_k\left(A_{ki}\frac{1}{Q^2}\phi^i\right)dx \\
&= -\int_{\partial\Omega}(p\delta^{ij} - (A_{kj}\partial_k v^i + A_{ki}\partial k v^j))A_{lj}n^l\phi^i\frac{d\sigma}{Q^2} + \int_\Omega f\phi\frac{1}{Q^2}dx,
\end{aligned}$$

where

$$\partial_k\left(A_{ki}\frac{1}{Q^2}\phi^i\right) = \frac{A_{ki}}{Q^2}\partial_k\phi^i = \frac{1}{Q^2}Tr(\nabla\phi A)$$

that we finally write as follows

$$\begin{aligned}
&\frac{1}{2}\int_\Omega Tr\left(\left(\nabla vA + A^*\nabla v^*\right)\left(\nabla\phi A + A^*\nabla\phi^*\right)\right)\frac{1}{Q^2}dx - \int_\Omega p\,Tr(\nabla\phi A)\frac{1}{Q^2}dx \\
&= \int_\Omega f\cdot\phi\frac{dx}{Q^2} - \int_{\partial\Omega}\left(q\mathbb{I} - \left(\nabla vA + A^*\nabla v^*\right)\right)A^*n\frac{1}{Q^2}d\sigma.
\end{aligned} \tag{30}$$

Therefore (30) is the weak formulation of our system.

Next we will define a kind of Leray projector. Let H^0_d the subspace of H^0_w formed by vectors $A^*\nabla\phi$ such that $\phi \in H^1_0$. Let H^0_σ the orthogonal complement of H^0_w with the following scalar product:

$$(f,g)_{H^0_w} = \int_\Omega fg\frac{1}{Q^2}dx.$$

Then is easy to check that if $v \in H^1 \cap H^0_\sigma$, v must satisfy

$$Tr(\nabla v A) = 0.$$

For $v \in L^2$ we define Rv to be the projection of v onto H^0_σ . For $v \in H^1$ we have that

$$Rv = v - A^*\nabla\psi, \tag{31}$$

where

$$\begin{aligned} Q^2\Delta\psi = Tr(\nabla v A) \quad &\text{in } \Omega_0 \\ \psi = 0 \quad &\text{on } \partial\Omega_0, \end{aligned} \tag{32}$$

The next lemma will deal with some properties about this projector R.

Lemma 4.1 *Let $0 \le s \le 4$. We have:*

i) R is a bounded operator on H^s.

ii) R is a bounded operator on $H^{ht,s}$, with norm bounded uniformly if T is bounded above.

iii) If $\phi \in H^{s+1}$, then $R(A^*\nabla\phi) = A^*\nabla\phi_1$, with $Q^2\Delta\phi_1 = 0$ in Ω_0, $\phi_1 = \phi$ on $\partial\Omega_0$.

Proof:

i) For $v \in H^s, 0 \le s \le 4$, it is easy to see that $Tr(\nabla v A) \in H^{s-1}$. Therefore the solution of the system (32) satisfies:

$$\|\psi\|_{H^{s+1}} \le C \left\| \frac{Tr(\nabla v A)}{Q^2} \right\|_{H^{s-1}} \le \|v\|_{H^s},$$

by elliptic theory, since both A and Q^2 are regular. The identity (31) provides

$$\|Rv\|_{H^s} \le \|v\|_{H^s} + \|\nabla\psi\|_{H^s} \le C\|v\|_{H^s}.$$

ii) It is easy to check that ∂_t commutes with R, since

$$\partial^k_t Rv = \partial^k_t v - A^*\nabla\partial^k_t\psi = R\partial^k_t v.$$

This proves the result for an integer number of derivatives. By interpolation we get the result for fractional derivatives.

iii) For $\phi \in H^{s+1}$, if $v = A^*\nabla\phi$, then

$$R(A^*\nabla\phi) = A^*\nabla\phi - A^*\nabla\psi = A^*\nabla(\phi - \psi).$$

Thus, we have that

$$0 = Tr(\nabla(R(A^*\nabla\phi))A) = Q^2\Delta(\phi - \psi), \quad \psi|_{\partial\Omega_0} = 0,$$

which implies that $\phi - \psi|_{\partial\Omega_0} = \phi$ and we can take $\phi_1 = \phi - \psi$.

□

Once we have obtained the energy identity (30) and Lemma 4.1 we pass to announce the main theorem of this section:

Theorem 4.2 *Let* $f \in H^{ht,s-1}$, $2 < s \leq 3$ *such that* $Rf(0) = 0$. *Then*

$$\|v\|_{H^{ht,s+1}} + \|\nabla q\|_{H^{ht,s-1}} + \|q\|_{H^{ht,s-\frac{1}{2}}} \leq C\|f\|_{H^{ht,s-1}}.$$

The constant is independent of T.

The rest of this subsection is devoted to prove this theorem.

First of all we will modify the equation by considering the new variables

$$u = e^{-t}v; \quad q = e^{-t}p; \quad \overline{f} = e^{-t}f$$

We should remark that $f \in H^{ht,s-1} \Leftrightarrow \overline{f} \in H^{ht,s-1}$. Then, the equation reads

$$\begin{aligned}\partial_t u &= -e^{-t}v + e^{-t}v_t = -u + e^{-t}(Q^2\Delta v + Q^2\nabla q + f)\\ &= -u + Q^2\Delta u - A^*\nabla q + \overline{f}.\end{aligned}$$

We will solve therefore

$$\partial_t u + u - Q^2\Delta u + A^*\nabla q = \overline{f}.$$

Let's start projecting onto H^0_σ to obtain

$$\partial_t u + u - Q^2\Delta u + A^*\nabla q_1 = R\overline{f},$$

since $Tr(\nabla(Q^2\Delta\tilde{v})A) = 0$ and therefore $R(Q^2\Delta u) = Q^2\Delta u$, where $A^*\nabla q_1 = RA^*\nabla q$.

We now introduce the operator

$$S_A : V^{s+1}(\Omega) \to RH^{s-1}$$

defined via:

$$S_A u = -Q^2\Delta u + u + A^*\nabla q_1, \quad A^*\nabla q_1 \equiv RA^*\nabla q,$$

where $V^r(\Omega) = \{u \in RH^r, A^*t_0\left(\nabla uA + A^*\nabla u^*\right)A^*n_0 = 0\}$ and $RH^{s-1} = \{Rf, f \in H^{s-1}\}$.

The following lemma deals with the invertibility of S_A.

Lemma 4.3 S_A *has a bounded inverse for* $1 \leq s \leq 3$, *and*

$$\|S_A^{-1}f\|_{H^{s+1}} \leq C\|f\|_{H^{s-1}}.$$

Proof: Let $f \in RH^{s-1}$. We will show that there exists $u \in V^{s+1}$ such that $S_A u = f$, i.e. $u - Q^2\Delta u + A^*\nabla q_1 = Rf$. (we will keep in the notation Rf instead of f, although $Rf = f$, to keep in mind this fact). Using the energy identity (30), we observe that

$$(u,\phi) + \langle u,\phi\rangle = -\int_{\partial\Omega} h\cdot\phi\frac{d\sigma}{Q^2} + \int_\Omega R\overline{f}\phi\frac{dx}{Q^2} + \int_\Omega q_1 Tr(\nabla\phi A)\frac{d\tilde{x}}{Q^2}$$

is the weak formulation of

$$\begin{aligned}u - Q^2\Delta u - A^*\nabla q_1 &= R\overline{f}\\ \left(q_1 - (\nabla uA + A^*\nabla u^*)\right)A^*n_0 &= h\end{aligned}$$

If we look for $h = 0$ we then have to solve

$$(u,\phi) + \langle u,\phi\rangle = \int_\Omega R\overline{f}\phi\frac{dx}{Q^2} + \int_\Omega q_1 Tr(\nabla\phi A)\frac{d\tilde{x}}{Q^2}, \tag{33}$$

where

$$(u,\phi) = \int u\phi\frac{dx}{Q^2} \tag{34}$$

$$\langle u,\phi\rangle = \int_\Omega Tr((\nabla u A + A^*\nabla u^*)(\nabla\phi A + A^*\nabla\phi^*))\frac{dx}{Q^2} \tag{35}$$

We will find a solution to this equation in RH^1. By this we mean that there exists $u \in RH^1$ such that (33) holds for all $\phi \in RH^1$. We notice that the last term vanishes since $\phi \in RH^1$, and henceforth, in RH^1, the equation (33) is equivalent to

$$(u,\phi) + \langle u,\phi\rangle = \int_\Omega R\overline{f}\phi\frac{dx}{Q^2}. \tag{36}$$

for all $\phi \in RH^1$.

Via [29, Corollary 4.7],

$$(u,\phi) + \langle u,\phi\rangle \geq C||u||_{H^1}$$

and therefore it is a coercive bilinear form. That implies Lax-Milgram's Theorem can be applied to obtain a solution $u \in RH^1$ of (36).

The next step is to show that there exists $p \in L^2$ such that

$$(u,\phi) + \langle u,\phi\rangle = \int_\Omega Rf\phi + \int_\Omega p\, Tr(\nabla\phi A) \quad \forall\phi \in H^1.$$

To do this, we decompose $\phi = R\phi + A^*\nabla\pi_\phi$, and then

$$\begin{aligned}(u,\phi) + \langle u,\phi\rangle &= (u,R\phi) + \langle u,R\phi\rangle + \langle u,A^*\nabla\pi_\phi\rangle + \underbrace{(u,A^*\nabla\pi_\phi)}_{0}\\ &= (Rf,R\phi) + \langle u,A^*\nabla\pi_\phi\rangle = (Rf,\phi) + \langle u,A^*\nabla\pi_\phi\rangle.\end{aligned}$$

Therefore we have to show that there exists $p \in L^2$ such that

$$(p, Tr(\nabla\phi A)) = \langle u, A^*\nabla\pi_\phi\rangle \quad \forall\phi \in H^1.$$

Let us assume that u is smooth and suppose we take p satisfying:

$$\begin{aligned}Q^2\Delta p &= 0\\ p|_{\partial\Omega} &= (A^*n, ((\nabla uA) + (\nabla uA)^*)A^*n)\\ &= Q^2(n(A\nabla u + \nabla u^* A^*)n).\end{aligned}$$

Then, on the one hand

$$\begin{aligned}\int pTr(\nabla\phi A)\frac{dx}{Q^2} &= \int p\Delta\pi_\phi dx = \int p\mathrm{div}\left(\frac{AA^*}{Q^2}\nabla\pi_\phi\right)dx\\ &= -\int_\Omega \nabla pAA^*\nabla\pi_\phi\frac{dx}{Q^2} + \int_{\partial\Omega} pnAA^*\nabla\pi_\phi\frac{dx}{Q^2}\\ &= -\underbrace{\int_\Omega \nabla pQ^2\nabla\pi_\phi\frac{dx}{Q^2}}_{0} + \int_{\partial\Omega} p(A^*n)\cdot A^*\nabla\pi_\phi\frac{dx}{Q^2}.\end{aligned} \tag{37}$$

On the other hand

$$\langle u, \nabla\pi_\phi\rangle = \frac{1}{2}\int Tr((\nabla uA + A^*\nabla u^*)(\nabla(A^*\nabla\pi_\phi)A + A^*\nabla(A^*\nabla\phi\pi_\phi)))\frac{dx}{Q^2},$$

where

$$\begin{aligned}(\nabla(A^*\nabla\pi_\phi)A)_{ij} &= \partial_l(A^*\nabla\pi_\phi)^i A_{lj} = \partial_l(A_{ki}\partial_k\pi_\phi)A_{lj}\\ &= A_{lj}\partial_l A_{ki}\partial_k\pi_\phi + A_{ki}A_{lj}\partial^2_{lk}\pi_\phi\end{aligned}$$

$$\begin{aligned}\langle u, \nabla\pi_\phi\rangle &= \int(\partial_l u^i A_{lj} + \partial_l u^j A_{li})A_{mj}\partial_m(\partial_k\pi\phi A_{ki})\frac{dx}{Q^2}\\ &= -\int\partial_m\left(\left(A_{mj}\partial_k u^i A_{lj} + A_{mj}\partial_l u^j A_{li}\right)\frac{1}{Q^2}\right)\partial_k\pi\phi A_{ki}dx\\ &+ \int_{\partial\Omega}(n_0)_m\left(A_{mj}\partial_k u^i A_{lj} + A_{mj}\partial_l u^j A_{li}\right)\partial_k\pi\phi A_{ki}\frac{dx}{Q^2}\\ &= -\int\partial_m\left(\left(\delta^{ml}\partial_l u^i + A_{mj}\partial_l u^j A_{li}\right)\frac{1}{Q^2}\right)\partial_k\pi\phi A_{ki}dx\\ &+ \int_{\partial\Omega}A^*n_0\cdot(\nabla uA + A^*\nabla u)A^*\nabla\pi\phi\frac{dx}{Q^2}.\end{aligned}$$

The first term is zero because of the orthogonality and because of the condition $Tr(\nabla uA) = 0$. We do the calculations for the second term:

$$\int_{\partial\Omega}A^*n_0(\nabla uA + A^*\nabla u^*)A^*\nabla\pi_\phi\frac{d\sigma}{Q^2} = \int n_0(\nabla uA + A^*\nabla u^*)\nabla\pi_\phi = \int n_0(\nabla uA + A^*\nabla u^*)n_0\partial_n\pi_\phi,$$

since $t_0\cdot\nabla\pi_\phi = 0$. Comparing with (37) we have that

$$(p, Tr(\nabla A)) = \langle u, A^*\nabla\pi\phi\rangle, \quad \forall\phi\in H^1.$$

for smooth u. Let $u\in H^1$ and $\{u_m\}_{m=1}^\infty$ such that $u_m\in H^\infty$, and $tr(\nabla u_m A) = 0$ for every m, and $u_m\to u$ strongly in H^1 (for example we extend u by zero to $\mathbb{R}^2$, we make the convolution with a mollifier $\rho_{\frac{1}{m}}$ and finally we project onto H_σ). Let p_m be given by

$$\begin{aligned}\Delta p_m &= 0\\ p_m|_{\partial\Omega} &= A^{-1}n_0(\nabla u_m A + A^*\nabla(u_m)^*)A^{-1}n_0.\end{aligned}$$

Then, we have that

$$(p_m, Tr(\nabla\phi A)) = \langle u_m, A^*\nabla\pi_\phi\rangle \quad \forall\phi\in H^1.$$

In particular, we take ϕ_m such that $Tr(\nabla\phi_m A) = p_m$. This implies that $Q^2\Delta\pi_{\phi_m} = p_m$, $\pi_{\phi_m}|_{\partial\Omega} = 0$. Showing the existence of such ϕ_m is trivial since one can choose $\phi_m = A^*\nabla\psi$, with $Q^2\Delta\psi = p_m$. Then, we can bound the L^2 norm of p_m in the following way:

$$\|p_m\|^2_{L^2} \le \langle u_m, A^*\nabla\pi_{\phi_m}\rangle \le \|u_m\|_{H^1}\|\pi_{\phi_m}\|_{H^2} \le C\|u\|_{H^1}\|p_m\|_{L^2},$$

which shows that p_m is bounded in L^2. Therefore there exists a subsequence p_{m_i} which converges weakly to a function p in L^2 and

$$(p, Tr(\nabla\phi A)) = \langle u, \nabla\pi_\phi\rangle \quad \forall\phi\in H^1.$$

We have shown that there exist $(u, p)\in H^1\times L^2$ such that

$$(u, \phi) + \langle u, \phi\rangle = (Rf, \phi) + \int ptr(\nabla\phi A)\frac{dx}{Q^2} \quad \forall\phi\in H^1.$$

Indeed, $u \in H^2, p \in H^1$. We now show that improvement on the regularity.

For every $\Omega^\flat \Subset \Omega$, it is easy to obtain the interior regularity estimate $\|u\|_{H^2(\Omega^\flat)} \leq C, \|p\|_{H^1(\Omega^\flat)} \leq C$. We focus here on the boundary estimates. We perform the following change of coordinates in Ω', where Ω' is a tubular neighbourhood of $\partial\Omega$:

$$x(s, \lambda) = z(s) - \lambda z_s^\perp(s)$$

We would like to check that it is indeed a diffeomorphism. We have following quantities:

$$\begin{aligned}
|z_s|^2 &= 1 \\
x_s(s) &= z_s(s) - \lambda z_{ss}^\perp(s) \\
z_{ss}^\perp &= \langle z_{ss}^\perp, z_s(s)\rangle z_s(s) = -\kappa(s) z_s(s) \\
x_s(s) &= (1 + \lambda\kappa(s)) z_s(s) \\
x_\lambda &= z_s^\perp(s)
\end{aligned}$$

Computing more,

$$\det \begin{pmatrix} x_s^1 & x_s^2 \\ x_\lambda^1 & x_\lambda^2 \end{pmatrix} = x_s^1 x_\lambda^2 - x_s^2 x_\lambda^2 = x_s^\perp x_\lambda = 1 - \lambda\kappa(s)$$

This fixes the width of the tubular neighbourhood to be $\lambda_0 < \sup_s \frac{1}{\kappa(s)}$. Under these assumptions $x(s, \lambda)$ is a diffeomorphism. Fix $x_0 \in \partial\Omega$, and consider the following cutoff function ψ, defined by

$$\psi(x) = \left\{ \begin{array}{ll} 1 & B\left(x_0, \frac{\lambda_0}{2}\right) \cap \Omega' \\ 0 & \left(B\left(x_0, \frac{3\lambda_0}{4}\right) \cap \Omega'\right)^c \end{array} \right\}, \quad 0 \leq \psi(x) \leq 1, \quad \psi(x) \in C^\infty.$$

We extend ψ to Ω by zero. We define the set C as $C = x^{-1}(B(x_0, \lambda_0) \cap \Omega')$, and the set $C/2 = x^{-1}(B(x_0, \frac{\lambda_0}{2}) \cap \Omega')$.

The energy identity can be written down as:

$$\int_\Omega (\partial_l u^i A_{lj} + \partial_l u^j A_{li})(\partial_k \phi^i A_{kj} + \partial_k \phi^j A_{ki}) \frac{dx}{Q^2} + \int_\Omega u^i \phi^i \frac{dx}{Q^2} = \int_\Omega f^i \phi^i \frac{dx}{Q^2} + \int_\Omega p \partial_l \phi^i A_{li} \frac{dx}{Q^2}$$

We choose as test function $\phi = \varphi\psi$. Then:

$$\int_\Omega (\partial_l u^i A_{lj} + \partial_l u^j A_{li})(\partial_k(\varphi^i\psi) A_{kj} + \partial_k(\varphi^j\psi) A_{ki}) \frac{dx}{Q^2} + \int_\Omega u^i \phi^i \frac{dx}{Q^2} = \int_\Omega f^i \phi^i \frac{dx}{Q^2} + \int_\Omega p \partial_l \phi^i A_{li} \frac{dx}{Q^2}$$
$$M_1 + M_2 = M_3 + M_4$$

We start developing each of the terms one by one

$$\begin{aligned}
M_1 &= \int_{\Omega'} (\partial_l u^i A_{lj} + \partial_l u^j A_{li})(\partial_k(\varphi^i) A_{kj} + \partial_k(\varphi^j) A_{ki}) \psi \frac{dx}{Q^2} \\
&+ \int_{\Omega'} (\partial_l u^i A_{lj} + \partial_l u^j A_{li})(\varphi^i \partial_k \psi A_{kj} + \varphi^j \partial_k \psi A_{ki}) \frac{dx}{Q^2} \\
&= \int_{\Omega'} (\partial_l u^i A_{lj} + \partial_l u^j A_{li}) \psi (\partial_k(\varphi^i) A_{kj} + \partial_k(\varphi^j) A_{ki}) \frac{dx}{Q^2} \\
&+ \int_{\Omega'} (\partial_l u^i A_{lj} + \partial_l u^j A_{li}) \partial_k \psi (\varphi^i A_{kj} + \varphi^j A_{ki}) \frac{dx}{Q^2}
\end{aligned}$$

We now do the change of variables

$$\underline{dx} = \left(\frac{dx}{dsd\lambda}\right) dsd\lambda = \underbrace{(1+\kappa(s)\lambda)}_{J(s,\lambda)} dsd\lambda,$$

and define

$$\begin{aligned} &\overline{A}(s,\lambda) = A \circ x(s,\lambda), \quad \overline{Q}(s,\lambda) = Q \circ x(s,\lambda), \quad \overline{u}(s,\lambda) = u \circ x(s,\lambda), \quad \overline{p}(s,\lambda) = p \circ x(s,\lambda), \\ &\overline{\varphi}(s,\lambda) = \varphi \circ x(s,\lambda), \quad \overline{\psi}(s,\lambda) = \psi \circ x(s,\lambda), \quad \overline{f}(s,\lambda) = f \circ x(s,\lambda). \end{aligned}$$

We can compute the derivatives in the bar-coordinates

$$\begin{aligned} u(x) = \overline{u} \circ s(x), \quad \partial_l u^i = \partial_k \overline{u} \circ s(x) \partial_l s^k \\ \partial_l u^i \circ x(s,\lambda) = \partial_k \overline{u}(s,\lambda) \partial_l s^k \circ x(s,\lambda) \end{aligned}$$

Setting $x = x(s,\lambda), \quad x^j = x^j(s,\lambda)$, we obtain:

$$\begin{aligned} \partial_l x^j &= \partial_l x^j(s,\lambda) \frac{\partial s^l}{\partial x^i} \\ \delta^{ij} &= \partial_l x^j(s,\lambda) \frac{\partial s^l}{\partial x^i} \\ \delta^{ij} &= \partial_l x^j \circ s(x) \frac{\partial s^l}{\partial x^i}(x) \\ \delta^{ij} &= (\nabla x)_{jl} (\nabla s)_{li} \circ x(s,\lambda) = (\nabla x \nabla s)_{ji} \\ 1 &= \nabla x \nabla s \circ x(s,\lambda), \quad \nabla s \circ x(s,\lambda) = (\nabla x)^{-1} = \Sigma \end{aligned}$$

Plugging this result into the equation for the derivatives of u, we get

$$\begin{aligned} \partial_l u^i \circ x(s,\lambda) = \Sigma_{kl} \partial_k \overline{u}^i \\ \Sigma = \frac{1}{1+\kappa(s)\lambda} \begin{pmatrix} x_\lambda^2 & -x_\lambda^1 \\ -x_s^2 & x_s^1 \end{pmatrix} \end{aligned}$$

Plugging this expression into M_1, and letting $\Sigma\overline{A} = B$, we obtain

$$\begin{aligned} M_1 = &\int_C (\partial_l u^i \circ x(s,\lambda) \overline{A}_{lj} + \partial_l u^j \circ x(s,\lambda) \overline{A}_{li}) \psi (\partial_k \overline{\varphi}^i \circ x(s,\lambda) \overline{A}_{kj} + \partial_k \overline{\varphi}^j \circ x(s,\lambda) \overline{A}_{ki} \frac{J}{\overline{Q}^2} dsd\lambda \\ &+ \int_C \partial_a \overline{\psi} (\partial_m \overline{u}^i B_{mj} + \partial_m \overline{u}^j B_{mi})(\overline{\varphi}^i B_{aj} + \overline{\varphi}^j B_{ai}) \frac{J}{\overline{Q}^2} dsd\lambda \\ &= \int_C (\partial_m \overline{u}^i \Sigma_{ml} \overline{A}_{lj} + \partial_m \overline{u}^j \Sigma_{ml} \overline{A}_{li}) \overline{\psi} (\partial_a \overline{\varphi}^i \Sigma_{ak} \overline{A}_{kj} + \partial_a \overline{\varphi}^j \Sigma_{ak} \overline{A}_{ki}) \frac{J}{\overline{Q}^2} dsd\lambda \\ &+ \int_C \partial_a \overline{\psi} (\partial_m \overline{u}^i B_{mj} + \partial_m \overline{u}^j B_{mi})(\overline{\varphi}^i B_{aj} + \overline{\varphi}^j B_{ai}) \frac{J}{\overline{Q}^2} dsd\lambda \\ &= \int_C \overline{\psi} (\partial_m \overline{u}^i B_{mj} + \partial_m \overline{u}^j B_{mi})(\partial_a \overline{\varphi}^i B_{aj} + \partial_a \overline{\varphi}^j B_{ai}) \frac{J}{\overline{Q}^2} dsd\lambda \\ &+ \int_C \partial_a \overline{\psi} (\partial_m \overline{u}^i B_{mj} + \partial_m \overline{u}^j B_{mi})(\overline{\varphi}^i B_{aj} + \overline{\varphi}^j B_{ai}) \frac{J}{\overline{Q}^2} dsd\lambda \end{aligned}$$

Moreover,

$$
\begin{aligned}
M_2 &= \int_{\Omega'} u^i \varphi^i \psi \frac{dx}{Q^2} = \int_C \overline{u}^i \overline{\varphi}^i \overline{\psi} \frac{J}{\overline{Q}^2} ds d\lambda \\
M_3 &= \int_C \overline{f}^i \overline{\varphi}^i \overline{\psi} \frac{J}{\overline{Q}^2} ds d\lambda \\
M_4 &= \int_C \overline{p} \partial_a (\overline{\varphi}^i \overline{\psi}) B_{ai} \frac{J}{\overline{Q}^2} ds d\lambda,
\end{aligned}
$$

with

$$
M_1 + M_2 = M_3 + M_4 \Rightarrow M_1 = -M_2 + M_3 + M_4.
$$

We start computing M_1. We have that

$$
\begin{aligned}
2M_1 &= \int_C (\partial_l \overline{u}^i B_{lj} + \partial_l \overline{u}_j B_{li})(\partial_k \overline{\varphi}^i B_{kj} + \partial_k \overline{\varphi}^j B_{ki}) \overline{\psi} \frac{J}{\overline{Q}^2} \\
&+ \int_C (\partial_l \overline{u}^i B_{lj} + \partial_l \overline{u}_j B_{li})(\overline{\varphi}^i B_{kj} + \overline{\varphi}^j B_{ki}) \partial_k \overline{\psi} \frac{J}{\overline{Q}^2} \\
&= m + l_1.
\end{aligned}
$$

We will use the convention that the m terms will denote bad terms from now on. We further split m into four terms:

$$
\begin{aligned}
m &= \int_C \partial_l \overline{u}^i B_{lj} \partial_k \overline{\varphi}^i B_{kj} \overline{\psi} \frac{J}{\overline{Q}^2} + \int_C \partial_l \overline{u}^j B_{li} \partial_k \overline{\varphi}^i B_{kj} \overline{\psi} \frac{J}{\overline{Q}^2} \\
&+ \int_C \partial_l \overline{u}^i B_{lj} \partial_k \overline{\varphi}^j B_{ki} \overline{\psi} \frac{J}{\overline{Q}^2} + \int_C \partial_l \overline{u}^j B_{li} \partial_k \overline{\varphi}^j B_{ki} \overline{\psi} \frac{J}{\overline{Q}^2} \\
&= m_1 + m_2 + m_3 + m_4.
\end{aligned}
$$

If we are able to estimate any one of them we can estimate all of them.

We will use the following formulas related to finite differences:

$$
\begin{aligned}
D_s^h(fg) &= \frac{f(s+h)g(s+h) - f(s)g(s)}{h} = \frac{(f(s+h) - f(s))g(s+h)}{h} + f(s)\frac{g(s+h) - g(s)}{h} \\
&= g(s+h) D_s^h f + f(s) D_s^h g \\
D_s^{-h} D_s^h(fg) &= D_s^{-h}(D_s^h f g(s+h)) + D_s^{-h}(f(s) D_s^h g) \\
&= (D_s^{-h} D_s^h f) g(s+h) + D_s^h f D_s^{-h}(g(s+h)) + D_s^{-h} f D_s^h g(s-h) + f(s) D_s^{-h} D_s^h g \\
D_s^{-h}(g(s+h)) &= \frac{g(s) - g(s+h)}{-h} = D_s^h g \\
(D_s^h g)(s-h) &= \frac{g(s) - g(s-h)}{h} = D_s^{-h} g \\
\Rightarrow D_s^{-h} D_s^h(fg) &= (D_s^{-h} D_s^h f) g(s+h) + D_s^h f D_s^h g + D_s^{-h} f D_s^h g + f(s) D_s^{-h} D_s^h g
\end{aligned}
$$

We take $\overline{\varphi} = D_s^{-h} D_s^h \overline{u}$. Thus, we have that

$$
D_s^{-h} D_s^h(B_{lj} \partial_l \overline{u}^j) = 0 = (D_s^{-h} D_s^h B_{lj}) \partial_l \overline{u}^j (h+s) + D_s^h B_{lj} D_s^{-h} \partial_l \overline{u}^j + D_s^{-h} B_{lj} D_s^h \partial_l \overline{u}^j + B_{lj} \partial_l D_s^{-h} D_s^h \overline{u}^j.
$$

Therefore

$$
\begin{aligned}
B_{lj} \partial_l D_s^{-h} D_s^h \overline{u}^j &= -(D_s^{-h} D_s^h B_{lj}) \partial_l \overline{u}^j (h+s) - D_s^h B_{lj} D_s^{-h} \partial_l \overline{u}^j - D_s^{-h} B_{lj} D_s^h \partial_l \overline{u}^j \\
B_{lj} \partial_l D_s^{-h} D_s^h (\overline{u}^j \overline{\psi}) &= D_s^{-h} D_s^h (B_{lj} \partial_j (\overline{u}^j \overline{\psi})) - D_s^{-h} D_s^h B_{lj} \partial_l (\overline{u}^j (s+h) \overline{\psi}(s+h)) \\
&- D_s^h B_{lj} D_s^{-h} \partial_l (\overline{u}^j \overline{\psi}) - D_s^{-h} B_{lj} D_s^h (\partial_l (\overline{u}^j \overline{\psi})),
\end{aligned}
$$

and

$$B_{lj}\partial_j(\overline{\psi}\overline{u}^j) = B_{lj}\partial_j\overline{\psi}\overline{u}^j.$$

Expanding the calculations, we obtain

$$\begin{aligned}
B_{lj}\partial_l D_s^{-h}D_s^h(\overline{u}^j\overline{\psi}) &= D_s^{-h}D_s^h(B_{lj}\partial_j\overline{\psi}\overline{u}^j) - D_s^{-h}D_s^hB_{lj}\partial_l(\overline{u}^j(s+h)\overline{\psi}(s+h)) \\
&\quad - D_s^hB_{lj}D_{-s}^h\partial_l(\overline{u}^j\overline{\psi}) - D_s^{-h}B_{lj}D_s^h(\partial_l\overline{u}^j\overline{\psi})) \\
\overline{\psi}B_{lj}\partial_l D_s^{-h}D_s^h(\overline{u}^j\overline{\psi}) &= \overline{\psi}D_s^{-h}D_s^h(B_{lj}\partial_j\overline{\psi}\overline{u}^j) - \overline{\psi}D_s^{-h}D_s^hB_{lj}\partial_l(\overline{u}^j(s+h)\overline{\psi}(s+h)) \\
&\quad - \overline{\psi}D_s^hB_{lj}D_s^{-h}\partial_l(\overline{u}^j\overline{\psi}) - \overline{\psi}D_s^{-h}B_{lj}D_s^h(\partial_l(\overline{u}^j\overline{\psi})).
\end{aligned}$$

Finally, we get to

$$\begin{aligned}
\overline{\psi}D_s^{-h}D_s^h(B_{lj}\partial_j\overline{\psi}\overline{u}^j) &= D_s^{-h}D_s^h(B_{lj}\partial_j\overline{\psi}\overline{u}^j) - D_s^{-h}D_s^h(B_{lj}\partial_j\overline{\psi})(\overline{u}^j\overline{\psi})(s+h) \\
&\quad - D_s^{-h}B_{lj}\partial_j\overline{\psi}D_s^h(\overline{u}^j\overline{\psi}) - D_s^h(B_{lj}\partial_j\overline{\psi})D_s^{-h}(\overline{u}^j\overline{\psi}).
\end{aligned}$$

Therefore

$$\begin{aligned}
\overline{\psi}B_{lj}\partial_l D_s^{-h}D_s^h(\overline{u}^j\overline{\psi}) &= D_s^{-h}D_s^h(B_{lj}\partial_l\overline{\psi}\overline{u}^j) - \overline{\psi}D_s^hB_{lj}D_s^{-h}\partial_l(\overline{u}^j\overline{\psi}) - \overline{\psi}D_s^{-h}B_{lj}D_s^h\partial_l(\overline{u}^j\overline{\psi}) + \text{LOW} \\
&= B_{lj}\partial_l\overline{\psi}D_s^{-h}D_s^h(\overline{u}^j\overline{\psi}) - \overline{\psi}D_s^hB_{lj}D_s^{-h}\partial_l(\overline{u}^j\overline{\psi}) - \overline{\psi}D_s^{-h}B_{lj}D_s^h\partial_l(\overline{u}^j\overline{\psi}) + \text{LOW},
\end{aligned}$$

where we say that a term T is LOW is $\|T\|_{L^2} \leq C\|\overline{u}\|_{H^1}$. We also say that a term T is SAFE if, for any $\delta > 0$,

$$\|T\|_{L^2} \leq C_\delta + \delta(\|\nabla D_s^{-h}(\overline{u}\overline{\psi})\|_{L^2}^2 + \|\nabla D_s^h(\overline{u}\overline{\psi})\|_{L^2}^2),$$

where C_δ may depend on $\|\overline{u}\|_{H^1}, \|\overline{p}\|_{L^2}, \|\overline{f}\|_{L^2}$ and $\overline{\psi}, B, \overline{Q}$ or J. We now look at what the terms $M_1, \ldots, M_4$ look like. We have that

$$\begin{aligned}
M_2 &= \int_C \overline{\psi}\overline{u}^i D_s^{-h}D_s^h\overline{u}^i\frac{J}{\overline{Q}^2} = \int_C D_s^h(\overline{\psi}\overline{u}^i)D_s^h\overline{u}^i\frac{J}{Q^2} + \text{LOW}. \\
M_3 &= \int_C \overline{f}^i D_s^{-h}D_s^h\overline{u}^i\frac{J}{\overline{Q}^2} \leq C_\delta\|\overline{f}^i\|_{L^2}^2 + \delta\|D_s^{-h}D_s^h(\overline{u}\overline{\psi})\|_{L^2}^2 \\
M_4 &= \int_C \overline{p}D_{lj}\partial_j\overline{\psi}D_s^{-h}D_s^h\overline{u}^i \leq C_\delta\|\overline{p}\|_{L^2}^2 + \delta\|D_s^{-h}D_s^h(\overline{u}\overline{\psi})\|_{L^2}^2 \\
l_1 &= \int_C \partial_l\overline{u}^iB_{lj}\phi^j\partial_k\overline{\psi}B_{kj}\frac{J}{\overline{Q}^2} \leq C_\delta\|\overline{u}\|_{H^1}^2 + \delta\|D_s^{-h}D_s^h(\overline{u}\overline{\psi})\|_{L^2}^2 \\
M_1 &= \int_C \partial_l\overline{u}^iB_{lj}B_{kj}\partial_kD_s^{-h}D_s^h(\overline{u}^i\overline{\psi})\overline{\psi}\frac{J}{\overline{Q}^2} = \int_C D_s^h\left(\overline{\psi}\partial_l\overline{u}^iB_{lj}B_{kj}\frac{J}{\overline{Q}^2}\right)D_s^h(\partial_k(\overline{u}^i\overline{\psi})) \\
&= \int_C B_{lj}B_{kj}D_s^h(\overline{\psi}\partial_l\overline{u}^i)D_s^h\partial_k(\overline{u}^i\overline{\psi})\frac{J}{\overline{Q}^2} + \text{SAFE} \\
&= \int_C B_{lj}B_{kj}D_s^h(\partial_l\overline{\psi}\overline{u}^i)D_s^h\partial_k(\overline{u}^i\overline{\psi})\frac{J}{\overline{Q}^2} + \text{SAFE}.
\end{aligned}$$

We can then bound

$$\begin{aligned}
\int_C |\nabla D_s^h(\overline{\psi}\overline{u})|^2 &\leq M_1 \leq C_\delta + \delta\|\nabla D_s^{-h}(\overline{u}\overline{\psi})\|_{L^2}^2 + \delta\|\nabla D_s^h(\overline{u}\overline{\psi})\|_{L^2}^2 \\
\int_C |\nabla D_s^{-h}(\overline{\psi}\overline{u})|^2 &\leq M_1 \leq C_\delta + \delta\|\nabla D_s^{-h}(\overline{u}\overline{\psi})\|_{L^2}^2 + \delta\|\nabla D_s^h(\overline{u}\overline{\psi})\|_{L^2}^2.
\end{aligned}$$

Similar computations show that we can control $\|\nabla \partial_s u\|_{L^2}$. This implies that

$$\|\partial_s u\|_{H^1(C/2)} \leq C.$$

We proceed to control the pressure terms. From the energy identity:

$$\frac{1}{2}\int_C (\partial_l \overline{u}^i B_{lj} + \partial_l \overline{u}^j B_{li})(\partial_k \overline{\varphi}^i B_{kj} + \partial_k \overline{\varphi}^j B_{ki})\frac{J}{\overline{Q}^2} + \int_C (\partial_l \overline{u}^i B_{lj} + \partial_l \overline{u}^j B_{li})(\overline{\varphi}^i B_{kj} + \overline{\varphi}^j B_{ki})\partial_k \overline{\psi}\frac{J}{\overline{Q}^2} + \int_C \overline{u}^i \overline{\partial}^i \psi \frac{J}{\overline{Q}^2}$$
$$= \int_C \overline{f}^i \overline{\varphi}^i \overline{\psi}\frac{J}{\overline{Q}^2} + \int_C \overline{p} Tr(\nabla \overline{\varphi} B)\overline{\psi}\frac{J}{\overline{Q}^2} + \overline{p} B \nabla \overline{\psi}\overline{\varphi}\frac{J}{\overline{Q}^2}$$
$$P_1 + P_2 = P_3 + P_4 + P_5$$

We choose $\overline{\varphi} = B^* \nabla \overline{\Pi}$, with

$$\frac{J}{\overline{Q}^2} D_s^{-h} D_s^h (\overline{p}\overline{\psi}) = \operatorname{div}(J \Sigma \Sigma^* \nabla \overline{\Pi}).$$

Then:

$$\begin{aligned} P_4 = \int_C \overline{p} Tr(\nabla \overline{\varphi} B)\overline{\psi}\frac{J}{\overline{Q}^2} &= \int_C \overline{\psi}\overline{p} D_s^{-h} D_s^h (\overline{p}\overline{\psi})\frac{J}{\overline{Q}^2} \\ &= -\int_C (D_s^h(\overline{\psi}\overline{p}))^2 \frac{J}{\overline{Q}^2} + \int_C \left(D_s^h \left(\frac{J}{\overline{Q}^2}\overline{\psi}\overline{p}\right) - \frac{J}{\overline{Q}^2} D_s^h(\overline{\psi}\overline{p})\right) \\ &= m_5 + l_1, \end{aligned}$$

where

$$|l_1| \leq c\|p\|_{L^2}.$$

So, if we control m_5 we can control $\|\partial_s \overline{p}\|_{L^2(C/2)}$. It is not hard to see that

$$|P_1| + |P_2| + |P_3| + |P_5| \leq C_\delta(\|\overline{u}\|_{L^2}^2 + \|\nabla \partial_s \overline{u}\|_{L^2}^2 + \|\nabla \overline{u}\|_{L^2}^2 + \|\overline{p}\|_{L^2}^2) + \delta\|D_s^h(\overline{p}\overline{\psi})\|_{L^2}^2$$

which implies that $\|D_s^h(\overline{p}\overline{\psi})\|_{L^2} \leq C$ independent of h. Because of the interior regularity estimate we have that the solution is strong in the interior of Ω and we can write

$$\frac{\overline{Q}^2}{J}\operatorname{div}\left(\frac{1}{J}\begin{pmatrix} 1 & 0 \\ 0 & J^2 \end{pmatrix} \nabla \overline{u}^i\right) + \overline{u}^i + (B^* \nabla \overline{p})^i = \overline{f}^i, \quad Tr(\nabla \overline{u} B) = 0$$

where we recall that $J = (1 + \lambda\kappa(s))$ and that we have used

$$\Delta u \circ x(s,\lambda) = \frac{1}{J(s,\lambda)}\operatorname{div}\left(J(s,\lambda)\Sigma\Sigma^* \nabla \overline{u}\right)(s,\lambda)$$

$$\Sigma\Sigma^* = \frac{1}{(1+\kappa(s)\lambda)^2}\begin{pmatrix} x_\lambda^2 & -x_\lambda^1 \\ -x_s^2 & x_s^1 \end{pmatrix}\begin{pmatrix} x_\lambda^2 & -x_s^2 \\ -x_\lambda^1 & x_s^1 \end{pmatrix} = \frac{1}{(1+\kappa(s)\lambda)^2}\begin{pmatrix} |x_\lambda|^2 & -x_\lambda \cdot x_s \\ -x_\lambda \cdot x_s & |x_s|^2 \end{pmatrix}$$

This implies that

$$\frac{1}{J(s,\lambda)}\operatorname{div}\left(\frac{1}{(1+\kappa(s)\lambda)^2}\begin{pmatrix} |x_\lambda|^2 & -x_\lambda \cdot x_s \\ -x_\lambda \cdot x_s & |x_s|^2 \end{pmatrix}\nabla \overline{u}^i\right) = \frac{1}{J(s,\lambda)}\operatorname{div}\left(\frac{1}{(1+\kappa(s)\lambda)^2}\begin{pmatrix} 1 & 0 \\ 0 & (1+\kappa(s)\lambda)^2 \end{pmatrix}\nabla \overline{u}^i\right).$$

Let us define

$$\begin{aligned}
\beta &= \begin{pmatrix} 1/J & 0 \\ 0 & J \end{pmatrix} \\
\operatorname{div}(\beta \nabla \overline{u}^i) &= \partial_k(\beta_{kl}\partial_l \overline{u}^i) = \underbrace{\partial_k B_{kl}}_{\Gamma_l} \partial_l \overline{u}^i + \beta_{kl}\partial_k \partial_l \overline{u}^i \\
&= \Gamma \nabla \overline{u}^i + \frac{1}{J}\partial_s^2 \overline{u}^i + J \partial_\lambda^2 \overline{u}^i .
\end{aligned}$$

And then:

$$\begin{aligned}
\frac{\overline{Q}^2}{J}\partial_s^2 \overline{u}^i + \overline{Q}^2 \partial_\lambda^2 \overline{u}^i + \Gamma \nabla \overline{u}^i + B_{li}\partial_l \overline{p} &= \overline{f}^i \\
\Rightarrow \overline{Q}^2 \partial_\lambda^2 \overline{u}^i + B_{li}\partial_l \overline{p} &= -\frac{\overline{Q}^2}{J}\partial_s^2 \overline{u}^i - \Gamma \nabla \overline{u}^i + \overline{f}^i = g^i,
\end{aligned}$$

and we know that $g^i \in L^2\left(C/2\right)$. We also have that

$$\begin{aligned}
(\partial_l \overline{u}^i B_{li}) &= 0 \\
\partial_s \overline{u}^1 B_{11} + \partial_\lambda \overline{u}^i B_{21} + \partial_s \overline{u}^2 B_{12} + \partial_\lambda \overline{u}^2 B_{22} &= 0
\end{aligned}$$

Thus

$$\partial_\lambda^2 \overline{u}^1 B_{21} + \partial_\lambda^2 \overline{u}^2 B_{22} = g^3, \text{ where } g^3 \in L^2$$

Plus

$$\begin{aligned}
\overline{Q}^2 \partial_\lambda^2 \overline{u}^1 + B_{11}\partial_s \overline{p} + B_{21}\partial_\lambda \overline{p} &= g^1 \\
\overline{Q}^2 \partial_\lambda^2 \overline{u}^2 + B_{12}\partial_s \overline{p} + B_{22}\partial_\lambda \overline{p} &= g^2 \\
\overline{Q}^2 B_{21}\partial_\lambda^2 \overline{u}^1 + B_{12}B_{11}\partial_s \overline{p} + B_{12}^2 \partial_\lambda \overline{p} &= B_{12} g^1 \\
\overline{Q}^2 B_{22}\partial_\lambda^2 \overline{u}^2 + B_{22}B_{12}\partial_s \overline{p} + B_{22}^2 \partial_\lambda \overline{p} &= B_{22} g^2
\end{aligned}$$

This implies

$$(B_{21}^2 + B_{22}^2)\partial_\lambda \overline{p} = g^4, \text{ where } g^4 \in L^2\left(C/2\right),$$

and since $(B_{21}^2 + B_{22}^2) > 0$ for a small enough λ_0, and we get that $\partial_\lambda \overline{p} \in L^2\left(C/2\right)$. Finally, we use this to get that

$$\begin{aligned}
\overline{Q}^2 \partial_\lambda^2 \overline{u}^1 &= g^5, \text{ where } g^5 \in L^2\left(C/2\right). \\
\overline{Q}^2 \partial_\lambda^2 \overline{u}^2 &= g^6, \text{ where } g^6 \in L^2\left(C/2\right).
\end{aligned}$$

This completes the regularity proof, since we can cover a $\frac{\lambda_0}{2}$ neighbourhood by a finite number of sets of type $C/2$.

We are only left to show that we can apply [2, Theorem 10.5, p.78] to our system in order to obtain higher regularity. Indeed we can apply this theorem to show:

$$\|u\|_{H^{s+1}} + \|p\|_{H^s} \leq \|f\|_{H^{s-1}}. \tag{38}$$

What follows is a confirmation that our problem fulfills the ellipticity conditions of [2]. To adapt our notation to the one in [2] we will write $(u^1, u^2, u^3) = (u^1, u^2, p)$.

In [2, pp.38, 42] the system is written like

$$l_{ij}(x, \partial)u^j = F_i,$$

with the boundary condition

$$B_{h,j}(x, \partial)u_j = \phi_h.$$

In our case we have the correspondence

$$\begin{aligned} l_{11} = 1 - Q^2\Delta, \quad l_{12} = 0, \quad l_{13} = A_{k1}\partial_l, \quad F_1 = f_1 \\ l_{21} = 0, \quad l_{22} = 1 - Q^2\Delta, \quad l_{23} = A_{k2}\partial_l, \quad F_2 = f_2 \\ l_{31} = A_{k1}\partial_k, \quad l_{32} = A_{k2}\partial_k, \quad l_{33} = 0, \quad F_3 = 0 \end{aligned}$$

And the indices t_j and s_j can be taken as

$$\begin{aligned} t_1 = 2, \quad t_2 = 2, \quad t_3 = 1 \\ s_1 = 0, \quad s_2 = 0, \quad s_3 = -1 \end{aligned}$$

It can be checked that with this choice l'_{ij} is given by

$$\begin{aligned} &l'_{11} = -Q^2\Delta, \quad l'_{12} = 0, \quad l'_{13} = A_{k1}\partial_l, \\ &l'_{21} = 0, \quad l'_{22} = -Q^2\Delta, \quad l'_{23} = A_{k2}\partial_l, \\ &l'_{31} = A_{k1}\partial_k, \quad l'_{32} = A_{k2}\partial_k, \quad l'_{33} = 0, \end{aligned}$$

where l'_{ij} is defined in pages 38-39 of [2]. Also

$$\begin{aligned} &B_{2l} = (t^i A_{il} n_j + t^j A_{il} n_i)\partial_j, \quad l = 1, 2 \\ &B_{23} = 0 \end{aligned}$$

and the indices $r_1 = -1$ and $r_2 = -1$. With this choice $B'_{ij} = B_{ij}$ (see [2, p.42]).

We can write

$$(l'(x,\xi))_{ij} = \begin{pmatrix} -Q^2(\xi_1^2 + \xi_2^2) & 0 & A_{k1}\xi_k \\ 0 & -Q^2(\xi_1^2 + \xi_2^2) & A_{k2}\xi_k \\ A_{k1}\xi_k & A_{k2}\xi_k & 0 \end{pmatrix}$$

and

$$B(x,\xi)_{ij} = \begin{pmatrix} -2n^i A_{i1}(n\cdot\xi) & -2n^i A_{i2}(n\cdot\xi) & 1 \\ t^i A_{i1}(\xi\cdot n) + n^i A_{i1}(t\cdot\xi) & t^i A_{i2}(\xi\cdot n) + n^i A_{i2}(t\cdot\xi) & 0 \end{pmatrix}$$

Let $L = \det(l'_{ij}(x,\xi))$, i.e.

$$\begin{aligned} L(x,\xi) = \det(l'(x,\xi)) &= Q^2(\xi_1^2 + \xi_2^2)(A_{k1}\xi_k)^2 + Q^2(\xi_1^2 + \xi_2^2)(A_{k2}\xi_k)^2 \\ (A_{11}\xi_1 + A_{21}\xi_2)^2 &= A_{11}^2\xi_1^2 + A_{21}^2\xi_2^2 + 2A_{11}A_{21}\xi_1\xi_2 \\ (A_{12}\xi_1 + A_{22}\xi_2)^2 &= A_{12}^2\xi_1^2 + A_{22}^2\xi_2^2 + 2A_{12}A_{22}\xi_1\xi_2 \\ \Rightarrow L(P,\xi) &= Q^2(\xi_1^2 + \xi_2^2)Q^2(\xi_1^2 + \xi_2^2) = Q^4(\xi_1^2 + \xi_2^2)^2 = Q^4|\xi|^4 \end{aligned}$$

To know whether the system is uniformly elliptic, we are left to check that it satisfies the Supplementary condition (see [2, p.39]). The degree is 4 and therefore $m = 2$. We need to compute the solutions of $L(P, \xi + \tau\xi') = 0$, which are the solutions to $|\xi + \tau\xi'|^2 = 0$.

$$|\xi + \tau\xi'|^2 = |\xi|^2 + \tau^2|\xi'|^2 + 2\tau\xi\cdot\xi' = 0$$

Solving in τ yields

$$\tau = \frac{-2\xi\cdot\xi' + \sqrt{4(\xi\cdot\xi')^2 - 4|\xi|^2|\xi'|^2}}{2|\xi'|^2}.$$

If ξ and ξ' are linearly independent, then the discriminant is strictly negative, which implies that there is a complex root with positive imaginary part. Since the roots have multiplicity 2, the Supplementary condition is satisfied. In addition uniform ellipticity is easy to obtain. Next we check that the Complementing Boundary Condition is satisfied ([2, p. 42]).

Let t_0 be tangential vector and n_0 the normal one.

Since i is a double root of $L(x, t_0 + \tau n)$ and $M^+(x, \xi, \tau)$ ([2, p. 42]) is given by

$$M^+(x, \xi, \tau) = (\tau - i)^2.$$

In addition

$$(l'(x, t_0 + \tau n))_{ij} = \begin{pmatrix} -Q^2(1+\tau^2) & 0 & A_{k1}(t_0^k + \tau n^k) \\ 0 & -Q^2(1+\tau^2) & A_{k2}(t_0^k + \tau n^k) \\ A_{k1}(t_0^k + \tau n^k) & A_{k2}(t_0^k + \tau n^k) & 0 \end{pmatrix}$$

We define L_{ij} as in [2, p.42], and we have that. $L(x, t_0 + \tau n) = \overline{l'(x, t + \tau n)}$. Also, it can be computed that

$$B(x, t + \tau n) = \begin{pmatrix} -2n^i A_{i1}\tau & -2n^i A_{i2}\tau & 1 \\ (n + \tau t)^i A_{i1} & (n + \tau t)^i A_{i2} & 0 \end{pmatrix}$$

$$L_{ij}(x, t_0 + \tau n) = \begin{pmatrix} -Q^2(1+\overline{\tau}^2) & 0 & A_{k1}(t + \overline{\tau} n)^k \\ 0 & -Q^2(1+\overline{\tau}^2) & A_{k2}(t + \overline{\tau} n)^k \\ A_{k1}(t + \overline{\tau} n)^k & A_{k2}(t + \overline{\tau} n)^k & 0 \end{pmatrix}$$

$$\begin{aligned} &(B(x, t + \tau n)L(x, t + \tau n))_{ij} \\ &= \begin{pmatrix} 2n_k A_{k1}\tau Q^2(1+\overline{\tau}^2) + A_{k1}(t+\overline{\tau}n)^k & 2n_k A_{k2}\tau Q^2(1+\overline{\tau}^2) + A_{k2}(t+\overline{\tau}n)^k & n^i A_{i1}\tau A_{k1}(t+\overline{\tau}n)^k + \tau n^i A_{i2}A_{k2}(t+\overline{\tau}n)^k \\ -(n+\tau t)^k A_{k1}Q^2(1+\overline{\tau}^2) & -(n+\tau t)^i A_{i2}Q^2(1+\overline{\tau}^2) & (n+\tau t)^i A_{i1}A_{k1}(t+\overline{\tau}n)^k + (n+\tau t)^i A_{i2}A_{k2}(t+\overline{\tau}n)^k \end{pmatrix} \end{aligned}$$

If the rows of $(BL)_{ij}$ are linearly independent modulo $(\tau - i)^2$, that means that the condition

$$C_h B_{hj} L_{jk} = 0 \qquad \text{mod} \quad M^+$$

implies that all $C_h = 0$ ([2, p.43]). If this condition is satisfied in particular

$$C_h B_{hj} L_{jk}|_{\tau = i} = 0.$$

Then c_3 must be zero and the following system of equations must be satisfied:

$$\begin{aligned} c_1 A_{k1} t^k + c_2 A_{k2} t^k &= 0 \\ c_1 A_{k1} n^k + c_2 A_{k2} n^k &= 0 \end{aligned}$$

In matricial form:

$$\begin{pmatrix} A_{k1}t^k & A_{k2}t^k \\ A_{k1}n^k & A_{k2}n^k \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

But the determinant of this matrix satifies

$$\begin{gathered} A_{k1}t^k A_{k2}n^k - A_{k2}t^k A_{k1}n^k \\ (A^*t)^1(A^*n)^2 - (A^*t)^2(A^*n)^1 \\ (A^*t, JA^*n) = (t, AJA^*n) \neq 0, \end{gathered}$$

and therefore $c_1 = c_2 = 0$. Thus we have checked that the complementing Boundary Condition is satisfied.

Finally we notice that our system can be written as in [2, p. 71], where the coefficients $a_{ij,P}$ are smooth. In our case the index l_1 in [2, p. 77] is $l_1 = 0$. The index l in [2] coincides with $s-1$. That means $l = 0, 1, 2$. The regularity we ask for the coefficient $b_{hj,\sigma}$ is C^{l-r_h} (cf. [2, p. 77]). The most we need is therefore $b_{hj,\sigma} \in C^3$. Since these coefficients are one derivative less regular that the boundary, a C^4 boundary is enough for our purpose. This fact finishes the proof of the inequality (38) if s is an integer. For the rest of the values we proceed by interpolation.

So far, we have solved the system with Rf. We need to do it with a general f

$$u - Q^2\Delta u + A^*\nabla p = \overline{f}$$

Decomposing $\overline{f} = R\overline{f} + \nabla\pi_{\overline{f}}$, we obtain

$$u - Q^2\Delta u + A^*\nabla p = R\overline{f} + \nabla\pi_{\overline{f}},$$

and we choose p in such a way that

$$A^*\nabla p = RA^*\nabla p + A^*\nabla\pi_{\overline{f}},$$

and we know that

$$RA^*\nabla p = A^*\nabla p_1.$$

Hence, we need to solve

$$u - Q^2\Delta u + A^*\nabla p_1 = R\overline{f},$$

which we have already done before. This concludes the proof of lemma 4.3. □

Once we have studied the operator S_A we will solve the time evolution. First we will show the following lemma.

Lemma 4.4 *Let $1 \le s \le 3$, $\lambda \in \mathbb{C}$, $\Re(\lambda) \ge 0$. Then the operator $\lambda + S_A : V^{s+1} \to RH^{s-1}$ is invertible. The inverse satisfies:*

$$\|(\lambda + S_A)^{-1}Rf\|_{H^{s+1}} \le C(\|Rf\|_{H^{s-1}} + |\lambda|^{\frac{s-1}{2}}\|Rf\|_{L^2}) \tag{39}$$

Proof: As before, we look for a weak solution of $(\lambda + S_A)v = Rf, \quad f \in L^2$. Therefore,

$$(1+\lambda)(v, w) + \langle v, w\rangle = \int_\Omega Rf\bar{w}\frac{dx}{Q^2} \quad \forall w \in RH^1.$$

The solution is given by Lax-Milgram's Theorem. For $w = v$, by virtue of Korn's inequality and $\Re(\lambda) \ge 0$, one obtains

$$|(1+\lambda)(v, v) + \langle v, v\rangle| \ge C((1 + |\lambda|)\|v\|^2_{L^2} + \|\nabla v\|^2_{L^2}),$$

with C independent of λ. Easily, the following bound is obtained:

$$\|v\|_{L^2} \le \frac{1}{1+|\lambda|}\|Rf\|_{L^2}. \tag{40}$$

If $f \in H^{s-1}$, then Lemma 4.3 gives $v = S_A^{-1}(Rf - \lambda v) \in H^{s+1}$, and we can get the following bounds:

$$\begin{aligned}\|v\|_{H^{s+1}} &\le C\|S_A v\|_{H^{s-1}} \le C(\|(S_A + \lambda)v\|_{H^{s-1}} + |\lambda|\|v\|_{H^{s-1}})\\ &\le C\|Rf\|_{H^{s-1}} + |\lambda|\|v\|^{\frac{s-1}{s+1}}_{H^{s+1}}\|v\|^{\frac{2}{s+1}}_{L^2}.\end{aligned}$$

The case $s = 1$ is already solved using (40). Young's inequality provides

$$\|v\|_{H^{s+1}} \le C\|Rf\|_{H^{s-1}} + \frac{1}{2}\|v\|_{H^{s+1}} + |\lambda|^{\frac{s+1}{2}}\|v\|_{L^2}.$$

and we can get (39) using (40).

□

In order to find the solution of

$$v_t + S_A(v) = Rf \quad \text{ in } V^{s+1},$$

we take Fourier transforms in time. Since $f \in H^{ht,s-1}, Rf(0) = 0$, we can extend Rf to a function $\overline{Rf}$ defined in $H^{ht,s-1}(\Omega \times \mathbb{R})$, with $\overline{Rf}(t) = 0$ for all $t < 0$. Since $\frac{s-1}{2} \leq \frac{3}{4} < 1$, using Lemma 3.2 (ii) we get

$$\|\overline{Rf}\|_{H^{ht,s-1}(\Omega\times\mathbb{R})} \leq C\|Rf\|_{H^{ht,s-1}(\Omega\times[0,T])},$$

with C independent of T. We look for a solution of

$$v_t + S_A(v) = \overline{Rf}, \quad \forall t \in \mathbb{R}, \quad v(0) = 0.$$

By Fourier, $i\tau\hat{v}(\tau) + S_A(\hat{v})(\tau) = \hat{\overline{Rf}}$, and therefore the solution is given by $\hat{v}(\tau) = (i\tau + S_A)^{-1}\hat{\overline{Rf}}$. Using (39) and (40) we can bound

$$\begin{aligned}\|v\|_{H^{ht,s+1}(\Omega\times\mathbb{R})} &= \int_{\mathbb{R}} (\|\hat{v}\|^2_{H^{s+1}}(\tau) + |\tau|^{s+1}\|\hat{v}\|^2_{L^2}(\tau))d\tau \leq C\int_{\mathbb{R}} (\|\overline{Rf}\|^2_{H^{s-1}}(\tau) + |\tau|^{s-1}\|\overline{Rf}\|^2_{L^2})d\tau \\ &\leq C\|\overline{Rf}\|_{H^{ht,s-1}} \leq C\|Rf\|_{H^{ht,s-1}}.\end{aligned}$$

Since $\overline{Rf}(t) = 0$ for every $t < 0$, $\hat{\overline{Rf}}(\tau)$ has an analytic extension in τ to $\Im(\tau) < 0$. Using Lemma 4.4, $\hat{v}(\tau)$ also has that extension. Moreover, (40) gives $\|\hat{v}(\tau)\|_{L^2} \leq C\|\hat{\overline{Rf}}\|_{L^2}$. Thus, Paley-Wiener provides $v(t) = 0 \ \forall t < 0$. Since $v \in H^{\frac{s+1}{2}}(\mathbb{R}; L^2)$ and $\frac{3}{2} < \frac{s+1}{2}$ we have continuity in time and hence $v(0) = 0$. Since $\hat{v} \in L^2(\mathbb{R}; V^{s+1}(\Omega))$ we have that $v \in L^2([0,T]; V^{s+1}(\Omega))$ and therefore $Tr(\nabla vA) = 0$ and $(q + \nabla vA + (\nabla vA)^*)A^{-1}n = 0$. We have already solved

$$v_t - Q^2\Delta v + A^*\nabla q_1 = Rf,$$

where q_1 satisfies

$$\begin{aligned}Q^2\Delta q_1 &= 0, \text{ in } \Omega_0 \times [0,T] \\ q_1 &= A^{-1}n(\nabla vA + (\nabla vA)^*)A^{-1}n, \text{ on } \partial\Omega \times [0,T].\end{aligned}$$

Definition

$$A^*\nabla q = (I - R)f + A^*\nabla q_1,$$

gives us the solution that we were looking for

$$v_t - Q^2\Delta v + A^*\nabla q = f.$$

The properties of R allow us to obtain

$$\begin{aligned}\|\nabla q\|_{H^{ht,s-1}} \leq C\|A^*\nabla q\|_{H^{ht,s-1}} &\leq \|(I-R)f\|_{H^{ht,s-1}} + \|A^*\nabla q_1\|_{H^{ht,s-1}} \\ &\leq C(\|f\|_{H^{ht,s-1}} + \|A^*\nabla q_1\|_{H^{ht,s-1}}).\end{aligned}$$

We have the following bounds:

$$\begin{aligned}\|A^*\nabla q_1\|_{L^2H^{s-1}} \leq \|\nabla q_1\|_{L^2H^{s-1}} &\leq |A^{-1}n((\nabla vA) + (\nabla vA)^*)A^{-1}n|_{L^2H^{s-\frac{1}{2}}} \\ &\leq \|(\nabla vA) + (\nabla vA)^*\|_{L^2H^s} \leq \|\nabla v\|_{L^2H^s} \leq \|v\|_{L^2([0,T];H^{s+1})},\end{aligned}$$

$$\|A^*\nabla q_1\|_{H^{\frac{s-1}{2}}L^2} \leq |A^{-1}n((\nabla vA) + (\nabla vA)^*)A^{-1}n|_{H^{\frac{s-1}{2}}H^{\frac{1}{2}}}.$$

Decomposing ∇v_i into the tangential and normal components: $\nabla v_i = (\nabla v_i \cdot n_0)n_0 + (\nabla v_i \cdot t_0)t_0$, we can bound each of them by

$$\begin{aligned}|\nabla v_i \cdot t_0|_{H^{\frac{s-1}{2}}H^{\frac{1}{2}}} &\leq C|v_i|_{H^{\frac{s-1}{2}}H^{\frac{3}{2}}} \leq C\|v_i\|_{H^{\frac{s-1}{2}}H^2} \leq C\|v_i\|_{H^{ht,s+1}}, \\ |\nabla v_i \cdot n_0|_{H^{\frac{s-1}{2}}H^{\frac{1}{2}}} &\leq |\nabla v_i \cdot n_0|_{H^{ht,s-\frac{1}{2}}} \leq C\|v_i\|_{H^{ht,s+1}}.\end{aligned}$$

Above we have used Lemma 3.1 (i). This yields

$$\|A^*\nabla q_1\|_{H^{\frac{s-1}{2}}L^2} \leq C\|v\|_{H^{ht,s+1}}.$$

and therefore

$$\|A^*\nabla q\|_{H^{ht,s-1}} \leq C\|f\|_{H^{ht,s-1}}.$$

Since $(I-R)f|_{\partial\Omega_0} = 0$, we have that $q = q_1$ on $\partial\Omega_0$. This implies

$$\begin{aligned} |q|_{H^{ht,s-\frac{1}{2}}} = |q_1|_{H^{ht,s-\frac{1}{2}}} &= |A^{-1}n((\nabla vA)+(\nabla vA)^*)A^{-1}n|_{H^{ht,s-\frac{1}{2}}} \\ &\leq |A^{-1}n((\nabla vA)+(\nabla vA)^*)A^{-1}n|_{L^2H^{s-\frac{1}{2}}} + |A^{-1}n((\nabla vA)+(\nabla vA)^*)A^{-1}n|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}. \end{aligned}$$

Decomposing into the tangential and normal components as before, we find finally

$$\begin{aligned} |q|_{H^{ht,s-\frac{1}{2}}} &\leq \|\nabla v\|_{L^2H^s} + |\nabla v_i \cdot t_0|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} + |\nabla v_i \cdot n_0|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} \\ &\leq \|v\|_{L^2H^{s+1}} + |v_i|_{H^{\frac{s}{2}-\frac{1}{4}}H^1} + |\nabla v_i \cdot n_0|_{H^{ht,s-\frac{1}{2}}} \\ &\leq \|v\|_{H^{ht,s+1}} + \|v\|_{H^{\frac{s}{2}-\frac{1}{4}}H^{1+\frac{1}{2}}} + \|v\|_{H^{ht,s+1}} \leq \|v\|_{H^{ht,s+1}}. \end{aligned}$$

4.2 Reduction for arbitrary g, h and v_0.

In this section we want to extend the result with $g = 0, h = 0$ and $v(x,0) = 0$ to the general case:

$$\begin{aligned} v_t - \nu Q^2\Delta v + A^*\nabla q &= f && \text{in } \Omega_0 \times [0,T] \\ Tr(\nabla vA) &= g && \text{in } \Omega_0 \times [0,T] \\ (q + (\nabla vA) + (\nabla vA)^*)A^{-1}n &= h && \text{on } \partial\Omega_0 \times [0,T] \\ v(x,0) &= v_0(x) && \text{in } \Omega_0 \end{aligned} \tag{41}$$

To get the space where g belongs we proceed as follows. For $\phi \in H^1_0$ it is easy to find

$$\int \partial_t^j(Tr(\nabla vA))\phi(x)\frac{dx}{Q^2(x)} = \int Tr(\nabla\partial_t^j vA)\phi(x)\frac{dx}{Q^2(x)} = \int \partial_t^j vA^*\nabla\phi\frac{dx}{Q^2(x)}.$$

Then

$$\left|\int \partial_t^j(Tr(\nabla vA))\phi(x)\frac{dx}{Q^2(x)}\right| \leq \|\partial_t^j v\|_{L^2}(t)\|\nabla\phi\|_{H^1},$$

and duality provides

$$\|\partial_t^j(Tr(\nabla vA))\|_{H^{-1}} \leq \|\partial_t^j v\|_{L^2}(t), \text{ with } H^{-1} = (H^1_0)^*.$$

For $j = \frac{s+1}{2}$, integration in time yields

$$\|Tr(\nabla vA)\|_{H^{\frac{s+1}{2}}H^{-1}} \leq \|v\|_{H^{ht,s+1}}.$$

Also, $Tr(\nabla vA) \in L^2([0,T];H^s)$, which implies:

$$Tr(\nabla vA) \in L^2([0,T];H^s) \cap H^{\frac{s+1}{2}}([0,T];H^{-1}) = \overline{H}^{ht,s}$$

To prove this fact, one can proceed for an integer number of derivatives, then interpolate for fractional ones (see [27]).

We check next the compatibility conditions of the initial data:

$$Tr(\nabla v_0A) = g(0) \qquad \text{in } \Omega_0 \tag{42}$$

$$(A^{-1}n)^{\perp}(\nabla v_0A + (\nabla v_0A)^*)A^{-1}n = h(0)(A^{-1}n)^{\perp} \qquad \text{on } \partial\Omega_0. \tag{43}$$

Taking $2 < s < \frac{5}{2}, \nabla q \in H^{ht,s-1}(\Omega \times [0,T]), q \in H^{ht,s-\frac{1}{2}}(\partial\Omega \times [0,T])$, and defining the following spaces:

$X := \{(v,q) : v \in H^{ht,s+1}, q \in H^{ht,s}_{pr}\}$

$Y := \{(f,g,h,v_0) : f \in H^{ht,s-1}, g \in \overline{H}^{ht,s}, h \in H^{ht,s-\frac{1}{2}}(\partial\Omega \times [0,T]), v_0 \in H^s(\Omega)$ and $(42)-(43)$ are satisfied$\}$

Then (41) is equivalent to solve:

$$L(v,q) = (f,g,h,v_0); \quad L : X \to Y, \quad 2 < s < \frac{5}{2}.$$

Theorem 4.5 *Let $2 < s < \frac{5}{2}$. Then $L : X \to Y$ has a bounded inverse:*

$$\|(v,q)\|_X \leq C\|(f,g,h,v_0)\|_Y$$

Proof: We structure the proof in 4 steps, in order to get to the previous case ($g = h = v_0 = 0$). Let $(f,g,h,v_0) \in Y$.

Step 1: Initial conditions. We want to find $(v^1,q^1) \in X$ such that $L(v^1,q^1) = (f^1,g^1,h^1,v^1_0)$ with

$$f^1(0) = f(0); \quad g^1(0) = g(0); \quad h^1(0) = h(0); \quad v^1_0 = v_0. \tag{44}$$

We can first define $q^1(0) \in H^{s-\frac{3}{2}}(\partial\Omega)$ using

$$q^1(0)|A^{-1}n|^2 = h(0)A^{-1}n - A^{-1}n(\nabla v_0 A + (\nabla v_0 A)^*)A^{-1}n. \tag{45}$$

By the Trace theorem we can extend it to $q^1(0) \in H^{s-1}(\Omega)$ and we can use the parabolic trace (Lemma 3.1) to obtain $q^1 \in H^{ht,s}$. We can use now the parabolic trace to obtain $v^1 \in H^{ht,s+1}$

$$\begin{aligned} v^1(0) &= v_0 \in H^s, \\ \partial_t v^1(0) &= Q^2\Delta v_0 - A^*\nabla q^1(0) + f(0), \qquad \partial_t v^1(0) \in H^{s-2}. \end{aligned} \tag{46}$$

Therefore (44) is satisfied, thanks to (42),(43),(45) and (46). We have

$$\|(v^1,q^1)\|_X \leq C\|(f,g,h,v_0)\|_Y.$$

Step 2: A-divergence adjustment. We want to find $(v^2,q^2) \in X$ such that $L(v^2,q^2) = (f^2,g^2,h^2,v^2_0)$ with

$$Rf^2(0) = Rf(0); \quad g^2(t) = g(t);\ \forall t \in [0,T] \quad h^2(0) = h(0); \quad v^2_0 = v_0. \tag{47}$$

We take $\overline{g}(t) = g(t) - g^1(t)$ in $\overline{H}^{ht,s}$, and we define ϕ by solving the following elliptic problem for every $t \in (-\infty,\infty)$, after extending $\overline{g}$ to the whole real line:

$$\begin{aligned} Q^2\Delta\phi &= \overline{g}(t) \qquad \text{in } \Omega_0 \times \mathbb{R} \\ \phi &= 0 \qquad \text{on } \partial\Omega_0 \times \mathbb{R} \end{aligned}$$

This system satisfies $\|\phi\|_{H^{s+2}}(t) \leq C\|g\|_{H^s}(t)$. In particular

$$\|\nabla\phi\|_{H^{s+1}}(t) \leq C\|g\|_{H^s}(t). \tag{48}$$

Taking the Fourier Transform in time, one has, for every τ:

$$\begin{aligned} Q^2\Delta\hat{\phi}(\tau) &= \hat{\overline{g}}(\tau) \qquad \text{in } \Omega_0 \times \mathbb{R}, \\ \hat{\phi}(\tau) &= 0 \qquad \text{on } \partial\Omega_0 \times \mathbb{R}. \end{aligned}$$

For $\lambda \in H^1_0$ it is possible to find

$$\int \hat{\overline{g}}(\tau)\lambda\frac{dx}{Q^2} = \int \Delta\hat{\phi}(\tau)\lambda dx = \int \nabla(\nabla\hat{\phi}A)A\lambda\frac{dx}{Q^2} = -\int \nabla\hat{\phi}A\underbrace{\nabla\lambda A}_{A^*\nabla\lambda}\frac{dx}{Q^2} = -\int \nabla\hat{\phi}\nabla\lambda dx$$

Therefore:

$$\int \nabla\hat{\phi}\nabla\lambda dx = -\int \hat{\overline{g}}(\tau)\lambda\frac{dx}{Q^2} \quad \forall\lambda \in H^1_0.$$

This implies:

$$\begin{gathered}\|\nabla\hat{\phi}\|_{L^2}(\tau) \leq C\|\hat{\overline{g}}\|_{H^{-1}}(\tau)\\ \Rightarrow \int |\tau|^{\frac{s+1}{2}}\|\nabla\hat{\phi}\|_{L^2}(\tau)d\tau \leq C\int |\tau|^{\frac{s+1}{2}}\|\hat{\overline{g}}\|_{H^{-1}}(\tau)d\tau.\end{gathered}$$

Using (48), we can conclude that $\nabla\phi \in H^{ht,s+1}$. Furthermore, since $\overline{g}(0) = g(0) - g^1(0)$, this implies via (44) that $\overline{g}(0) = 0$, and therefore $\phi(0) = 0$. We now define

$$v^2 = v^1 + A^*\nabla\phi \in H^{ht,s+1}, \quad q^2 = q^1.$$

It is easy to check that

$$\begin{aligned} f^2(0) &= \partial_t v^2(0) - Q^2\Delta v^2(0) + A^*\nabla q^1(0)\\ &= \partial_t v^1(0) - Q^2\Delta v^1(0) + A^*\nabla q^1(0) + A^*\nabla\phi_t(0) - Q^2\Delta(A^*\nabla\phi(0))\\ &= f(0) + A^*\nabla\phi_t(0)\end{aligned}$$

This implies $Rf^2(0) = Rf(0)$, since $R(A^*\nabla\phi_t) = 0$ because of Lemma 4.1. By construction, it is obvious that $g^2(t) = g(t)$ for every $t \in [0,T]$. Since $\phi(0) = 0$, $h^2 = h(0)$ and $v^2(0) = v_0$. This shows that (47) is satisfied.

Step 3: Adjusting the boundary conditions in the tangential direction without modifying the A-divergence.

We want to find $(v^3, q^3) \in X$ such that $L(v^3, q^3) = (f^3, g^3, h^3, v^3_0)$ with

$$\begin{aligned} &Rf^3(0) = Rf(0); \quad g^3(t) = g(t); \ \forall t \in [0,T]\\ &h^3(t)\cdot(A^{-1}n)^\perp = h(t)\cdot(A^{-1}n)^\perp; \ \forall t \in [0,T], \quad h^3(0) = 0; \quad v^3_0 = v_0 \end{aligned} \tag{49}$$

We will use the following Lemma:

Lemma 4.6 *Let* $\eta \in H^{ht,s-\frac{1}{2}}(\partial\Omega_0 \times [0,T])$, $2 < s < \frac{5}{2}$, $\eta(0) = 0$. *Then there exists* $w \in H^{ht,s+1}$ *such that* $\|w\|_{H^{ht,s+1}} \leq C|\eta|_{H^{ht,s-\frac{1}{2}}}$, $w(0) = w_t(0) = Tr(\nabla wA) = 0$ *and*

$$(A^{-1}n)^\perp(\nabla wA + (\nabla wA)^*)A^{-1}n = \eta \text{ on } \partial\Omega_0 \times [0,T].$$

Proof:

Let $\psi \in H^{ht,s+2}$, with $\psi(0) = \psi_t(0) = 0, \psi(x) = \partial_n\psi(x) = 0$, $\forall x \in \partial\Omega_0$, and moreover $\partial^2_n\psi(x) = \eta(x)$ $\forall x \in \partial\Omega_0 \times [0,T]$. This choice is possible because of the parabolic trace. All that is left is to check that the compatibility conditions from Lemma 3.1. Defining

$$w = \nabla^\perp_A\psi = (-\partial_2P_1 \circ P^{-1}\partial_1\psi - \partial_2P_2 \circ P^{-1}\partial_2\psi, \partial_1P_1 \circ P^{-1}\partial_1\psi + \partial_1P_2 \circ P^{-1}\partial_2\psi),$$

it is immediate that $w(0) = w_t(0) = 0$. A straightforward, but long calculation gives:

$$Tr(\nabla(\nabla^\perp_A\psi)A) = 0 = Tr(\nabla wA).$$

We will now show that $(A^{-1}n)^\perp(\nabla wA + (\nabla wA)^*)(A^{-1}n) = \eta$. Let $x_0 \in \partial\Omega_0$. We perform an euclidean change of coordinates in a way that $x_0 = 0, n_0 = (0,1)$. Thus, $\psi(0,0) = 0$, as well as

$\partial_1\psi(0,0)$ and $\partial_1^2\psi(0,0)$. The condition $\partial_n\psi(0,0)=0$ implies $\partial_2\psi(0,0)=0, \partial_1\partial_2\psi(0,0)=0$. That gives:

$$\nabla(\nabla_A^\perp\psi)=\begin{pmatrix} 0 & -\partial_2 P_2\circ P^{-1}\partial_2^2\psi(0,0) \\ 0 & \partial_1 P_2\circ P^{-1}\partial_2^2\psi(0,0)\end{pmatrix}.$$

Computing further:

$$\nabla(\nabla_A^\perp\psi)A=\partial_2^2\psi(0,0)\begin{pmatrix} -\partial_2 P_2\partial_1 P_2 & -(\partial_2 P_2)^2 \\ (\partial_1 P_2)^2 & \partial_1 P_2\partial_2 P_2\end{pmatrix}\circ P^{-1}$$

Thus

$$((\nabla(\nabla_A^\perp\psi)A)+(\nabla(\nabla_A^\perp\psi)A)^*)=\partial_2^2\psi(0,0)\begin{pmatrix} -2\partial_2 P_2\partial_1 P_2 & (\partial_1 P_2)^2-(\partial_2 P_2)^2 \\ (\partial_1 P_2)^2-(\partial_2 P_2)^2 & 2\partial_1 P_2\partial_2 P_2\end{pmatrix}\circ P^{-1}.$$

We also have that

$$A^{-1}n_0=\frac{1}{Q^2}\begin{pmatrix} -\partial_2 P_1 \\ \partial_1 P_1\end{pmatrix}\circ P^{-1};\quad (A^{-1}n_0)^\perp=\frac{1}{Q^2}\begin{pmatrix} -\partial_1 P_1 \\ -\partial_2 P_1\end{pmatrix}\circ P^{-1};$$

Combining everything, we get

$$\begin{aligned}((\nabla(\nabla_A^\perp\psi)A)+(\nabla(\nabla_A^\perp\psi)A)^*)A^{-1}n_0 &= \frac{\partial_2\psi(0,0)}{Q^2}(-\partial_1 P_1\circ P^{-1}Q^2, \partial_1 P_2\circ P^{-1}Q^2) \\ \Rightarrow (A^{-1}n)^\perp((\nabla(\nabla_A^\perp\psi)A)+(\nabla(\nabla_A^\perp\psi)A)^*)A^{-1}n_0 &= \partial_2^2\psi(0,0)\end{aligned}$$

Since $\partial_n^2\psi(0,0)=\sum_{i,j}(n_0)_i\partial_i\partial_j\psi(n_0)_j=\partial_2^2\psi(0,0)$, we are done. □

We now apply Lemma 4.6 to $\eta=h(t)(A^{-1}n)^\perp-h^2(t)(A^{-1}n)^\perp$. Equation (47) shows that $\eta(0)=0$. Then, we can take:

$$v^3=v^2+w,\quad q^3=q^2.$$

Since $w(0)=w_t(0)=0$, we have that $Rf^3(0)=Rf(0)$. Also, since $Tr(\nabla wA)=0$, by (47), $g^3(t)=g(t)$ $\forall t\in[0,T]$. By construction, one has that $h^3(t)(A^{-1}n)^\perp=h(t)(A^{-1}n)^\perp$, and since $\eta(0)=0$, that implies that $h^3(0)=h(0)$. Finally, since $w(0)=0$ we obtain $v^3(0)=v_0$ and (49) is satisfied.

Step 4: $h(t)=h^4(t)\forall t$ without modifying the rest. We want to have $(v^4,q^4)\in X$. We have that $L(v^4,q^4)=(f^4,g^4,h^4,v_0^4)$ with

$$\begin{aligned} Rf^4(0)=Rf(0);\quad g^4(t)=g(t);\ \forall t\in[0,T] \\ h^4(t)=h(t);\ \forall t\in[0,T]\quad v_0^4=v_0.\end{aligned}\tag{50}$$

We first take $v^4=v^3$ and define $\overline{q}$ by

$$\overline{q}=h(t)\cdot\frac{A^{-1}n_0}{|A^{-1}n_0|^2}-\underbrace{(q^3I+(\nabla v^3A)+(\nabla v^3A)^*)A^{-1}n_0}_{h^3(t)}\cdot\frac{A^{-1}n_0}{|A^{-1}n_0|^2}\text{ on }\partial\Omega_0\times[0,T],$$

$$\overline{q}(x,0)=0,\quad\text{in }\Omega_0.$$

Using once again the parabolic trace, the compatibility condition is satisfied since $\overline{q}(x,0)=0$ in $\partial\Omega\times[0,T]$ by means of (49). Therefore, we take $q^4=q^3+\overline{q}$.

Since $\nabla\overline{q}(x,0)=0$ we have as before $Rf^4(0)=Rf^3(0)=f(0)$, and $Tr(\nabla v^4 A)=0$ because the velocity was not modified. At the boundary we find:

$$\begin{aligned}((q^4I+(\nabla v^4A)+(\nabla v^4A)^*)A^{-1}n &= \overline{q}A^{-1}n+(q^3I+(\nabla v^3A)+(\nabla v^3A)^*)A^{-1}n\\ &= \overline{q}A^{-1}n+h^3(t)=h(t)\cdot\frac{A^{-1}n}{|A^{-1}n|}\frac{A^{-1}n}{|A^{-1}n|}-h^3(t)\cdot\frac{A^{-1}n}{|A^{-1}n|}\frac{A^{-1}n}{|A^{-1}n|}+h^3(t)\\ &= h(t)\cdot\frac{A^{-1}n}{|A^{-1}n|}\frac{A^{-1}n}{|A^{-1}n|}+h^3(t)\cdot\frac{(A^{-1}n)^\perp}{|A^{-1}n|}\frac{(A^{-1}n)^\perp}{|A^{-1}n|}\\ &\overset{(49)}{=} h(t)\cdot\frac{A^{-1}n}{|A^{-1}n|}\frac{A^{-1}n}{|A^{-1}n|}+h(t)\cdot\frac{(A^{-1}n)^\perp}{|A^{-1}n|}\frac{(A^{-1}n)^\perp}{|A^{-1}n|}=h(t).\end{aligned}$$

We find $v^4(0)=v_0$ trivially. By construction, we conclude that

$$\|(v^4,q^4)\|_X\le C\|(f,g,h,v_0)\|_Y.$$

If we consider the variables $v_{\tilde f}=v-v^4, q_{\tilde f}=q-q^4$, we obtain the following problem:

$$\begin{aligned}\partial_t v_{\tilde f}-Q^2\Delta v_{\tilde f}+A^*\nabla q_{\tilde f}&=\tilde f && \text{in } \Omega\times[0,T]\\ Tr(\nabla v_{\tilde f}A)&=0 && \text{in } \Omega\times[0,T]\\ (q_{\tilde f}+(\nabla v_{\tilde f}A)+(\nabla v_{\tilde f}A)^*)A^{-1}n&=0 && \text{in } \partial\Omega\times[0,T]\\ v_{\tilde f}(x,0)&=0 && \text{in } \Omega,\end{aligned}\tag{51}$$

with $R\tilde f(0)=0$. Using the results from the previous section, the Theorem is proved.

□

Let us define the following spaces:

$$\begin{aligned}X_0&:=\{(v,q)\in X: v(0)=0, v_t(0)=0, q(0)=0\}\\ Y_0&:=\{(f,g,h,0)\in Y: f(0)=0, g(0)=0, g_t(0)=0, h(0)=0\}\end{aligned}$$

Theorem 4.7 *$L: X_0\to Y_0$ is invertible for $2<s<\frac{5}{2}$. Moreover, $\|L^{-1}\|$ is bounded uniformly if T is bounded above.*

Proof: It is clear that $L(X_0)\subset Y_0$. We take now $(f,g,h,0)\in Y_0$ and consider $L^{-1}(f,g,h,0)=(v,q)$. Then, $v(0)=0$ and for every $x\in\partial\Omega$:

$$0=h(0)A^{-1}n=A^{-1}n(q(0)I+\underbrace{\nabla v(0)A}_{0}+\underbrace{(\nabla v(0)A)^*}_{0})A^{-1}n=q(0)|A^{-1}n|^2,$$

which implies $q(0)=0$ on $\partial\Omega$.
We also have that $v(t)=A^*\nabla\pi(t)+R(v)(t)$, with $\pi\in H^1_0(\Omega)\ \forall t$ and $Q^2\Delta\pi(t)=g(t)$.
Since $g_t(0)=0$, that implies that

$$\begin{aligned}Q^2\Delta\pi_t(0)&=0 \text{ in } \Omega\\ \pi_t(0)&=0 \text{ in } \partial\Omega,\end{aligned}$$

and therefore $\pi_t(0)=0$, which means:

$$v_t(0)=A^*\nabla\pi_t(0)+R(v_t)(0)=R(v_t)(0).$$

Since $f(0)=v(0)=0$, we have that

$$v_t(0) + A^*\nabla q(0) = 0.$$

But $R(A^*\nabla q(0)) = 0$ because of the condition $q(0) = 0$ for every $x \in \partial\Omega$. Therefore $R(v_t)(0) = 0$, and $v_t(0) = 0$. This shows that $A^*\nabla q(0) = 0$ and from that it follows that $q(0) = 0$. We have proved that $(v, q) \in X_0$.

We are left to see that the norm of L^{-1} is uniformly bounded if T is bounded above. To do so, we use that in the case $f(0) = 0, g(t) = h(t) = v_0 = 0$, L^{-1} is invertible and its norm is uniformly bounded if T is bounded above (see Subsection 4.1). For the case (41), let's see that in our construction the norm will not depend on time. Since the results are based on the parabolic trace, we have to see that it is uniformly bounded if T is bounded above. In order to do it, we shall observe that the intrinsic conditions of the space Y_0 already give us the result for $2 < s < \frac{5}{2}$. Since $f(0) = 0$ and $f \in H^{ht,s-1}$, and $\frac{1}{2} < \frac{s-1}{2} < \frac{3/2}{2} \Leftrightarrow 0 < \frac{s-2}{2} < \frac{1}{4}$, the trace p does not depend on time in the extension of f (see Lemma 3.1).

We also have that $g \in H^{\frac{s+1}{2}}([0,T]; H^{-1})$ and $\frac{3}{2} < \frac{s+1}{2} < \frac{7/2}{2} \Leftrightarrow 1 < \frac{s}{2} < \frac{5}{4}$. Because of that, $g(0) = g_t(0) = 0$ allows us to take a parabolic trace independent of time.

Finally, $h \in H^{s-\frac{1}{2}}$, and $\frac{3}{4} < \frac{s}{2} - \frac{1}{4} < 1$ Therefore, we only require $h(0) = 0$ to obtain a parabolic trace independent of time.

All this reasoning allows us to conclude that $\|L^{-1}\|$ is uniformly bounded if T is bounded above.

□

5 Fixed point argument

In section 4 the system (17), (18) and (19) was solved uniformly in T (for small T) but with the initial conditions $\tilde{v}|_{t=0} = 0$ and $\partial_t\tilde{v}|_{t=0} = 0$. Then we still need to carry out a small modification of (17-20) to be able to apply the result of that section.

Let's define the approximated solution ϕ in $\tilde{\Omega}_0 \times \mathbb{R}$ as follows

$$\phi = \tilde{v}_0 + t\exp(-t^2)\left(Q^2\Delta\tilde{v}_0 - A^*\nabla\tilde{q}_\phi\right), \tag{52}$$

We shall show we can choose $\tilde{q}_\phi$ in such a way that $\partial_t\phi|_{t=0} = \partial_t v^{(n)}|_{t=0}$ for all n.

We specify who $\tilde{q}_\phi$ is. Given the system (17-20), for

$$\begin{aligned} v^{(n)} =& \tilde{v}^{(n)} \circ P \\ q^{(n)} =& \tilde{q}^{(n)} \circ P \end{aligned}$$

we have that

$$\begin{aligned} \partial_t v^{(n+1)} - \Delta v^{(n+1)} =& -\nabla q^{(n+1)} + f^{(n)} \quad \text{in } \Omega_0 \\ \nabla\cdot v^{(n+1)} =& g^{(n)} \quad \text{in } \Omega_0 \\ \left(q^{(n+1)}\mathbb{I} - (\nabla v^{(n+1)} + (\nabla v^{(n+1)})^*)\right)n_0 =& h^{(n)} \quad \text{on } \partial\Omega_0 \\ v|_{t=0} =& v_0, \end{aligned}$$

where $v_0 = \tilde{v}_0 \circ P$, $f^{(n)} = \tilde{f}^{(n)} \circ P$, $g^{(n)} = \tilde{g}^{(n)} \circ P$ and $h^{(n)} = \tilde{h}^{(n)} \circ P$. Taking the divergence on the first equation yields

$$\partial_t\nabla\cdot v^{(n+1)} - \Delta\nabla\cdot v^{(n+1)} = -\Delta q^{(n+1)} + \nabla\cdot f^{(n)}.$$

Thus, taking into account the third equation, we find $q^{(n+1)}$ by solving

$$\begin{aligned} -\Delta q^{(n+1)} =& \partial_t g^{(n)} - \Delta g^{(n)} - \nabla\cdot f^{(n)} \quad \text{in } \Omega_0 \\ q^{(n+1)}|_{\partial\Omega_0} =& n_0\left(\nabla v^{(n)} + (\nabla v^{(n)})^*\right)|_{\partial\Omega_0}\cdot n_0 + h^{(n)}|_{\partial\Omega_0}\cdot n_0. \end{aligned}$$

Next notice that $\tilde{f}^{(n)}$, $\tilde{g}^{(n)}$ and $\tilde{h}^{(n)}$ in (21), (22) and (23) are equal to zero at $t = 0$. Thus, calling $q_0^{(n)} = q^{(n)}|_{t=0}$, we have that

$$\begin{aligned} -\Delta q_0^{(n+1)} =& (\partial_t g^{(n)})|_{t=0} \\ q_0^{(n)}|_{\partial\Omega_0} =& n_0\left(\nabla v_0 + (\nabla v_0)^*\right)n_0. \end{aligned}$$

Now we focus our attention in the structure of $\tilde{g}^{(n)}$ in (22). It easy to check that $\tilde{g}^{(n)} \circ P$ can be written as

$$\tilde{g}^{(n)} \circ P = -Tr\left(\nabla v'^{(n)} \left(\nabla X^{(n)}\right)^{-1}\right) + Tr\left(\nabla v'^{(n)}\right),$$

where

$$v'^{(n)} = \tilde{v}^{(n)} \circ \tilde{X}^{-1} \circ P \circ X$$

with

$$\begin{aligned} \frac{dX^{(n)}}{dt} =& v'^{(n)}(\alpha, t) \\ X^{(n)}(\alpha, 0) =& \alpha \quad \alpha \in \Omega_0 \end{aligned}$$

Therefore

$$(\partial_t g^{(n)})|_{t=0} = Tr\left(\nabla u_0 \nabla u_0\right) = \nabla \cdot ((u_0 \cdot \nabla) u_0).$$

Because of the previous discussion we will choose q_ϕ solving

$$\begin{aligned} -\Delta q_\phi =& \nabla \cdot ((u_0 \cdot \nabla) u_0) \quad \text{in } \Omega_0 \\ q_\phi|_{\partial\Omega_0} =& n_0 \left(\nabla u_0 + (\nabla u_0)^*\right)|_{\partial\Omega_0} n_0. \end{aligned}$$

which is independent on the superscript n.

Finally one finds $\tilde{q}_\phi$ changing from Ω_0 to $\tilde{\Omega}_0$,

$$-Q^2 \Delta \tilde{q}_\phi = Tr\left(\nabla \tilde{u}_0 A \nabla \tilde{u}_0 A\right) \quad \text{in } \tilde{\Omega}_0 \tag{53}$$

$$\tilde{q}_\phi|_{\partial\tilde{\Omega}_0} \left(A^{-1}\tilde{n}_0 \cdot A^{-1}\tilde{n}_0\right) = A^{-1}\tilde{n}_0 \left(\nabla \tilde{u}_0 A + (\nabla \tilde{u}_0 A)^*\right) A^{-1}\tilde{n}_0 \quad \text{on } \partial\tilde{\Omega}_0. \tag{54}$$

Once we have defined ϕ we define the velocity $\tilde{w}^{(n)}$ by the expression

$$\tilde{w}^{(n)} = \tilde{v}^{(n)} - \phi.$$

Now it is easy to check that

$$\tilde{w}^{(n)}|_{t=0} = 0, \quad \left(\partial_t \tilde{w}^{(n)}\right)|_{t=0} = 0.$$

Then it is better for our purpose to write the system (17-20) in terms of $\tilde{w}^{(n)}$ rather than in terms of $\tilde{v}^{(n)}$. We obtain that

$$\partial_t \tilde{w}^{(n+1)} - Q^2 \Delta \tilde{w}^{(n+1)} = -A^* \nabla \tilde{q}_w^{(n+1)} + \tilde{f}^{(n)} - \partial_t \phi + Q^2 \Delta \phi - A^* \nabla \tilde{q}_\phi \quad \text{in } \tilde{\Omega}_0, \tag{55}$$

$$Tr\left(\nabla \tilde{w}^{(n+1)} A\right) = \tilde{g}^{(n)} - Tr\left(\nabla \phi A\right) \quad \text{in } \tilde{\Omega}_0 \tag{56}$$

$$\begin{aligned} &\left(\tilde{q}_w^{(n+1)} \mathbb{I} - \left(\left(\nabla \tilde{w}^{(n+1)} A\right) + \left(\nabla \tilde{w}^{(n+1)} A\right)^*\right)\right) A^{-1}\tilde{n}_0 \\ &= \tilde{h}^{(n)} - \tilde{q}_\phi A^{-1}\tilde{n}_0 + \left((\nabla \phi A) + (\nabla \phi A)^*\right) A^{-1}\tilde{n}_0 \quad \text{on } \partial\tilde{\Omega}_0, \end{aligned} \tag{57}$$

$$\tilde{w}^{(n+1)}|_{t=0} = 0 \tag{58}$$

where $\tilde{f}^{(n)}$, $\tilde{g}^{(n)}$ and $\tilde{h}^{(n)}$ are given by (21), (22) and (23) with $\tilde{v}^{(n)} = \tilde{w}^{(n)} + \phi$.

Remark 5.1 *With this choice of the function ϕ we lose regularity of the solution with respect to the initial data. One could look for more sophisticated choices of ϕ, like Beale in [5], in order to avoid this loss. However, this exceeds the scope of this paper.*

We now prove the following theorem:

Theorem 5.2 *Let $\left(\{\tilde{w}^{(n)}\}_{n=0}^\infty, \{\tilde{q}_w^{(n)}\}_{n=0}^\infty, \{\tilde{X}^{(n)}\}_{n=0}^\infty\right)$ be the sequence given by the system (55), (56), (57) and (58) and (24) with*

$$\tilde{w}^{(0)} = 0, \qquad \tilde{q}_w^{(0)} = 0, \qquad \tilde{X}^{(0)} = \alpha,$$

where $\tilde{q}_\phi$ is given by (53) *and* (54), *ϕ is given by* (52) *and $\tilde{v}^n = \tilde{w}^{(n)} + \phi$. Then*

$$\left(\{\tilde{w}^{(n)}\}_{n=0}^{\infty},\ \{\tilde{q}_w^{(n)}\}_{n=0}^{\infty},\ \{\tilde{X}^{(n)}\}_{n=0}^{\infty} - \alpha\right)$$

is a Cauchy sequence in

$$H^{ht,s+1}\left([0,T],\, \Omega_0\right) \times H_{pr}^{ht,s}\left([0,T],\, \Omega_0\right) \times F^{s+1}\left([0,T],\, \Omega_0\right)$$

for $1 < \gamma < s-1$, $2 < s < 2.5$ and

$$||\tilde{u}_0||_{H^{100}(\tilde{\Omega}_0)} < C \qquad\qquad |\partial\tilde{\Omega}_0|_{C^{100}} < C,$$

for T sufficiently small.

In order to prove this result we use propositions 5.3 and 5.4 presented below concerning the system (55-58), together with (24). We start by writing this system in the more concise form

$$L(w^{(n+1)}, q_w^{(n+1)}) = \left(f^{(n)}, g^{(n)}, h^{(n)}\right) + \left(f_\phi^L, g_\phi^L, h_\phi^L\right). \tag{59}$$

where

$$\begin{aligned} f_\phi^L &= -\partial_t\phi + Q^2\Delta\phi - A^*\nabla q, \\ g_\phi^L &= -Tr(\nabla\phi A), \\ h_\phi^L &= -q_\phi A^{-1}n_0 + (\nabla\phi A + (\nabla\phi A)^*)A^{-1}n_0. \end{aligned} \tag{60}$$

For technical reasons we rewrite the right hand side in a different way. First we notice that

$$\partial_t\zeta^{(n)} = \partial_t\left[\left(\nabla X^{(n)}\right)^{-1}\right] = -\left(\nabla X^{(n)}\right)^{-1}\nabla\partial_t X^{(n)}\left(\nabla X^{(n)}\right)^{-1},$$

and therefore

$$\partial_t\zeta^{(n)}|_{t=0} = -\nabla\left(Av_0\right).$$

We define ζ_ϕ, independent of n by the following expression

$$\zeta_\phi = \mathbb{I} + te^{-t^2}\left(\partial_t\zeta^{(n)}|_{t=0}\right),$$

and also we define A_ϕ as the matrix with entries

$$\left(A_\phi\right)_{ij} = A_{ij} + te^{-t^2}\left.\left(\frac{d}{dt}\left(A^{ij}\circ X\right)\right)\right|_{t=0} = A_{ij} + te^{-t^2}\partial_k A_{ij}A_{kl}v_{0l}.$$

By using ζ_ϕ, A_ϕ, we will write the system (59) in the following way

$$L(w^{(n+1)}, q_w^{(n+1)}) = (f^{(n)}, \overline{g}^{(n)}, h^{(n)}) + (f_\phi^L, \overline{g}_\phi^L, h_\phi^L),$$

with

$$\overline{g}^{(n)} = g^{(n)} + tr\left(\nabla\phi\zeta_\phi A_\phi\right) - tr\left(\nabla\phi A\right)$$

and

$$\overline{g}_\phi^L = g_\phi^L - tr\left(\nabla\phi\zeta_\phi A_\phi\right) + tr\left(\nabla\phi A\right)$$

In this situation we have the following result.

Proposition 5.3 *1. Let $X^{(n)} - \alpha \in F^{s+1}$, and $w^{(n)} \in H^{ht,s+1}$ and such that*

$$\begin{aligned} X^{(n)} - \alpha \in &\left\{X - \alpha \in F^{s+1} : \left|\left|X - \alpha - \int_0^t A\phi\, d\tau\right|\right|_{F^{s+1}} \leq \left|\left|\int_0^t A\phi\, d\tau\right|\right|_{F^{s+1}}\right\} \\ &\equiv B_{A\phi} \\ \left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}} &\leq N \end{aligned}$$

Then, for small enough $T > 0$, depending only on N and v_0.

$$X^{(n+1)} - \alpha \in B_{A\phi}$$

2. Let $X^{(n)}-\alpha, X^{(n-1)}-\alpha \in F^{s+1}$ *with* $v^{(n)}, v^{(n-1)} \in H^{ht,s+1}$ *and such that*

$$\begin{aligned}\left|\left|v^{(n)}\right|\right|_{H^{ht,s+1}}, \left|\left|v^{(n-1)}\right|\right|_{H^{ht,s+1}} \leq & M\\ ||X^{(n)}-\alpha||_{F^{s+1}}, ||X^{(n-1)}-\alpha||_{F^{s+1}} \leq & M\end{aligned}$$

for some $M>0$*. Then*

$$||X^{(n+1)}-X^{(n)}||_{F^{s+1}} \leq C(M)T^{\varepsilon}\left(\left|\left|v^{(n)}-v^{(n-1)}\right|\right|_{H^{ht,s+1}}+||X^{(n)}-X^{(n-1)}||_{F^{s+1}}\right)$$

for a small enough ε*.*

Proof: Part 1:

We start by remarking that if the norm $\left|\left|\int_0^t A\phi\right|\right|_{F^{s+1}}=0$, then the solution trivializes and $X=\alpha$. It will be enough to show that

$$\left|\left|X^{(n+1)}-\alpha-\int_0^t A\phi\, d\tau\right|\right|_{F^{s+1}} \leq C\left(N, \left|\left|X^{(n)}-\alpha\right|\right|_{F^{s+1}}, v_0\right)T^{\varepsilon},$$

since, as $X^{(n)} \in B_{A\phi}$, this inequality implies that

$$\left|\left|X^{(n+1)}-\alpha-\int_0^t A\phi\, d\tau\right|\right|_{F^{s+1}} \leq C\left(N, v_0\right)T^{\varepsilon}.$$

Here C will be a constant that may depend on N, and v_0 and we recall that C can mean different things in different lines. To conclude, we will only need to take

$$T^{\varepsilon} \leq \frac{\left|\left|\int_0^t A\phi d\tau\right|\right|_{F^{s+1}}}{C\left(N, v_0\right)}.$$

The definition of F^{s+1} and $v^{(n)}$ imply

$$\begin{aligned}&\left|\left|X^{(n+1)}-\alpha-\int_0^t A\phi\right|\right|_{F^{s+1}} \leq \left|\left|\int_0^t A\circ X^{(n)}w^{(n)}\right|\right|_{F^{s+1}}+\left|\left|\int_0^t\left(A\circ X^{(n)}-A\right)\phi d\tau\right|\right|_{F^{s+1}}\\ &\leq \left|\left|\int_0^t A\circ X^{(n)}w^{(n)}\right|\right|_{L^{\infty}_{1/4}H^{s+1}}+\left|\left|\int_0^t A\circ X^{(n)}w^{(n)}\right|\right|_{H^2H^{\gamma}}\\ &+\left|\left|\int_0^t\left(A\circ X^{(n)}-A\right)\phi d\tau\right|\right|_{L^{\infty}_{1/4}H^{s+1}}+\left|\left|\int_0^t\left(A\circ X^{(n)}-A\right)\phi d\tau\right|\right|_{H^2H^{\gamma}}\\ &\equiv I_1+I_2+I_3+I_4.\end{aligned}$$

To estimate I_1 we proceed as follows. Applying Hölder and Minkowski inequalities yields

$$\left|\left|\int_0^t A\circ X^{(n)}w^{(n)}d\tau\right|\right|_{L^{\infty}_{1/4}H^{s+1}} \leq T^{\frac{1}{4}}\left|\left|A\circ X^{(n)}w^{(n)}\right|\right|_{H^0H^{s+1}} \leq T^{\frac{1}{4}}\left|\left|A\circ X^{(n)}\right|\right|_{L^{\infty}H^{s+1}}\left|\left|w^{(n)}\right|\right|_{H^0H^{s+1}},$$

Lemma 3.9 implies

$$\left|\left|A\circ X^{(n)}\right|\right|_{L^{\infty}H^{s+1}} \leq C\left(1+\left|\left|X^{(n)}-\alpha\right|\right|_{F^{s+1}}\right).$$

Therefore

$$I_1 \leq T^{\frac{1}{4}}C\left(N, \left|\left|X^{(n)}-\alpha\right|\right|_{F^{s+1}}, v_0\right).$$

For I_2 we have that

$$I_2 \leq \left|\left|\int_0^t A\circ X^{(n)}w^{(n)}\right|\right|_{H^2H^{\gamma}} \leq \left|\left|A\circ X^{(n)}w^{(n)}\right|\right|_{H^1H^{\gamma}},$$

where we have applied Lemma 3.4 with $\varepsilon = 0$. In addition, applying the third part of Lemma 3.7 and Lemmas 3.3 and 3.4 yields

$$\begin{aligned}\left|\left|A \circ X^{(n)} w^{(n)}\right|\right|_{H^1 H^\gamma} &\leq C(M) \left|\left|\int_0^t \partial_t w^{(n)}\right|\right|_{H^1 H^\gamma} \leq C(M) \left|\left|\int_0^t \partial_t w^{(n)}\right|\right|_{H^{1+\delta-\varepsilon} H^\gamma} \\ &\leq C(M) T^\varepsilon \left|\left|w^{(n)}\right|\right|_{H^{1+\delta} H^\gamma} \leq C(M) T^\varepsilon \left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}.\end{aligned}$$

Here $\varepsilon > 0$, $\varepsilon < \delta$, $\delta < \frac{s-1-\gamma}{2}$. Similar computations show that

$$I_3 \leq T^{\frac{1}{4}} C\left(N, v_0\right).$$

For I_4 by applying the second part of Lemma 3.4 with $\varepsilon = 0$ and the third part of Lemma 3.7 we have that

$$I_4 \leq \left|\left|(A \circ X^{(n)} - A)\phi\right|\right|_{H^1 H^\gamma} \leq C(M) \left|\left|A \circ X^{(n)} - A\right|\right|_{H^1 H^\gamma}$$

To finish the estimate for I_2 we will apply the second part of Lemma 3.4 in such a way that

$$\begin{aligned}\left|\left|A \circ X^{(n)} - A\right|\right|_{H^1 H^\gamma} &= \left|\left|\int_0^t \partial_t (A \circ X^{(n)} - A)\right|\right|_{H^1 H^\gamma} \leq \left|\left|\int_0^t \partial_t (A \circ X^{(n)} - A)\right|\right|_{H^{1+\delta-\varepsilon} H^\gamma} \\ &\leq C T^\varepsilon \left|\left|A \circ X^{(n)} - A\right|\right|_{H^{1+\delta} H^\gamma} \leq C T^\varepsilon \left|\left|X^{(n)} - \alpha\right|\right|_{H^{1+\delta} H^\gamma}.\end{aligned}$$

In order to obtain this last estimate we have applied Lemma 3.9 and part 3 of Lemma 3.7, for $\varepsilon > 0$ and $\varepsilon < \delta < \frac{1}{2}$.

This concludes the proof of part 1 of proposition 5.3.

Part 2:

From (24) we have that

$$\begin{aligned}\left|\left|X^{(n+1)} - X^{(n)}\right|\right|_{F^{s+1}} &= \left|\left|\int_0^t \left(A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)}\right) d\tau\right|\right|_{F^{s+1}} \\ &\leq \left|\left|\int_0^t \left(A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)}\right) d\tau\right|\right|_{L^\infty_{1/4} H^{s+1}} \\ &+ \left|\left|\int_0^t \left(A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)}\right) d\tau\right|\right|_{H^2 H^\gamma} \\ &\equiv I_1 + I_2\end{aligned}$$

In order to bound I_1 we notice that

$$\begin{aligned}&\left|\left|\int_0^t \left(A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)}\right) d\tau\right|\right|_{H^{s+1}} \\ &\leq t^{\frac{1}{2}} \left|\left|A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)} d\tau\right|\right|_{H^0 H^{s+1}}\end{aligned}$$

Therefore

$$\begin{aligned}I_1 &\leq T^{\frac{1}{4}} \left|\left|\left(A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)}\right)\right|\right|_{H^0 H^{s+1}} \\ &\leq T^{\frac{1}{4}} \left|\left|\left(A \circ X^{(n)} - A \circ X^{(n-1)}\right) v^{(n)}\right|\right|_{H^0 H^{s+1}} \\ &+ T^{\frac{1}{4}} \left|\left|\left(v^{(n)} - v^{(n-1)}\right) A \circ X^{(n-1)}\right|\right|_{H^0 H^{s+1}} \\ &\equiv I_{11} + I_{12}\end{aligned}$$

In addition

$$\begin{aligned}
I_{11} &\leq T^{\frac{1}{4}} \left|\left|A \circ X^{(n)} - A \circ X^{(n-1)}\right|\right|_{L^\infty H^{s+1}} \left|\left|v^{(n)}\right|\right|_{H^0 H^{s+1}} \\
&\leq MT^{\frac{1}{4}} \left|\left|RR(X^{(n)}, X^{(n-1)})(X^{(n)} - X^{(n-1)})\right|\right|_{L^\infty H^{s+1}} \\
&\leq MT^{\frac{1}{4}} \left|\left|RR(X^{(n)}, X^{(n-1)})\right|\right|_{L^\infty H^{s+1}} \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^{s+1}} \\
&\leq C(M)T^{\frac{1}{4}} \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^{s+1}} \\
&\leq C(M)T^{\frac{1}{2}} \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty_{1/4} H^{s+1}},
\end{aligned}$$

where we have applied Lemma 3.9 and that H^{s+1} is an algebra as stated in Lemma 3.5. $RR(x, y)$ is defined via

$$RR(x, y) = \frac{A(x) - A(y)}{x - y}.$$

Also we have that

$$\begin{aligned}
I_{12} &\leq CT^{\frac{1}{4}} \left|\left|A \circ X^{(n-1)}\right|\right|_{L^\infty H^{s+1}} \left|\left|v^{(n)} - v^{(n-1)}\right|\right|_{H^0 H^{s+1}} \\
&\leq CT^{\frac{1}{4}} \left(1 + \left|\left|X^{(n-1)} - \alpha\right|\right|_{L^\infty H^{s+1}}\right) \left|\left|v^{(n)} - v^{(n-1)}\right|\right|_{H^0 H^{s+1}} \\
&\leq C(M)T^{\frac{1}{4}} \left|\left|v^{(n)} - v^{(n-1)}\right|\right|_{H^0 H^{s+1}}
\end{aligned}$$

Thus

$$I_1 \leq C(M)T^{\frac{1}{4}} \left(\left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty_{1/4} H^{s+1}} + \left|\left|v^{(n)} - v^{(n-1)}\right|\right|_{H^{ht,s+1}}\right).$$

It remains to bound I_2.

$$I_2 = \left|\left|\int_0^t (A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)}) d\tau\right|\right|_{H^2 H^\gamma}$$

and applying Lemma 3.4 with $\varepsilon = 0$ we have that

$$I_2 \leq \left|\left|A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)}\right|\right|_{H^1 H^\gamma}.$$

We will decompose that term in the following way

$$\begin{aligned}
A \circ X^{(n)} v^{(n)} - A \circ X^{(n-1)} v^{(n-1)} =& (A \circ X^{(n)} - A \circ X^{(n-1)}) w^{(n)} + (A \circ X^{(n)} - A \circ X^{(n-1)}) \phi \\
&+ A \circ X^{(n-1)} (w^{(n)} - w^{(n-1)}).
\end{aligned}$$

Thus

$$I_2 \leq I_{21} + I_{22} + I_{23}$$

with

$$\begin{aligned}
I_{21} &= \left|\left|(A \circ X^{(n)} - A \circ X^{(n-1)}) w^{(n)}\right|\right|_{H^1 H^\gamma} \\
I_{22} &= \left|\left|(A \circ X^{(n)} - A \circ X^{(n-1)}) \phi\right|\right|_{H^1 H^\gamma} \\
I_{23} &= \left|\left|A \circ X^{(n-1)} (w^{(n)} - w^{(n-1)})\right|\right|_{H^1 H^\gamma}.
\end{aligned}$$

For I_{21}, by applying Lemma 3.7 ($1 < \gamma < s - 1$), we have that

$$\begin{aligned}
&\left|\left|(A \circ X^{(n)} - A \circ X^{(n-1)}) w^{(n)}\right|\right|_{H^1 H^\gamma} \leq C \left|\left|A \circ X^{(n)} - A \circ X^{(n-1)}\right|\right|_{H^1 H^\gamma} \left|\left|w^{(n)}\right|\right|_{H^1 H^\gamma} \\
&\leq C(M) \left|\left|A \circ X^{(n)} - A \circ X^{(n-1)}\right|\right|_{H^1 H^\gamma} \leq C(M) \left|\left|RR(X^{(n)}, X^{(n-1)})(X^{(n)} - X^{(n-1)})\right|\right|_{H^1 H^\gamma}.
\end{aligned}$$

Then the third part of Lemma 3.7 yields

$$\begin{aligned} I_{21} \leq & C(M) \left|\left| X^{(n)} - X^{(n-1)} \right|\right|_{H^1 H^\gamma} \leq \left|\left| \int_0^t (\partial_t \left(X^{(n)} - X^{(n-1)} \right) d\tau) \right|\right|_{H^{1+\delta-\varepsilon} H^\gamma} \\ & \leq C(M) T^\varepsilon \left|\left| X^{(n)} - X^{(n-1)} \right|\right|_{H^{1+\delta} H^\gamma}, \end{aligned}$$

where we have applied Lemma 3.3 with $\varepsilon > 0$, $\delta > \varepsilon$ and $\delta < \frac{1}{2}$.

The term I_{22} is quite similar to I_{21}. The only difference is that we have to apply the third part of Lemma 3.7 to obtain that

$$\left|\left| (A \circ X^{(n)} - A \circ X^{(n-1)}) \phi \right|\right|_{H^1 H^\gamma} \leq C(M) \left|\left| A \circ X^{(n)} - A \circ X^{(n-1)} \right|\right|_{H^1 H^\gamma}.$$

Finally

$$I_{23} \leq C(M) \left|\left| w^{(n)} - w^{(n-1)} \right|\right|_{H^1 H^\gamma},$$

by applying the third part of Lemma 3.7. Also we can compute

$$\begin{aligned} I_{23} \leq & C(M) \left|\left| w^{(n)} - w^{(n-1)} \right|\right|_{H^{1+\delta-\varepsilon} H^\gamma} \leq C(M) T^\varepsilon \left|\left| w^{(n)} - w^{(n-1)} \right|\right|_{H^{1+\delta} H^\gamma} \\ \leq & C(M) T^\varepsilon \left|\left| w^{(n)} - w^{(n-1)} \right|\right|_{H^{ht, s+1}}, \end{aligned}$$

for $\varepsilon > 0$, $\delta > \varepsilon$ and $\delta < \frac{s-1-\gamma}{2}$.

This concludes the proof of part 2 of proposition 5.3. Therefore proposition 5.3 is shown. □

Proposition 5.4 *1. Let $X^{(n)} - \alpha \in F^{s+1}$, $q_w^{(n)} \in H_{pr}^{ht,\, s}$ and $w^{(n)} \in H^{ht, s+1}$, and such that*

$$\begin{aligned} & \left|\left| X^{(n)} - \alpha \right|\right|_{F^{s+1}}, \leq N \\ & (w^{(n)}, q_w^{(n)}) \in \big\{ (w, q) \in H^{ht,\, s+1} \times H_{pr}^{ht,s} \,:\, w|_{t=0} = 0, \partial_t w|_{t=0} = 0, \\ & \qquad ||(w, q) - L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)||_{H^{ht,s+1} \times H_{pr}^{ht,s}} \\ & \qquad \leq ||L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)||_{H^{ht,s+1} \times H_{pr}^{ht,s}} \big\} \\ & \qquad \equiv B_{L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)}. \end{aligned}$$

Then

$$(w^{(n+1)}, q_w^{(n+1)}) \in B_{L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)}.$$

2. Let $X^{(n)} - \alpha, X^{(n-1)} - \alpha \in F^{(s+1)}$ and $w^{(n)}, w^{(n-1)} \in H^{ht,\, s+1}$, with $w^{(n)}|_{t=0} = w^{(n-1)}|_{t=0} = \partial_t w^{(n)}|_{t=0} = \partial_t w^{(n-1)}|_{t=0} = 0$ and such that

$$\begin{aligned} \left|\left| X^{(n)} - \alpha \right|\right|_{F^{s+1}}, \left|\left| X^{(n-1)} - \alpha \right|\right|_{F^{s+1}} & \leq M \\ \left|\left| v^{(n)} \right|\right|_{H^{ht,s+1}}, \left|\left| v^{(n-1)} \right|\right|_{H^{ht,s+1}} & \leq M \\ ||q_w^{(n)}||_{H_{pr}^{ht,s}}, ||q_w^{(n-1)}||_{H_{pr}^{ht,s}} & \leq M \end{aligned}$$

Then

$$\begin{aligned} & \left|\left| w^{(n+1)} - w^{(n)} \right|\right|_{H^{ht,s+1}} + ||q^{(n+1)} - q^{(n)}||_{H_{pr}^{ht,s}} \\ & \leq C(M) T^\varepsilon \left(\left|\left| w^{(n)} - w^{(n-1)} \right|\right|_{H^{ht,s+1}} + \left|\left| X^{(n)} - X^{(n-1)} \right|\right|_{F^{s+1}} + ||q^{(n)} - q^{(n-1)}||_{H_{pr}^{ht,s}} \right), \end{aligned}$$

for ε small enough.

Proof of part 1:

By applying Theorem 4.5 we see it is enough to prove that

$$\left|\left|f^{(n)}\right|\right|_{H^{ht,s-1}} \leq C\left(N, \left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}, ||q_w^{(n)}||_{H^{ht,s}_{pr}}\right)T^{\varepsilon}, \tag{61}$$

$$\left|\left|\overline{g}^{(n)}\right|\right|_{\overline{H}^{ht,s}} \leq C\left(N, \left|\left|w^{(n)}\right|\right|_{H^{ht,s}}, ||q_w^{(n)}||_{H^{ht,s}_{pr}}\right)T^{\varepsilon}, \tag{62}$$

$$\left|h^{(n)}\right|_{H^{ht,s-\frac{1}{2}}} \leq C\left(N, \left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}, ||q_w^{(n)}||_{H^{ht,s}_{pr}}\right)T^{\varepsilon} \tag{63}$$

for all $(w^{(n)}, q_w^{(n)})$ in $B_{L^{-1}(f_\phi,\overline{g}_\phi,h_\phi)}$. Indeed

$$\begin{aligned}
&\left|\left|(w^{(n+1)}, q_w^{(n+1)}) - L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)\right|\right|_{H^{ht,s+1}\times H^{ht,s}_{pr}} \leq C\left|\left|L^{-1}(f^{(n)}, \overline{g}^{(n)}, h^{(n)})\right|\right|_{H^{ht,s+1}\times H^{ht,s}_{pr}} \\
&\leq C\left(\left|\left|f^{(n)}\right|\right|_{H^{ht,s-1}} + ||\overline{g}^{(n)}||_{\overline{H}^{ht,s}} + \left|h^{(n)}\right|_{H^{ht,s-\frac{1}{2}}}\right) \\
&\leq C\left(N, \left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}, ||q_w^{(n)}||_{H^{ht,s}_{pr}}\right)T^{\varepsilon} \\
&\leq C\left(N, ||L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)||_{H^{ht,s+1}\times H^{ht,s}_{pr}}\right)T^{\varepsilon}
\end{aligned}$$

(here we recall that C can mean different things in different lines) and we just need to take

$$T^{\varepsilon} \leq \frac{||L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)||_{H^{ht,s+1}}}{C\left(N, ||L^{-1}(f_\phi, \overline{g}_\phi, h_\phi)||_{H^{ht,s+1}}\right)}.$$

We split the proof in three parts, corresponding with the functions $f^{(n)}$, $\overline{g}^{(n)}$ and $h^{(n)}$:

P1.1. Estimate for $f^{(n)}$:

In this section we have to deal with $f^{(n)}$ to estimate

$$\left|\left|f^{(n)}\right|\right|_{H^{ht,s-1}} = \left|\left|f^{(n)}\right|\right|_{L^2H^{s-1}} + \left|\left|f^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2}.$$

We will gather terms by writing $f^{(n)} = f_w^{(n)} + f_\phi^{(n)} + f_q^{(n)}$ as follows,

$$f_w^{(n)} = Q^2 \circ X^{(n)}\zeta^{(n)}\partial\left(\zeta^{(n)}\partial w^{(n)}\right) - Q^2\Delta w^{(n)},$$

$$f_\phi^{(n)} = Q^2 \circ X^{(n)}\zeta^{(n)}\partial\left(\zeta^{(n)}\partial\phi\right) - Q^2\Delta\phi, \tag{64}$$

and

$$f_q^{(n)} = -A \circ X^{(n)}\zeta^{(n)}\partial q^{(n)} + A\partial q^{(n)}.$$

Above we removed subscripts to alleviate notation. Next we also ignore the superscripts for the same reason. We firstly bound $||\cdot||_{L^2H^{s-1}}$:

$$\begin{aligned}
||f_w||_{L^2H^{s-1}} \leq &\left|\left|(Q^2\circ X - Q^2)\zeta\partial(\zeta\partial w)\right|\right|_{L^2H^{s-1}} + \left|\left|Q^2(\zeta - \mathbb{I})\partial(\zeta\partial w)\right|\right|_{L^2H^{s-1}} \\
&+ \left|\left|Q^2\partial((\zeta - \mathbb{I})\partial w)\right|\right|_{L^2H^{s-1}} \equiv I_1 + I_2 + I_3.
\end{aligned}$$

We can deal with these three terms as follows. For I_1 one can get

$$\begin{aligned}
I_1 \leq &\left|\left|(Q^2\circ X - Q^2)\zeta\partial\zeta\partial w\right|\right|_{L^2H^{s-1}} + \left|\left|(Q^2\circ X - Q^2)\zeta\zeta\partial^2 w\right|\right|_{L^2H^{s-1}} \\
\leq &\left|\left|Q^2\circ X - Q^2\right|\right|_{L^\infty H^{s-1}} ||\zeta||_{L^\infty H^{s-1}} ||\partial\zeta||_{L^\infty H^{s-1}} ||\partial w||_{L^2H^{s-1}} \\
&+ \left|\left|Q^2\circ X - Q^2\right|\right|_{L^\infty H^{s-1}} ||\zeta||^2_{L^\infty H^{s-1}} \left|\left|\partial^2 w\right|\right|_{L^2H^{s-1}},
\end{aligned}$$

and therefore

$$\begin{aligned}
I_1 \leq &C\left|\left|Q^2\circ X - Q^2\right|\right|_{L^\infty H^{s-1}} ||X||^2_{L^\infty H^{s+1}} ||w||_{H^{ht,s+1}} \\
\leq &C(M)\,||X - \alpha||_{L^\infty H^{s-1}} ||w||_{H^{ht,s+1}} \leq C(M)T^{\frac{1}{4}}.
\end{aligned}$$

Also

$$\begin{aligned}I_2 \leq & \left|\left|Q^2(\zeta-\mathbb{I})\partial\zeta\partial w\right|\right|_{L^2H^{s-1}} + \left|\left|Q^2(\zeta-\mathbb{I})\zeta\partial^2 w\right|\right|_{L^2H^{s-1}}\\ \leq & C\,||\zeta-\mathbb{I}||_{L^\infty H^{s-1}}\left(||\partial\zeta||_{L^\infty H^{s-1}}||\partial w||_{L^2H^{s-1}} + ||\zeta||_{L^\infty H^{s-1}}\left|\left|\partial^2 w\right|\right|_{L^2H^{s-1}}\right)\\ \leq & C(M)\,||\zeta-\mathbb{I}||_{L^\infty H^{s-1}}||w||_{H^{ht,s+1}}.\end{aligned}$$

Identity

$$\zeta-\mathbb{I} = (\nabla X)^{-1}(\mathbb{I}-\nabla X) = (\nabla X)^{-1}\nabla(\alpha - X),$$

allows us to get

$$||\zeta-\mathbb{I}||_{L^\infty H^s} \leq \left|\left|(\nabla X)^{-1}\right|\right|_{L^\infty H^s}||X-\alpha||_{L^\infty H^{s+1}} \leq C(M)T^{\frac14}||X-\alpha||_{L^\infty_{1/4}H^{s+1}},$$

thus

$$I_2 \leq C(M)T^{\frac14}.$$

It remains I_3 for which we compute

$$\begin{aligned}I_3 = & \left|\left|Q^2\partial((\zeta-\mathbb{I})\partial w)\right|\right|_{L^2H^{s-1}} \leq \left|\left|Q^2\partial\zeta\partial w\right|\right|_{L^2H^{s-1}} + \left|\left|Q^2(\zeta-\mathbb{I})\partial^2 w\right|\right|_{L^2H^{s-1}}\\ \leq & C(||\partial\zeta||_{L^\infty H^{s-1}}||w||_{L^2H^s} + ||\zeta-\mathbb{I}||_{L^\infty H^{s-1}}||w||_{L^2H^{s+1}})\\ \leq & C\,||\zeta-\mathbb{I}||_{L^\infty H^s}||w||_{H^{ht,s+1}} \leq C(M)T^{\frac14}.\end{aligned}$$

We are done with $||f_w||_{L^2H^{s-1}}$. At this point it is easy to check that an analogous procedure yields

$$||f_\phi||_{L^2H^{s-1}} \leq C(M)T^{\frac14}||\phi||_{H^{ht,s+1}} \leq C(M)T^{\frac14}.$$

Next we deal with $||f_q||_{L^2H^{s-1}}$. Indeed,

$$\begin{aligned}||f_q||_{L^2H^{s-1}} \leq & ||A\circ X(\zeta-\mathbb{I})\partial q||_{L^2H^{s-1}} + ||(A - A\circ X)\partial q||_{L^2H^{s-1}}\\ \leq & (||A\circ X||_{L^\infty H^{s-1}}||\zeta-\mathbb{I}||_{L^\infty H^{s-1}} + ||A\circ X - A||_{L^\infty H^{s-1}})\,||\partial q||_{L^2H^{s-1}}\\ \leq & C(M)T^{\frac14}||q||_{H^{ht,s}_{pr}} \leq C(M)T^{\frac14}.\end{aligned}$$

Therefore we are done with $||f||_{L^2H^{s-1}}$.

Using the same splitting we now deal with $||f||_{H^{\frac{s-1}{2}}L^2}$:

$$\begin{aligned}||f_w||_{H^{\frac{s-1}{2}}L^2} \leq & \left|\left|(Q^2\circ X - Q^2)\zeta\partial(\zeta\partial w)\right|\right|_{H^{\frac{s-1}{2}}L^2} + \left|\left|Q^2(\zeta-\mathbb{I})\partial(\zeta\partial w)\right|\right|_{H^{\frac{s-1}{2}}L^2}\\ & + \left|\left|Q^2\partial((\zeta-\mathbb{I})\partial w)\right|\right|_{H^{\frac{s-1}{2}}L^2} \equiv I_4 + I_5 + I_6.\end{aligned}$$

We bound I_4 by using lemma 3.7 as follows,

$$I_4 \leq \left|\left|Q^2\circ X - Q^2\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}(||\zeta||_{H^{\frac{s-1}{2}}H^{1+\delta}}+1)(||\partial\zeta\partial w||_{H^{\frac{s-1}{2}}L^2} + \left|\left|\zeta\partial^2 w\right|\right|_{H^{\frac{s-1}{2}}L^2}).$$

Using lemma 3.6, with $\frac{1}{q}=\varepsilon$, yields

$$\begin{aligned}||\partial\zeta\partial w||_{H^{\frac{s-1}{2}}L^2} \leq & ||\partial\zeta||_{H^{\frac{s-1}{2}}H^\varepsilon}||\partial w||_{H^{\frac{s-1}{2}}H^{1-\varepsilon}}\\ \leq & C(M)\,||X-\alpha||_{H^{\frac{s-1}{2}}H^{2+\varepsilon}}||w||_{H^{\frac{s-1}{2}}H^{2-\varepsilon}} \leq C(M),\end{aligned}$$

for $\varepsilon > 0$ and small enough. Also

$$\left|\left|\zeta\partial^2 w\right|\right|_{H^{\frac{s-1}{2}}L^2} \leq (||\zeta-\mathbb{I}||_{H^{\frac{s-1}{2}}H^{1+\delta}} + 1)\left|\left|\partial^2 w\right|\right|_{H^{\frac{s-1}{2}}L^2} \leq C(M).$$

Finally,

$$\begin{aligned}\left|\left|Q^2\circ X - Q^2\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}} \leq & C(M)\,||X-\alpha||_{H^{\frac{s-1}{2}}H^{1+\delta}} \leq C(M)\left|\left|\partial_t\int_0^t(X-\alpha)\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\\ \leq & C(M)\left|\left|\int_0^t(X-\alpha)\right|\right|_{H^{\frac{s-1}{2}+\varepsilon+1-\varepsilon}H^{1+\delta}} \leq C(M)T^\varepsilon\,||X-\alpha||_{H^{\frac{s-1}{2}+\varepsilon}H^{1+\delta}}\\ \leq & C(M)T^\varepsilon\,||X-\alpha||_{F^{s+1}} \leq C(M)T^\varepsilon,\end{aligned}$$

where we have used lemma 3.9 and the second part of lemma 3.4 with $\frac{1}{2} < \frac{s-1}{2} + \varepsilon < 1$. With this last inequality we conclude the estimate for I_4.

For I_5 we have that

$$I_5 \leq ||Q^2||_{H^{1+\delta}} ||\zeta - \mathbb{I}||_{H^{\frac{s-1}{2}}H^{1+\delta}} ||\partial(\zeta\partial w)||_{H^{\frac{s-1}{2}}L^2}.$$

The norm $||\partial(\zeta\partial w)||_{H^{\frac{s-1}{2}}L^2}$ has already been estimated for I_4. Using lemma 3.8 and the second part of lemma 3.4 we find that

$$\begin{aligned} ||\zeta - \mathbb{I}||_{H^{\frac{s-1}{2}}H^{1+\delta}} &\leq CT^{\varepsilon} ||\zeta - \mathbb{I}||_{H^{\frac{s-1}{2}+\varepsilon}H^{1+\delta}} \leq C(M)T^{\varepsilon} ||X - \alpha||_{H^{\frac{s-1}{2}+\varepsilon}H^{2+\delta}} \\ &\leq C(M)T^{\varepsilon}. \end{aligned}$$

We are done with I_5. For I_6 we have that

$$I_6 \leq \left|\left|Q^2\partial\zeta\partial w\right|\right|_{H^{\frac{s-1}{2}}L^2} + \left|\left|Q^2(\zeta - \mathbb{I})\partial^2 w\right|\right|_{H^{\frac{s-1}{2}}L^2}.$$

Both terms can be handled as before. In fact

$$\begin{aligned} \left|\left|Q^2(\zeta - \mathbb{I})\partial^2 w\right|\right|_{H^{\frac{s-1}{2}}L^2} \leq& ||Q^2||_{H^{1+\delta}} ||\zeta - \mathbb{I}||_{H^{\frac{s-1}{2}}H^{1+\delta}} \left|\left|\partial^2 w\right|\right|_{H^{\frac{s-1}{2}}L^2} \\ \leq& C(M)T^{\varepsilon} ||w||_{H^{ht,s+1}} \leq C(M)T^{\varepsilon}, \end{aligned}$$

and

$$\begin{aligned} \left|\left|Q^2\partial\zeta\partial w\right|\right|_{H^{\frac{s-1}{2}}L^2} &\leq ||Q^2||_{H^{1+\delta}} ||\partial\zeta||_{H^{\frac{s-1}{2}}H^{\varepsilon}} ||\partial w||_{H^{\frac{s-1}{2}}H^{1-\varepsilon}} \\ &\leq C ||\zeta - \mathbb{I}||_{H^{\frac{s-1}{2}}H^{1+\epsilon}} ||w||_{H^{\frac{s-1}{2}}H^{2-\varepsilon}} \leq C(M)T^{\varepsilon} ||w||_{H^{ht,s+1}} \\ &\leq C(M)T^{\varepsilon}, \end{aligned}$$

for $\varepsilon > 0$ and small enough. We are done with I_6 and therefore with $||f_w||_{H^{\frac{s-1}{2}}L^2}$. The same procedure allows us to get

$$||f_\phi||_{H^{\frac{s-1}{2}}L^2} \leq C(M)T^{\epsilon}.$$

The estimate of $||f_q||_{H^{\frac{s-1}{2}}L^2}$ is obtained as follows

$$\begin{aligned} ||f_q||_{H^{\frac{s-1}{2}}L^2} \leq& ||(A \circ X - A)\zeta\partial q||_{H^{\frac{s-1}{2}}H^{1+\delta}} + ||A(\zeta - \mathbb{I})\partial q||_{H^{\frac{s-1}{2}}H^{1+\delta}} \\ \leq& ||A \circ X - A||_{H^{\frac{s-1}{2}}H^{1+\delta}} (||\zeta||_{H^{\frac{s-1}{2}}H^{1+\delta}} + 1)(||\partial q||_{H^{\frac{s-1}{2}}L^2} + 1) \\ &+ ||A||_{H^{1+\delta}} ||\zeta - \mathbb{I}||_{H^{\frac{s-1}{2}}H^{1+\delta}} (||\partial q||_{H^{\frac{s-1}{2}}L^2} + 1) \\ \leq& C(M)T^{\varepsilon}(\|q\|_{H^{ht,s}_{pr}} + 1) \leq C(M)T^{\varepsilon}. \end{aligned}$$

This finishes the bounds for $||f||_{H^{\frac{s-1}{2}}L^2}$ and therefore we are done with $||f||_{H^{ht,s-1}}$.

P1.2. Estimate for $\overline{g}^{(n)}$:

We recall that

$$\left|\left|\overline{g}^{(n)}\right|\right|_{\overline{H}^{ht,s}} = \left|\left|\overline{g}^{(n)}\right|\right|_{H^0H^s} + \left|\left|\overline{g}^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}.$$

We will first estimate the H^0H^s-norm and after that the $H^{\frac{s+1}{2}}H^{-1}$-norm. We will split $\overline{g}^{(n)}$ in the following terms,

$$\begin{aligned} \overline{g}^{(n)} =& - Tr\left(\nabla v^{(n)}\zeta^{(n)}A \circ X^{(n)}\right) + Tr\left(\nabla v^{(n)}A\right) \\ &+ Tr\left(\nabla\phi\zeta_\phi A_\phi\right) - Tr\left(\nabla\phi A\right) \\ =& - Tr\left(\nabla w^{(n)}\zeta^{(n)}A \circ X^{(n)}\right) - Tr\left(\nabla\phi(\zeta^{(n)}A \circ X^{(n)} - \zeta_\phi A_\phi)\right) \\ &+ Tr\left(\nabla w^{(n)}A\right) + Tr\left(\nabla\phi\zeta_\phi A_\phi\right) \\ =& - Tr\left(\nabla w^{(n)}\left(\zeta^{(n)} - \mathbb{I}\right)A \circ X^{(n)}\right) - Tr\left(\nabla\phi\left(\zeta^{(n)} - \zeta_\phi\right)A \circ X^{(n)}\right) \\ &+ Tr\left(\nabla w^{(n)}\left(A - A \circ X^{(n)}\right)\right) + Tr\left(\nabla\phi\zeta_\phi\left(A_\phi - A \circ X^{(n)}\right)\right), \end{aligned}$$

From the partition of $\overline{g}^{(n)}$ we have that

$$\begin{aligned}\left|\left|\overline{g}^{(n)}\right|\right|_{H^0H^s} =& \left|\left|\nabla w^{(n)}\left(\zeta^{(n)}-\mathbb{I}\right)A\circ X^{(n)}\right|\right|_{H^0H^s}+\left|\left|\nabla\phi\left(\zeta^{(n)}-\zeta_\phi\right)A\circ X^{(n)}\right|\right|_{H^0H^s}\\ &+\left|\left|\nabla w^{(n)}\left(A-A\circ X^{(n)}\right)\right|\right|_{H^0H^s}+\left|\left|\nabla\phi\left(A_\phi-A\circ X^{(n)}\right)\right|\right|_{H^0H^s}\\ &\equiv I_1+I_2+I_3+I_4.\end{aligned}$$

Since H^s is an algebra for $s>1$ as stated in lemma 3.5 we have that

$$I_1\leq\left|\left|\zeta^{(n)}-\mathbb{I}\right|\right|_{L^\infty H^s}\left|\left|A\circ X^{(n)}\right|\right|_{L^\infty H^s}\left|\left|\nabla w^{(n)}\right|\right|_{H^0H^s}.$$

Therefore, applying lemma 3.9 yields

$$I_1\leq C(N)\left|\left|\zeta^{(n)}-\mathbb{I}\right|\right|_{L^\infty H^s}\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}.$$

In order to bound the $L^\infty H^s$-norm of $\zeta^{(n)}-\mathbb{I}$ we proceed as follows. First we notice that

$$\begin{aligned}\zeta^{(n)}-\mathbb{I}=&\left(\nabla X^{(n)}\right)^{-1}-\mathbb{I}=\left(\nabla X^{(n)}\right)^{-1}\left(\mathbb{I}-\nabla X^{(n)}\right)\\ =&\left(\nabla X^{(n)}\right)^{-1}\nabla\left(\alpha-X^{(n)}\right).\end{aligned}$$

Therefore

$$\begin{aligned}\left|\left|\zeta^{(n)}-\mathbb{I}\right|\right|_{L^\infty H^s}\leq&\left|\left|\left(\nabla X^{(n)}\right)^{-1}\right|\right|_{L^\infty H^s}\left|\left|X^{(n)}-\alpha\right|\right|_{L^\infty H^{s+1}}\\ \leq& C(N)T^{\frac{1}{4}}\left|\left|X^{(n)}-\alpha\right|\right|_{L^\infty_{1/4}H^{s+1}}.\end{aligned}\tag{65}$$

where we have used lemma 3.9. Thus

$$I_1\leq C(N)\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}T^{\frac{1}{4}}.$$

Similarly we obtain that

$$I_2\leq C(N,v_0)T^{\frac{1}{4}}.$$

In addition, by using the differentiability of A and lemma 3.9 it is easy to check that

$$\begin{aligned}I_3\leq& C(N)T^{\frac{1}{4}}\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}\\ I_4\leq& C(N,v_0)T^{\frac{1}{4}}.\end{aligned}$$

Then we have proved that

$$\left|\left|\overline{g}^{(n)}\right|\right|_{H^0H^s}\leq C\left(N,v_0,\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}\right)T^{\frac{1}{4}}.$$

To estimate the $H^{\frac{s+1}{2}}H^{-1}$-norm of $\overline{g}^{(n)}$, we split in a different way

$$\begin{aligned}\overline{g}^{(n)}=&-Tr\left(\nabla w^{(n)}\zeta^{(n)}A\circ X^{(n)}\right)+Tr\left(\nabla w^{(n)}A\right)\\ &-Tr\left(\nabla\phi(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right)+Tr\left(\nabla\phi\zeta_\phi(A_\phi-A\circ X^{(n)})\right)\\ &\equiv-\overline{g}^{(n)}_w-\overline{g}^{(n)}_\phi,\end{aligned}$$

where

$$\begin{aligned}\overline{g}^{(n)}_w=&Tr\left(\nabla w^{(n)}\zeta^{(n)}A\circ X^{(n)}\right)-Tr\left(\nabla w^{(n)}A\right)\\ =&Tr\left(\nabla w^{(n)}(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right)+Tr\left(\nabla w^{(n)}\zeta_\phi(A\circ X^{(n)}-A_\phi)\right)\\ &+Tr\left(\nabla w^{(n)}\zeta_\phi(A_\phi-A)\right)+Tr\left(\nabla w^{(n)}(\zeta_\phi-\mathbb{I})A\right).\end{aligned}$$

and

$$\overline{g}_\phi^{(n)} = Tr\left(\nabla\phi(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right) - Tr\left(\nabla\phi\zeta_\phi(A_\phi - A\circ X^{(n)})\right)$$

First we will bound $\overline{g}_w^{(n)}$ and then we will do the same with $\overline{g}_\phi^{(n)}$. Taking $H^{\frac{s+1}{2}}H^{-1}$-norms yields,

$$\begin{aligned}\left|\left|\overline{g}_w^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}} \leq & \left|\left|\nabla w^{(n)}(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\\ & + \left|\left|\nabla w^{(n)}\zeta_\phi(A\circ X^{(n)}-A_\phi)\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\\ & + \left|\left|\nabla w^{(n)}\zeta_\phi(A_\phi - A)\right|\right|_{H^{\frac{s+1}{2}}H^{-1}} + \left|\left|\nabla w^{(n)}(\zeta_\phi-\mathbb{I})A\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\\ \equiv & I_1+I_2+I_3+I_4. \end{aligned} \tag{66}$$

For I_1 we have that

$$\begin{aligned} I_1 = & \left|\left|\nabla w^{(n)}(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}} = \left|\left|\int_0^t \partial_t\left(\nabla w^{(n)}(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right)\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\\ \leq & \left|\left|\int_0^t \partial_t\nabla w^{(n)}(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}} + \left|\left|\int_0^t \nabla w^{(n)}\partial_t(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\\ + & \left|\left|\int_0^t \nabla w^{(n)}(\zeta^{(n)}-\zeta_\phi)\partial_t A\circ X^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\\ \equiv & I_{11}+I_{12}+I_{13}. \end{aligned}$$

And we bound I_{11}, I_{12} and I_{13} as follows

$$I_{11} \leq \left|\left|\partial_t\nabla w^{(n)}(\zeta^{(n)}-\zeta_\phi)A\circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}},$$

since $0<\frac{s-1}{2}<1$ and we can apply part 2 of lemma 3.4 with $\varepsilon=0$. Moreover, applying part 3 of lemma 3.7 we obtain that

$$I_{11} \leq C(M)\left|\left|\nabla\partial_t w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}\left(1+\left|\left|A\circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\right)\left|\left|\zeta^{(n)}-\zeta_\phi\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}.$$

In addition

$$\begin{aligned} \left|\left|\zeta^{(n)}-\zeta_\phi\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}} \leq & \left|\left|\partial_t\int_0^t \zeta^{(n)}-\zeta_\phi\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\\ \leq C & \left|\left|\int_0^t \zeta^{(n)}-\zeta_\phi\right|\right|_{H^{\frac{s-1}{2}+\varepsilon+1-\varepsilon}H^{1+\delta}} \leq CT^\varepsilon\left|\left|\zeta^{(n)}-\zeta_\phi\right|\right|_{H^{\frac{s-1}{2}+\varepsilon}H^{1+\delta}} \end{aligned}$$

by the second part of lemma 3.4, for $0<\frac{s-1}{2}+\varepsilon<1$. Then lemmas 3.8, 3.9 and 3.3 imply the estimate we need.

For I_{12} we have that, applying part 4 of lemma 3.5 and part 3 of lemma 3.7,

$$I_{12} \leq C(M)\left|\left|\nabla w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^1}\left(1+\left|\left|A\circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\right)\left|\left|\partial_t(\zeta^{(n)}-\zeta_\phi)\right|\right|_{H^{\frac{s-1}{2}}H^0}.$$

By lemmas 3.3 and 3.9 we obtain

$$I_{12} \leq C(M)\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}\left|\left|\partial_t(\zeta^{(n)}-\zeta_\phi)\right|\right|_{H^{\frac{s-1}{2}}H^0}.$$

Again we can apply lemma 3.4 to get

$$\begin{aligned} I_{12} \leq & C(M)\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}T^\varepsilon\left|\left|\partial_t(\zeta^{(n)}-\zeta_\phi)\right|\right|_{H^{\frac{s-1}{2}+\varepsilon}H^0}\\ \leq & C(M)\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}T^\varepsilon\left|\left|\zeta^{(n)}-\zeta_\phi\right|\right|_{H^{\frac{s+1}{2}+\varepsilon}H^0} \end{aligned}$$

for $0<\frac{s-1}{2}+\varepsilon<1$. Therefore lemmas 3.9 and 3.8 yield a suitable estimate for I_{12}. The term I_{13} is even easier once we have lemmas 3.9, 3.8 and 3.3.

Next we bound the second term in (66),

$$\left|\left|\nabla w^{(n)}\zeta_\phi(A\circ X^{(n)}-A_\phi)\right|\right|_{H^{\frac{s+1}{2}}H^{-1}} \leq \left|\left|\nabla w^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\left(1+||\zeta_\phi||_{H^{\frac{s+1}{2}}H^{1+\delta}}\right) \\ \times\left|\left|A\circ X^{(n)}-A_\phi\right|\right|_{H^{\frac{s+1}{2}}H^{1+\delta}}$$

But

$$\begin{aligned}\left|\left|A\circ X^{(n)}-A_\phi\right|\right|_{H^{\frac{s+1}{2}}H^{1}} &= \left|\left|\int_0^t \partial_t(A\circ X^{(n)}-A_\phi)\right|\right|_{H^{\frac{s-1}{2}+1}H^{1+\delta}}\\ &\leq CT^{\varepsilon}\left|\left|\partial_t(A\circ X^{(n)}-A_\phi)\right|\right|_{H^{\frac{s-1}{2}+\varepsilon}H^{1+\delta}},\end{aligned}$$

for $0<\frac{s-1}{2}+\varepsilon<1$. To finish the estimate for I_{12} we just need to apply lemmas 3.8 and 3.9.

The way to bound the third and fourth term in (66) is somehow different. For example

$$\begin{aligned}I_3 &= \left|\left|\nabla w^{(n)}\zeta_\phi(A_\phi-A)\right|\right|_{H^{\frac{s+1}{2}}H^{-1}}\\ &\leq \left|\left|\partial_t\nabla w^{(n)}\zeta_\phi(A_\phi-A)\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}+\left|\left|\nabla w^{(n)}\partial_t\zeta_\phi(A_\phi-A)\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\qquad +\left|\left|\nabla w^{(n)}\zeta_\phi\partial_t A_\phi\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}.\end{aligned}$$

This splitting allows us to find

$$\begin{aligned}I_3 \leq&\left|\left|\partial_t\nabla w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}(1+||\zeta_\phi||_{H^{\frac{s-1}{2}}H^{1+\delta}})\,||A_\phi-A||_{H^{\frac{s-1}{2}}H^{1+\delta}}\\ &+\left|\left|\nabla w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}(1+||\partial_t\zeta_\phi||_{H^{\frac{s-1}{2}}H^{1+\delta}})\,||A_\phi-A||_{H^{\frac{s-1}{2}}H^{1+\delta}}\\ &+\left|\left|\nabla w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}(1+||\zeta_\phi||_{H^{\frac{s-1}{2}}H^{1+\delta}})(||\partial_t A_\phi||_{H^{\frac{s-1}{2}}H^{1+\delta}}+1).\end{aligned}$$

Here the most difficult term to bound is the last one. We proceed as follows

$$\left|\left|\nabla w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}\leq\left|\left|w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{0}}\leq CT^{\varepsilon}\left|\left|w^{(n)}\right|\right|_{H^{\frac{s-1}{2}+\varepsilon}H^{0}}\leq CT^{\varepsilon}\left|\left|w^{(n)}\right|\right|_{H^{ht,s+1}}.$$

The same argument applies to I_4.

The estimate of $||\overline{g}_\phi||_{H^{\frac{s+1}{2}}H^{-1}}$ follows similar (simpler) steps.

P1.3. Estimate for $h^{(n)}$:

We will show the appropriate estimate for $h^{(n)}$. We decompose $h^{(n)}=h_v^{(n)}+h_{v*}^{(n)}+h_q^{(n)}$ given by

$$\begin{aligned}h_v^{(n)} &= (\nabla v^{(n)}\zeta^{(n)}\nabla_J X^{(n)}-\nabla v^{(n)})n_0,\\ h_{v*}^{(n)} &= ((\nabla v^{(n)}\zeta^{(n)}A\circ X^{(n)})^*A^{-1}\circ X^{(n)}\nabla_J X^{(n)}-(\nabla v^{(n)}A)^*A^{-1})n_0,\\ h_q^{(n)} &= (-q^{(n)}A^{-1}\circ X^{(n)}\nabla_J X^{(n)}+q^{(n)}A^{-1})n_0.\end{aligned}\tag{67}$$

As before, we ignore the superscripts for simplicity. We deal first with the $|\cdot|_{L^2H^{s-\frac{1}{2}}}$. Then

$$|h_v|_{L^2H^{s-\frac{1}{2}}}\leq|\nabla v(\zeta-\mathbb{I})\nabla_J X|_{L^2H^{s-\frac{1}{2}}}+|\nabla v(\nabla_J X-\mathbb{I})|_{L^2H^{s-\frac{1}{2}}}\equiv I_1+I_2.$$

For I_1 we find

$$\begin{aligned}I_1 &\leq C\,|\nabla v|_{L^2H^{s-\frac{1}{2}}}\,|\zeta-\mathbb{I}|_{L^\infty H^{s-\frac{1}{2}}}\,|\nabla X|_{L^\infty H^{s-\frac{1}{2}}}\\ &\leq C\,||v||_{L^2H^{s+1}}\,||\zeta-\mathbb{I}||_{L^\infty H^{s}}\,||X||_{L^\infty H^{s+1}}\\ &\leq C(M)\,||X-\alpha||_{L^\infty H^{s+1}}\leq C(M)T^{\frac{1}{4}}\,||X-\alpha||_{L^\infty_{1/4}H^{s+1}}\leq C(M)T^{\frac{1}{4}}.\end{aligned}$$

For I_2 the computation is analogous:

$$\begin{aligned}I_2 &\leq C\,|\nabla v|_{L^2H^{s-\frac{1}{2}}}\,|\nabla X-\mathbb{I}|_{L^\infty H^{s-\frac{1}{2}}}\leq C(M)\,||X-\alpha||_{L^\infty H^{s+1}}\\ &\leq C(M)T^{\frac{1}{4}}\,||X-\alpha||_{L^\infty_{1/4}H^{s+1}}\leq C(M)T^{\frac{1}{4}}.\end{aligned}$$

We are done with $|h_v|_{L^2H^{s-\frac{1}{2}}}$. Next we deal with $|h_{v*}|_{L^2H^{s-\frac{1}{2}}}$. Indeed

$$\begin{aligned}|h_{v*}|_{L^2H^{s-\frac{1}{2}}} &\leq \left|(\nabla v(\zeta-\mathbb{I})A\circ X)^*A^{-1}\circ X\nabla_J X\right|_{L^2H^{s-\frac{1}{2}}}\\ &+\left|(\nabla v(A\circ X-A))^*A^{-1}\circ X\nabla_J X\right|_{L^2H^{s-\frac{1}{2}}}+\left|(\nabla vA)^*(A^{-1}\circ X-A^{-1})\nabla_J X\right|_{L^2H^{s-\frac{1}{2}}}\\ &+\left|(\nabla vA)^*A^{-1}(\nabla_J X-\mathbb{I})\right|_{L^2H^{s-\frac{1}{2}}}\equiv I_3+I_4+I_5+I_6.\end{aligned}$$

It is possible to obtain

$$\begin{aligned}I_3 &\leq C(M)\,|\nabla v|_{L^2H^{s-\frac{1}{2}}}\,|\zeta-\mathbb{I}|_{L^\infty H^{s-\frac{1}{2}}}\,|X|^2_{L^\infty H^{s-\frac{1}{2}}}\,|\nabla X|_{L^\infty H^{s-\frac{1}{2}}}\\ &\leq C\,||v||_{L^2H^{s+1}}\,||\zeta-\mathbb{I}||_{L^\infty H^s}\,||X||^3_{L^\infty H^{s+1}}\leq C(M)\,||X-\alpha||_{L^\infty H^{s+1}}\\ &\leq C(M)T^{\frac{1}{4}}.\end{aligned}$$

Similarly

$$\begin{aligned}I_4+I_5+I_6 &\leq C(M)\,|\nabla v|_{L^2H^{s-\frac{1}{2}}}\,||X-\alpha||_{L^\infty H^{s+1}}\,(||X||^2_{L^\infty H^{s+1}}+||X||_{L^\infty H^{s+1}}+1)\\ &\leq C(M)T^{\frac{1}{4}}\,||X-\alpha||_{L^\infty_{1/4}H^{s+1}}\leq C(M)T^{\frac{1}{4}}.\end{aligned}$$

It remains to control $|h_q|_{L^2H^{s-\frac{1}{2}}}$. We proceed as follows:

$$\begin{aligned}|h_q|_{L^2H^{s-\frac{1}{2}}} &\leq\left|q(A^{-1}\circ X-A^{-1})\nabla_J X\right|_{L^2H^{s-\frac{1}{2}}}+\left|qA^{-1}(\nabla_J X-\mathbb{I})\right|_{L^2H^{s-\frac{1}{2}}}\\ &\leq C(M)\,|q|_{L^2H^{s-\frac{1}{2}}}\,||X-\alpha||_{L^\infty H^{s+1}}\,(||\nabla X||_{L^\infty H^{s+1}}+1)\\ &\leq C(M)T^{1/4},\end{aligned}$$

to end with the bounds for $|h|_{L^2H^{s-\frac{1}{2}}}$.

Next step is to deal with $|h|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}$. We find

$$|h_v|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\leq|\nabla v(\zeta-\mathbb{I})\nabla_J X|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}+|\nabla v(\nabla_J X-\mathbb{I})|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\equiv K_1+K_2.$$

This splitting provides

$$K_1\leq C(|\nabla v|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}+1)\,|\zeta-\mathbb{I}|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}\,(|\nabla X|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}+1).$$

We remark that the constant above is independent of time due to Lemma 3.7. It yields

$$\begin{aligned}K_1 &\leq C(||v||_{H^{ht,s+1}}+1)\,||\zeta-\mathbb{I}||_{H^{\frac{s}{2}-\frac{1}{4}}H^{1+\delta}}\,(||X||_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}+1)\\ &\leq C(M)\,||X-\alpha||_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}\leq C(M)T^\epsilon\,||X-\alpha||_{H^{\frac{s}{2}-\frac{1}{4}+\epsilon}H^{2+\delta}}\\ &\leq C(M)T^\epsilon\,||X-\alpha||_{F^{s+1}}\leq C(M)T^\epsilon.\end{aligned}$$

Above we have used that $\nabla v_i=(\nabla v_i\cdot n_0)n_0+(\nabla v_i\cdot t_0)t_0$ for $i=1,2$ and

$$|\nabla v_i\cdot t_0|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\leq C\,|v_i|_{H^{\frac{s}{2}-\frac{1}{4}}H^1}\leq C\,||v_i||_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{3}{2}}}\leq C\,||v_i||_{H^{ht,s+1}}\,,$$

together with

$$|\nabla v_i\cdot n_0|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\leq|\nabla v_i\cdot n_0|_{H^{ht,s-\frac{1}{2}}}\leq C\,||v_i||_{H^{ht,s+1}}\,.$$

These two yield

$$|\nabla v|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\leq C\,||v||_{H^{ht,s+1}}\,. \tag{68}$$

For K_2 we find similarly

$$\begin{aligned}K_2 &\leq C(|\nabla v|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}+1)\,|\nabla X-\mathbb{I}|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}\\ &\leq C(||v||_{H^{ht,s+1}}+1)\,||X-\alpha||_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}\leq C(M)T^\epsilon.\end{aligned}$$

We are done with $|h_v|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}$. Next we deal with $|h_{v*}|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}$. Indeed

$$\begin{aligned}|h_{v*}|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} &\leq\left|(\nabla v(\zeta-\mathbb{I})A\circ X)^*A^{-1}\circ X\nabla_J X\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\\ &+\left|(\nabla v(A\circ X-A))^*A^{-1}\circ X\nabla_J X\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\\ &+\left|(\nabla vA)^*(A^{-1}\circ X-A^{-1})\nabla_J X\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\\ &+\left|(\nabla vA)^*A^{-1}(\nabla_J X-\mathbb{I})\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\equiv K_3+K_4+K_5+K_6.\end{aligned}$$

One can get

$$\begin{aligned}K_3 \leq & C(|\nabla v|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}+1)\,|\zeta-\mathbb{I}|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}\,(|A\circ X|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}+1)\\ & \times(\left|A^{-1}\circ X\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}+1)(\left|\nabla^{\perp}X\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}+1),\end{aligned}$$

where the constant is independent of time due to Lemma 3.7. It yields

$$\begin{aligned}K_3 &\leq C(M)(||v||_{H^{ht,s+1}}+1)\,||X-\alpha||_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}\,(||X||^3_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}+1)\\ &\leq C(M)T^{\epsilon}\,||X-\alpha||_{F^{s+1}} \leq C(M)T^{\epsilon}.\end{aligned}$$

Similarly

$$\begin{aligned}K_4+K_5+K_6 &\leq C(M)(||v||_{H^{ht,s+1}}+1)\,||X-\alpha||_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}\,(||X||^2_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}+1)\\ &\leq C(M)T^{\epsilon}\,||X-\alpha||_{F^{s+1}} \leq C(M)T^{\epsilon}.\end{aligned}$$

It remains to control $|h_q|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}$. We proceed as follows:

$$\begin{aligned}|h_q|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} \leq & \left|q(A^{-1}\circ X-A^{-1})\nabla_J X\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}+\left|qA^{-1}(\nabla_J X-\mathbb{I})\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}\\ \leq & C(M)(|q|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}+1)\,||X-\alpha||_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}\,(||X||_{H^{\frac{s}{2}-\frac{1}{4}}H^{2+\delta}}+1)\\ \leq & C(M)(|q|_{H^{ht,s-\frac{1}{2}}}+1)T^{\epsilon} \leq C(M)(\|q\|_{H^{ht,s}_{pr}}+1)T^{\epsilon} \leq C(M)T^{\epsilon},\end{aligned}$$

to end with the bounds for $|h|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}$. We are done with h.

Proof of part 2:

It will be enough to show that

$$\begin{aligned}\left|\left|f^{(n)}-f^{(n-1)}\right|\right|_{H^{ht,s-1}} \leq & C(M)\left(\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}+\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}}\right.\\ & \left.+||q^{(n)}-q^{(n-1)}||_{H^{ht,s}_{pr}}\right),\\ \left|\left|g^{(n)}-g^{(n-1)}\right|\right|_{\overline{H}^{ht,s}} \leq & C(M)\left(\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}+\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}}\right),\\ \left|h^{(n)}-h^{(n-1)}\right|_{H^{ht,s-\frac{1}{2}}} \leq & C(M)\left(\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}+\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}}\right.\\ & \left.+||q^{(n)}-q^{(n-1)}||_{H^{ht,s}_{pr}}\right).\end{aligned}$$

Again we split the proof in three parts:

P2.1. Estimate for $f^{(n)}-f^{(n-1)}$

We split as in (64): $f^{(j)}=f^{(j)}_w+f^{(j)}_\phi+f^{(j)}_q$. In $f^{(n)}_w-f^{(n-1)}_w$ we split further $f^{(n)}_w-f^{(n-1)}_w=d_1+...+d_6$ with the following differences

$$\begin{aligned}d_1 &= (Q^2\circ X^{(n)}-Q^2\circ X^{(n-1)})\zeta^{(n)}\partial\left(\zeta^{(n)}\partial w^{(n)}\right),\\ d_2 &= Q^2\circ X^{(n-1)}(\zeta^{(n)}-\zeta^{(n-1)})\partial\left(\zeta^{(n)}\partial w^{(n)}\right),\\ d_3 &= Q^2\circ X^{(n-1)}\zeta^{(n-1)}\partial\left((\zeta^{(n)}-\zeta^{(n-1)})\partial w^{(n)}\right),\\ d_4 &= (Q^2\circ X^{(n-1)}-Q^2)\zeta^{(n-1)}\partial\left(\zeta^{(n-1)}\partial(w^{(n)}-w^{(n-1)})\right),\\ d_5 &= Q^2(\zeta^{(n-1)}-\mathbb{I})\partial\left(\zeta^{(n-1)}\partial(w^{(n)}-w^{(n-1)})\right),\\ d_6 &= Q^2\partial\left((\zeta^{(n-1)}-\mathbb{I})\partial(w^{(n)}-w^{(n-1)})\right).\end{aligned} \tag{69}$$

Above we do not distinguish from coordinates and partial derivatives, as all the cases can be handled in the same manner. Next we estimate d_1. In order to do that we split further $d_1=d_{11}+d_{12}$ with

$$d_{11}=(Q^2\circ X^{(n)}-Q^2\circ X^{(n-1)})\zeta^{(n)}\partial\zeta^{(n)}\partial w^{(n)},$$

$$d_{12} = (Q^2 \circ X^{(n)} - Q^2 \circ X^{(n-1)})\zeta^{(n)}\zeta^{(n)}\partial^2 w^{(n)}.$$

We take

$$\begin{aligned}
||d_{11}||_{L^2H^{s-1}} &\leq C \left|\left|Q^2 \circ X^{(n)} - Q^2 \circ X^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\zeta^{(n)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial\zeta^{(n)}\partial w^{(n)}\right|\right|_{L^2H^{s-1}} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial\zeta^{(n)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial w^{(n)}\right|\right|_{L^2H^{s-1}} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \leq C(M)T^{1/4} \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{F^{s+1}},
\end{aligned}$$

and

$$\begin{aligned}
||d_{11}||_{H^{\frac{s-1}{2}}L^2} &\leq C \left|\left|Q^2 \circ X^{(n)} - Q^2 \circ X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \\
&\quad \times (\left|\left|\zeta^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} + 1) \left|\left|\partial\zeta^{(n)}\partial w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \left|\left|\partial\zeta^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{\epsilon}} \left|\left|\partial w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1-\epsilon}}.
\end{aligned}$$

We deal with above terms as before to get

$$||d_{11}||_{H^{\frac{s-1}{2}}L^2} \leq C(M)T^\epsilon \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{F^{s+1}}.$$

For d_{12} we find

$$\begin{aligned}
||d_{12}||_{L^2H^{s-1}} &\leq C \left|\left|Q^2 \circ X^{(n)} - Q^2 \circ X^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\zeta^{(n)}\right|\right|^2_{L^\infty H^{s-1}} \left|\left|\partial^2 w^{(n)}\right|\right|_{L^2H^{s-1}} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \leq C(M)T^{1/4} \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{F^{s+1}}
\end{aligned}$$

and

$$\begin{aligned}
||d_{12}||_{H^{\frac{s-1}{2}}L^2} &\leq C \left|\left|Q^2 \circ X^{(n)} - Q^2 \circ X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \\
&\quad \times (\left|\left|\zeta^{(n)} - \mathbb{I}\right|\right|^2_{H^{\frac{s-1}{2}}H^{1+\epsilon}} + 1) \left|\left|\partial^2 w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \leq C(M)T^\epsilon \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{F^{s+1}}.
\end{aligned}$$

In order to continue we decompose the next term, $d_2 = d_{21} + d_{22}$ where

$$\begin{aligned}
d_{21} &= Q^2 \circ X^{(n-1)}(\zeta^{(n)} - \zeta^{(n-1)})\partial\zeta^{(n)}\partial w^{(n)}, \\
d_{22} &= Q^2 \circ X^{(n-1)}(\zeta^{(n)} - \zeta^{(n-1)})\zeta^{(n)}\partial^2 w^{(n)}.
\end{aligned}$$

We take

$$\begin{aligned}
||d_{21}||_{L^2H^{s-1}} &\leq C \left|\left|Q^2 \circ X^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\zeta^{(n)} - \zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial\zeta^{(n)}\partial w^{(n)}\right|\right|_{L^2H^{s-1}} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^{s}} \left|\left|\partial\zeta^{(n)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial w^{(n)}\right|\right|_{L^2H^{s-1}} \\
&\leq C(M)T^{1/4} \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{F^{s+1}}.
\end{aligned}$$

$$\begin{aligned}
||d_{21}||_{H^{\frac{s-1}{2}}L^2} &\leq C(\left|\left|Q^2 \circ X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} + 1) \\
&\quad \times \left|\left|\zeta^{(n)} - \zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \left|\left|\partial\zeta^{(n)}\partial w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{2+\epsilon}} \left|\left|\partial\zeta^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{\epsilon}} \left|\left|\partial w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1-\epsilon}} \\
&\leq C(M)T^\epsilon \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{F^{s+1}}.
\end{aligned}$$

For d_{22} we find

$$\begin{aligned}
||d_{22}||_{L^2H^{s-1}} &\leq C(M) \left|\left|\zeta^{(n)} - \zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial^2 w^{(n)}\right|\right|_{L^2H^{s-1}} \\
&\leq C(M) \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^{s}} \leq C(M)T^{1/4} \left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{F^{s+1}},
\end{aligned}$$

and

$$\begin{aligned}
||d_{22}||_{H^{\frac{s-1}{2}}L^2} &\leq C(M)\left|\left|\zeta^{(n)}-\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}\left|\left|\partial^2 w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2}\\
&\leq C(M)\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{2+\epsilon}}\leq C(M)T^{\epsilon}\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}.
\end{aligned}$$

In d_3 we split as follows $d_3=d_{31}+d_{32}$ with

$$d_{31}=Q^2\circ X^{(n-1)}\zeta^{(n-1)}\partial(\zeta^{(n)}-\zeta^{(n-1)})\partial w^{(n)},$$

and

$$d_{32}=Q^2\circ X^{(n-1)}\zeta^{(n-1)}(\zeta^{(n)}-\zeta^{(n-1)})\partial^2 w^{(n)}.$$

For d_{31} it is possible to get

$$\begin{aligned}
||d_{31}||_{L^2H^{s-1}} &\leq C\left|\left|Q^2\circ X^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\partial(\zeta^{(n)}-\zeta^{(n-1)})\partial w^{(n)}\right|\right|_{L^2H^{s-1}}\\
&\leq C(M)\left|\left|\partial(\zeta^{(n)}-\zeta^{(n-1)})\right|\right|_{L^\infty H^{s-1}}\left|\left|\partial w^{(n)}\right|\right|_{L^2H^{s-1}}\\
&\leq C(M)T^{1/4}\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}},
\end{aligned}$$

and

$$\begin{aligned}
||d_{31}||_{H^{\frac{s-1}{2}}L^2} &\leq C(\left|\left|Q^2\circ X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}+1)\\
&\qquad\times\left(\left|\left|\zeta^{(n-1)}-\mathbb{I}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}+1\right)\left|\left|\partial(\zeta^{(n)}-\zeta^{(n-1)})\partial w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2}\\
&\leq C(M)\left|\left|\partial(\zeta^{(n)}-\zeta^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}H^{\delta}}\left|\left|\partial w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1-\delta}}\\
&\leq C(M)\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{2+\delta}}\leq C(M)T^{\epsilon}\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}.
\end{aligned}$$

Next, for d_{32} we obtain

$$\begin{aligned}
||d_{32}||_{L^2H^{s-1}} &\leq C\left|\left|Q^2\circ X^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|(\zeta^{(n)}-\zeta^{(n-1)})\partial^2 w^{(n)}\right|\right|_{L^2H^{s-1}}\\
&\leq C(M)\left|\left|\zeta^{(n)}-\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\partial^2 w^{(n)}\right|\right|_{L^2H^{s-1}}\\
&\leq C(M)\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{L^\infty H^{s}}\leq C(M)T^{1/4}\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}.
\end{aligned}$$

The usual approach also gives

$$\begin{aligned}
||d_{32}||_{H^{\frac{s-1}{2}}L^2} &\leq C\Big(\left|\left|Q^2\circ X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}+1\Big)\\
&\qquad\times\Big(\left|\left|\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}+1\Big)\left|\left|(\zeta^{(n)}-\zeta^{(n-1)})\partial^2 w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2}\\
&\leq C(M)\left|\left|\zeta^{(n)}-\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\left|\left|\partial^2 w^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2}\\
&\leq C(M)\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{2+\delta}}\leq C(M)T^{\epsilon}\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}.
\end{aligned}$$

In d_4 we proceed by considering $d_4=d_{41}+d_{42}$ with

$$d_{41}=(Q^2\circ X^{(n-1)}-Q^2)\zeta^{(n-1)}\partial\zeta^{(n-1)}\partial(w^{(n)}-w^{(n-1)}),$$

and

$$d_{42}=(Q^2\circ X^{(n-1)}-Q^2)\zeta^{(n-1)}\zeta^{(n-1)}\partial^2(w^{(n)}-w^{(n-1)}).$$

For d_{41} it is possible to get

$$\begin{aligned}
||d_{41}||_{L^2H^{s-1}} &\leq C\left|\left|Q^2\circ X^{(n-1)}-Q^2\right|\right|_{L^\infty H^{s-1}}\\
&\qquad\times\left|\left|\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\partial\zeta^{(n-1)}\partial(w^{(n)}-w^{(n-1)})\right|\right|_{L^2H^{s-1}}\\
&\leq C(M)\left|\left|X^{(n-1)}-\alpha\right|\right|_{L^\infty H^{s-1}}\left|\left|\partial\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\partial(w^{(n)}-w^{(n-1)})\right|\right|_{L^2H^{s-1}}\\
&\leq C(M)T^{1/4}\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}},
\end{aligned}$$

and

$$\begin{aligned}
||d_{41}||_{H^{\frac{s-1}{2}}L^2} \leq C &\left|\left|Q^2 \circ X^{(n-1)} - Q^2\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \\
&\times \left(\left|\left|\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} + 1\right) \left|\left|\partial\zeta^{(n-1)}\partial(w^{(n)} - w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}L^2} \\
\leq C(M) &\left|\left|X^{(n-1)} - \alpha\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \left|\left|\partial\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{\delta}} \left|\left|\partial(w^{(n)} - w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}H^{1-\delta}} \\
\leq C(M)&T^{\epsilon} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}.
\end{aligned}$$

For d_{42} it is possible to get

$$\begin{aligned}
||d_{42}||_{L^2H^{s-1}} &\leq C \left|\left|Q^2 \circ X^{(n-1)} - Q^2\right|\right|_{L^\infty H^{s-1}} \left|\left|\zeta^{(n-1)}\right|\right|^2_{L^\infty H^{s-1}} \left|\left|\partial^2(w^{(n)} - w^{(n-1)})\right|\right|_{L^2H^{s-1}} \\
&\leq C(M) \left|\left|X^{(n)} - \alpha\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial^2(w^{(n)} - w^{(n-1)})\right|\right|_{L^2H^{s-1}} \\
&\leq C(M)T^{1/4} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}},
\end{aligned}$$

and

$$\begin{aligned}
||d_{42}||_{H^{\frac{s-1}{2}}L^2} \leq C &\left|\left|Q^2 \circ X^{(n-1)} - Q^2\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \\
&\times \left(\left|\left|\zeta^{(n-1)}\right|\right|^2_{H^{\frac{s-1}{2}}H^{1+\epsilon}} + 1\right) \left|\left|\partial^2(w^{(n)} - w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}L^2} \\
\leq C(M) &\left|\left|X^{(n)} - \alpha\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \left|\left|\partial^2(w^{(n)} - w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}L^2} \\
\leq C(M)&T^{\epsilon} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}.
\end{aligned}$$

Analogously, one could take $d_5 = d_{51} + d_{52}$ with

$$d_{51} = Q^2(\zeta^{(n-1)} - \mathbb{I})\partial\zeta^{(n-1)}\partial(w^{(n)} - w^{(n-1)}),$$

and

$$d_{52} = Q^2(\zeta^{(n-1)} - \mathbb{I})\zeta^{(n-1)}\partial^2(w^{(n)} - w^{(n-1)}).$$

The splitting yields

$$\begin{aligned}
||d_{51}||_{L^2H^{s-1}} &\leq C \left|\left|\zeta^{(n-1)} - \mathbb{I}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial(w^{(n)} - w^{(n-1)})\right|\right|_{L^2H^{s-1}} \\
&\leq C(M)T^{1/4} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}},
\end{aligned}$$

and

$$\begin{aligned}
||d_{51}||_{H^{\frac{s-1}{2}}L^2} &\leq C \left|\left|\zeta^{(n-1)} - \mathbb{I}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \left|\left|\partial\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{\delta}} \left|\left|\partial(w^{(n)} - w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}H^{1-\delta}} \\
&\leq C(M)T^{\epsilon} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}.
\end{aligned}$$

For d_{52} we proceed as follows

$$\begin{aligned}
||d_{52}||_{L^2H^{s-1}} &\leq C \left|\left|\zeta^{(n-1)} - \mathbb{I}\right|\right|_{L^\infty H^{s-1}} \left|\left|\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}} \left|\left|\partial^2(w^{(n)} - w^{(n-1)})\right|\right|_{L^2H^{s-1}} \\
&\leq C(M)T^{1/4} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}},
\end{aligned}$$

$$\begin{aligned}
||d_{52}||_{H^{\frac{s-1}{2}}L^2} \leq C &\left|\left|\zeta^{(n-1)} - \mathbb{I}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} \\
&\times \left(\left|\left|\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}} + 1\right) \left|\left|\partial^2(w^{(n)} - w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}L^2} \\
\leq C(M)&T^{\epsilon} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}.
\end{aligned}$$

Finally, writing $d_6 = d_{61} + d_{62}$ where

$$d_{61} = Q^2\partial\zeta^{(n-1)}\partial(w^{(n)} - w^{(n-1)}), \text{ and } d_{62} = Q^2(\zeta^{(n-1)} - \mathbb{I})\partial^2(w^{(n)} - w^{(n-1)}),$$

it is possible to find

$$\begin{aligned}||d_{61}||_{L^2H^{s-1}} &\leq C\left|\left|\partial\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\partial(w^{(n)}-w^{(n-1)})\right|\right|_{L^2H^{s-1}}\\ &\leq C(M)T^{1/4}\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}},\end{aligned}$$

and

$$\begin{aligned}||d_{61}||_{H^{\frac{s-1}{2}}L^2} &\leq C\left|\left|\partial\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^\delta}\left|\left|\partial(w^{(n)}-w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}H^{1-\delta}}\\ &\leq C(M)T^\epsilon\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}}.\end{aligned}$$

For d_{62} we proceed as for d_{52} to get

$$\begin{aligned}||d_{62}||_{L^2H^{s-1}} &\leq C(M)T^{1/4}\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}},\\ ||d_{62}||_{H^{\frac{s-1}{2}}L^2} &\leq C(M)T^\epsilon\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}}.\end{aligned}$$

We are done with $f_w^{(n)}-f_w^{(n-1)}$. To continue with $f_\phi^{(n)}-f_\phi^{(n-1)}$ we decompose as before but this time using $f_\phi^{(n)}-f_\phi^{(n-1)}=d_7+d_8+d_9$ where

$$\begin{aligned}d_7 &= (Q^2\circ X^{(n)}-Q^2\circ X^{(n-1)})\zeta^{(n)}\partial\left(\zeta^{(n)}\partial\phi\right),\\ d_8 &= Q^2\circ X^{(n-1)}(\zeta^{(n)}-\zeta^{(n-1)})\partial\left(\zeta^{(n)}\partial\phi\right),\\ d_9 &= Q^2\circ X^{(n-1)}\zeta^{(n-1)}\partial\left((\zeta^{(n)}-\zeta^{(n-1)})\partial\phi\right).\end{aligned}$$

If we proceed as for d_1, d_2 and d_3 it is possible to find

$$||d_7+d_8+d_9||_{H^{ht,s-1}}\leq C(M)T^\epsilon\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}.$$

We move to the f term involving q. The splitting $f_q^{(n)}-f_q^{(n-1)}=d_1^q+d_2^q+d_3^q+d_4^q$ with

$$\begin{aligned}d_1^q &= -(A\circ X^{(n)}(\zeta^{(n)}-\zeta^{(n-1)}))^*\nabla q^{(n)},\\ d_2^q &= -((A\circ X^{(n)}-A\circ X^{(n-1)})\zeta^{(n-1)})^*\nabla q^{(n)},\\ d_3^q &= -(A\circ X^{(n-1)}(\zeta^{(n-1)}-\mathbb{I}))^*\nabla(q^{(n)}-q^{(n-1)}),\\ d_4^q &= -(A\circ X^{(n-1)}-A)^*\nabla(q^{(n)}-q^{(n-1)}),\end{aligned}\tag{70}$$

allows us to do the work. In fact

$$\begin{aligned}||d_1^q||_{L^2H^{s-1}} &\leq C\left|\left|\zeta^{(n)}-\zeta^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|A\circ X^{(n)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\nabla q^{(n)}\right|\right|_{L^2H^{s-1}}\\ &\leq C(M)T^{1/4}\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}},\end{aligned}$$

$$\begin{aligned}||d_1^q||_{H^{\frac{s-1}{2}}L^2} &\leq C\left|\left|\zeta^{(n)}-\zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}\\ &\quad\times\left(\left|\left|A\circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}+1\right)\left(\left|\left|\nabla q^{(n)}\right|\right|_{H^{\frac{s-1}{2}}L^2}+1\right)\\ &\leq C(M)T^\epsilon\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}.\end{aligned}$$

Sharing norms in the same manner gives

$$||d_2^q||_{L^2H^{s-1}}\leq C(M)T^{1/4}\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}},$$

and

$$||d_2^q||_{H^{\frac{s-1}{2}}L^2}\leq C(M)T^\epsilon\left|\left|X^{(n)}-X^{(n-1)}\right|\right|_{F^{s+1}}.$$

In a similar way

$$\begin{aligned}||d_3^q||_{L^2H^{s-1}} &\leq C\left|\left|\zeta^{(n-1)}-\mathbb{I}\right|\right|_{L^\infty H^{s-1}}\left|\left|A\circ X^{(n-1)}\right|\right|_{L^\infty H^{s-1}}\left|\left|\nabla(q^{(n)}-q^{(n-1)})\right|\right|_{L^2H^{s-1}}\\ &\leq C(M)T^{1/4}\left|\left|\nabla(q_w^{(n)}-q_w^{(n-1)})\right|\right|_{L^2H^{s-1}}\leq C(M)T^{1/4}\|q_w^{(n)}-q_w^{(n-1)}\|_{H^{ht,s}_{pr}},\end{aligned}$$

and

$$\begin{aligned}||d_3^q||_{H^{\frac{s-1}{2}}L^2} &\leq C\left|\left|\zeta^{(n-1)}-\mathbb{I}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}\\ &\quad\times\left(\left|\left|A\circ X^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}+1\right)\left|\left|\nabla(q^{(n)}-q^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}L^2}\\ &\leq C(M)T^\epsilon\left|\left|\nabla(q_w^{(n)}-q_w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}L^2}\\ &\leq C(M)T^\epsilon\|q_w^{(n)}-q_w^{(n-1)}\|_{H^{ht,s}_{pr}}.\end{aligned}$$

Finally

$$\begin{aligned}||d_4^q||_{L^2H^{s-1}} &\leq C\left|\left|A\circ X^{(n-1)}-A\right|\right|_{L^\infty H^{s-1}}\left|\left|\nabla(q^{(n)}-q^{(n-1)})\right|\right|_{L^2H^{s-1}}\\ &\leq C(M)T^{1/4}\|q_w^{(n)}-q_w^{(n-1)}\|_{H^{ht,s}_{pr}},\end{aligned}$$

and

$$\begin{aligned}||d_4^q||_{H^{\frac{s-1}{2}}L^2} &\leq C\left|\left|A\circ X^{(n-1)}-A\right|\right|_{H^{\frac{s-1}{2}}H^{1+\epsilon}}\left|\left|\nabla(q^{(n)}-q^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}L^2}\\ &\leq C(M)T^\epsilon\|q_w^{(n)}-q_w^{(n-1)}\|_{H^{ht,s}_{pr}}.\end{aligned}$$

P2.2. Estimate for $g^{(n)}-g^{(n-1)}$:

We are concerned with the estimates of the norms

$$\left|\left|g^{(n)}-g^{(n-1)}\right|\right|_{\overline{H}^{ht,s}}\leq\left|\left|g^{(n)}-g^{(n-1)}\right|\right|_{H^0H^s}+\left|\left|g^{(n)}-g^{(n-1)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}},$$

where we can write $g^{(n-1)}-g^{(n)}=d_1+d_2+d_3+d_4$ where

$$\begin{aligned}d_1&=-Tr((\nabla v^{(n)}-\nabla v^{(n-1)})(\zeta^{(n)}-\mathbb{I})A\circ X^{(n)}),\\ d_2&=-Tr(\nabla v^{(n-1)}(\zeta^{(n)}-\zeta^{(n-1)})A\circ X^{(n)})\\ d_3&=-Tr(\nabla v^{(n-1)}\zeta^{(n-1)}(A\circ X^{(n)}-A\circ X^{(n-1)}))\\ d_4&=-Tr((\nabla v^{(n)}-\nabla v^{(n-1)})(A\circ X^{(n)}-A)).\end{aligned}\tag{71}$$

First we will estimate $\left|\left|g^{(n)}-g^{(n-1)}\right|\right|_{H^0H^s}$. With d_1 we proceed as follows

$$\begin{aligned}||d_1||_{H^0H^s} &=\left|\left|(\nabla v^{(n)}-\nabla v^{(n-1)})(\zeta^{(n)}-\mathbb{I})A\circ X^{(n)}\right|\right|_{H^0H^s}\\ &\leq\left|\left|\zeta^{(n)}-\mathbb{I}\right|\right|_{L^\infty H^s}\left|\left|A\circ X^{(n)}\right|\right|_{H^0H^s}\left|\left|\nabla w^{(n)}-\nabla w^{(n-1)}\right|\right|_{H^0H^s}.\end{aligned}$$

Then lemma 3.9 implies

$$||d_1||_{H^0H^s}\leq C(M)\left|\left|\zeta^{(n)}-\mathbb{I}\right|\right|_{L^\infty H^s}\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}}.$$

And therefore, by using (65) we obtain that

$$||d_1||_{H^0H^s}\leq C(M)T^{\frac{1}{4}}\left|\left|w^{(n)}-w^{(n-1)}\right|\right|_{H^{ht,s+1}}.$$

For d_2 we have that

$$||d_2||_{H^0H^s}\leq\left|\left|\nabla v^{(n-1)}\right|\right|_{H^0H^s}\left|\left|A\circ X^{(n)}\right|\right|_{L^\infty H^s}\left|\left|\zeta^{(n)}-\zeta^{(n-1)}\right|\right|_{L^\infty H^s}.$$

Thus, using that

$$\zeta^{(n)} - \zeta^{(n-1)} = (\nabla X^{(n)})^{-1}\left(\nabla X^{(n-1)} - \nabla X^{(n)}\right)(\nabla X^{(n-1)})^{-1}$$

yields,

$$||d_2||_{H^0H^s} \le C(M)T^{\frac{1}{4}}\left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty_{1/4}H^{s+1}}.$$

The estimate for d_3 follows the next steps

$$\begin{aligned}
||d_3||_{H^0H^s} &\le \left|\left|\nabla v^{(n-1)}\right|\right|_{H^0H^s}\left|\left|\zeta^{(n-1)}\right|\right|_{L^\infty H^s}\left|\left|A\circ X^{(n)} - A\circ X^{(n-1)}\right|\right|_{L^\infty H^s}\\
&\le C(M)\left|\left|RR(X^{(n)},X^{(n-1)})(X^{(n)} - X^{(n-1)})\right|\right|_{L^\infty H^s}\\
&\le C(M)\left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty H^s} \le C(M)T^{\frac{1}{4}}\left|\left|X^{(n)} - X^{(n-1)}\right|\right|_{L^\infty_{1/4}H^s}.
\end{aligned}$$

And for d_4 we have that

$$\begin{aligned}
||d_4||_{H^0H^s} &\le \left|\left|v^{(n)} - v^{(n-1)}\right|\right|_{H^{ht,s+1}}\left|\left|A\circ X^{(n)} - A\right|\right|_{L^\infty H^s}\\
&\le C(M)\left|\left|v^{(n)} - v^{(n-1)}\right|\right|_{H^{ht,s+1}}\left|\left|X^{(n)} - \alpha\right|\right|_{L^\infty H^s}\\
&\le C(M)T^{\frac{1}{4}}\left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}.
\end{aligned}$$

It remains to estimate the $||\cdot||_{H^{\frac{s+1}{2}}H^{-1}}$ norm. For d_1 we have that

$$d_1 = -\int_0^t Tr(\partial_t((\nabla v^{(n)} - \nabla v^{(n-1)})(\zeta^{(n)} - \mathbb{I})A\circ X^{(n)}))d\tau = d_{11} + d_{12} + d_{13}$$

where

$$\begin{aligned}
d_{11} &= -\int_0^t Tr(\partial_t(\nabla v^{(n)} - \nabla v^{(n-1)})(\zeta^{(n)} - \mathbb{I})A\circ X^{(n)})d\tau,\\
d_{12} &= -\int_0^t Tr((\nabla v^{(n)} - \nabla v^{(n-1)})\partial_t(\zeta^{(n)} - \mathbb{I})A\circ X^{(n)})d\tau,\\
d_{13} &= -\int_0^t Tr((\nabla v^{(n)} - \nabla v^{(n-1)})(\zeta^{(n)} - \mathbb{I})\partial_t(A\circ X^{(n)}))d\tau.
\end{aligned} \tag{72}$$

By applying lemma 3.4 with $\varepsilon = 0$ we obtain

$$||d_{11}||_{H^{\frac{s+1}{2}}H^{-1}} \le \left|\left|\partial_t(\nabla v^{(n)} - \nabla v^{(n-1)})(\zeta^{(n)} - \mathbb{I})A\circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}.$$

Now we use part 3 of lemma 3.5 to yield

$$||d_{11}||_{H^{\frac{s+1}{2}}H^{-1}} \le \left|\left|\partial_t(\nabla v^{(n)} - \nabla v^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}\left|\left|\zeta^{(n)} - \mathbb{I}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\left(1 + \left|\left|A\circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\right)$$

In addition lemmas 3.9, 3.8 and 3.3 imply

$$||d_{11}||_{H^{\frac{s+1}{2}}H^{-1}} \le C(M)\left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}\left|\left|\zeta^{(n)} - \mathbb{I}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}.$$

And we can apply the first part of lemma 3.4 to get

$$\begin{aligned}
\left|\left|\zeta^{(n)} - \mathbb{I}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}} &\le \left|\left|\partial_t\int_0^t(\zeta^{(n)} - \mathbb{I})\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}\left|\left|\int_0^t(\zeta^{(n)} - \mathbb{I})\right|\right|_{H^{\frac{s-1}{2}+\varepsilon+1-\varepsilon}H^{1+\delta}}\\
&\le CT^\varepsilon\left|\left|\zeta^{(n)} - \mathbb{I}\right|\right|_{H^{\frac{s-1}{2}+\varepsilon}H^{1+\delta}},
\end{aligned}$$

for $0 < \frac{s-1}{2} + \varepsilon < 1$. Thus, by lemma 3.8,

$$||d_{11}||_{H^{\frac{s+1}{2}}H^{-1}} \le C(M)T^\varepsilon\left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}.$$

Next we bound d_{12}. We use a similar argument but using part 4 of lemma 3.5 instead of part 3 and using a small modification of part 3 of lemma 3.7. Indeed

$$\begin{aligned}||d_{12}||_{H^{\frac{s+1}{2}}H^{-1}} &= \left|\left|(\nabla w^{(n)} - \nabla w^{(n-1)})\partial_t \zeta^{(n)} A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}} \\ &\leq C \left|\left|(\nabla w^{(n)} - \nabla w^{(n-1)})\partial_t \zeta^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}} (1 + \left|\left|A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}).\end{aligned}$$

In addition we can make the following splitting

$$\begin{aligned}||d_{12}||_{H^{\frac{s+1}{2}}H^{-1}} \leq& C \left|\left|(\nabla w^{(n)} - \nabla w^{(n-1)})(\partial_t \zeta^{(n)} - \partial_t \zeta^{(n)}|_{t=0})\right|\right|_{H^{\frac{s-1}{2}}H^{-1}} \\ &+ C \left|\left|(\nabla w^{(n)} - \nabla w^{(n-1)})\partial_t \zeta^{(n)}|_{t=0}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}} \\ \leq C \left|\left|(\nabla w^{(n)} - \nabla w^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}H^{1}} &\left|\left|\partial_t \zeta^{(n)} - \partial_t \zeta^{(n)}|_{t=0}\right|\right|_{H^{\frac{s-1}{2}}H^{0}} + C \left|\left|\nabla w^{(n)} - \nabla w^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}.\end{aligned}$$

Since

$$\left|\left|\partial_t \zeta^{(n)} - \partial_t \zeta^{(n)}|_{t=0}\right|\right|_{H^{\frac{s-1}{2}}H^{0}} \leq C(M)T^{\varepsilon},$$

and

$$\left|\left|\nabla w^{(n)} - \nabla w^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}} \leq C(M)T^{\varepsilon} \left|\left|w^{(n)} - w^{(n-1)}\right|\right|_{H^{ht,s+1}}$$

we obtain a suitable estimate for d_{12}.

The term d_{13} does not cause any difficulty and it can be handled as before.

The estimate for d_2 follows similar steps,

$$\begin{aligned}||d_2||_{H^{\frac{s+1}{2}}H^{-1}} &\leq \left|\left|\nabla v^{(n-1)}(\zeta^{(n)} - \zeta^{(n-1)})A \circ X^{(n)}\right|\right|_{H^{\frac{s+1}{2}}H^{-1}} \\ &\leq \left|\left|\int_0^t \partial_t(\nabla v^{(n-1)}(\zeta^{(n)} - \zeta^{(n-1)})A \circ X^{(n)})\right|\right|_{H^{\frac{s+1}{2}}H^{-1}},\end{aligned} \tag{73}$$

and therefore

$$\begin{aligned}||d_2||_{H^{\frac{s+1}{2}}H^{-1}} \leq& \left|\left|\partial_t \nabla v^{(n-1)}(\zeta^{(n)} - \zeta^{(n-1)})A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}} \\ &+ \left|\left|\nabla v^{(n-1)}\partial_t(\zeta^{(n)} - \zeta^{(n-1)})A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}} \\ &+ \left|\left|\nabla v^{(n-1)}(\zeta^{(n)} - \zeta^{(n-1)})\partial_t A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}.\end{aligned}$$

Each addend is estimated in a different manner as follows

$$\begin{aligned}||d_2||_{H^{\frac{s+1}{2}}H^{-1}} \leq& C(M)(1 + \left|\left|\nabla \partial_t v^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{-1}}) \left|\left|\zeta^{(n)} - \zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}} (1 + \left|\left|A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}) \\ &+ C(M)(1 + \left|\left|\nabla v^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1}}) \left|\left|\partial_t(\zeta^{(n)} - \zeta^{(n-1)})\right|\right|_{H^{\frac{s-1}{2}}H^{0}} (1 + \left|\left|A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}) \\ &+ C(M)(1 + \left|\left|\nabla v^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1}}) \left|\left|\zeta^{(n)} - \zeta^{(n-1)}\right|\right|_{H^{\frac{s-1}{2}}H^{0}} (1 + \left|\left|\partial_t A \circ X^{(n)}\right|\right|_{H^{\frac{s-1}{2}}H^{1+\delta}}),\end{aligned}$$

and we have already bounded every term in here.

We will give full detail for d_3. Again we can split d_3 in the following way

$$d_3 = -\int_0^t Tr \partial_t (\nabla v^{(n-1)} \zeta^{(n-1)} (A \circ X^{(n)} - A \circ X^{(n-1)}))d\tau = d_{31} + d_{32} + d_{33}$$

where

$$\begin{aligned}d_{31} &= -\int_0^t Tr(\partial_t \nabla v^{(n-1)} \zeta^{(n-1)} (A \circ X^{(n)} - A \circ X^{(n-1)}))d\tau \\ d_{32} &= -\int_0^t Tr(\nabla v^{(n-1)} \partial_t \zeta^{(n-1)} (A \circ X^{(n)} - A \circ X^{(n-1)}))d\tau \\ d_{33} &= -\int_0^t Tr(\nabla v^{(n-1)} \zeta^{(n-1)} \partial_t (A \circ X^{(n)} - A \circ X^{(n-1)}))d\tau.\end{aligned} \tag{74}$$

The estimates of these three terms follow similar steps as those for d_1 and d_2. First we apply part 2 of lemma 3.4 with $\varepsilon=0$ since $0<\frac{s-1}{2}<1$ to find

$$\begin{aligned}\|d_{31}\|_{H^{\frac{s+1}{2}}H^{-1}} \le&\left\|\partial_t\nabla v^{(n-1)}\zeta^{(n-1)}(A\circ X^{(n)}-A\circ X^{(n-1)})\right\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\le C(M)(1+\left\|\nabla\partial_t v^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{-1}})(1+\left\|\zeta^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}})\left\|A\circ X^{(n)}-A\circ X^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}}\\ &\le C(M)\left\|A\circ X^{(n)}-A\circ X^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}}\le C(M)\left\|X^{(n)}-X^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}}.\end{aligned}$$

Above we use part 3 of lemma 3.5, part 3 of lemma 3.7 and lemma 3.9 as we did in the proof of proposition 5.3. In addition

$$\begin{aligned}\|d_{31}\|_{H^{\frac{s+1}{2}}H^{-1}} \le& C(M)\left\|\partial_t\int_0^t X^{(n)}-X^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}}\le\left\|\int_0^t X^{(n)}-X^{(n-1)}\right\|_{H^{\frac{s-1}{2}+\varepsilon+1-\varepsilon}H^{1+\delta}}\\ \le& CT^{\varepsilon}\left\|X^{(n)}-X^{(n-1)}\right\|_{H^{\frac{s-1}{2}+\varepsilon}H^{1+\delta}}\le CT^{\varepsilon}\left\|X^{(n)}-X^{(n-1)}\right\|_{F^{s+1}},\end{aligned}$$

for $0<\frac{s-1}{2}+\varepsilon<1$. The estimate for d_{32} is quite similar. Indeed, first we use part 2 of lemma 3.4 with $\varepsilon=0$ to obtain

$$\begin{aligned}\|d_{32}\|_{H^{\frac{s+1}{2}}H^{-1}} \le& C\left\|\nabla v^{(n-1)}\partial_t\zeta^{(n-1)}(A\circ X^{(n)}-A\circ X^{(n-1)})\right\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\le C(M)\left(1+\left\|\nabla w^{(n)}\right\|_{H^{\frac{s-1}{2}}H^{1}}\right)(1+\left\|\partial_t\zeta^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{0}})\left\|A\circ X^{(n)}-A\circ X^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}}.\end{aligned}$$

Above we apply part 4 of lemma 3.5 and the rest of the estimate follows the same steps. Next we notice that $\partial_t(A\circ X^{(n)}-A\circ X^{(n-1)})|_{t=0}=0$ and in d_{33} we can make use of part 2 of lemma 3.4 with $\varepsilon=0$ since $0<\frac{s-1}{2}<1$. We have that

$$\begin{aligned}\|d_{33}\|_{H^{\frac{s+1}{2}}H^{-1}} \le C(M)(1+&\left\|\nabla w^{(n)}\right\|_{H^{\frac{s-1}{2}}H^{1}})(1+\left\|\zeta^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}})\\ &\times\left\|\partial_t(A\circ X^{(n)}-A\circ X^{(n-1)})\right\|_{H^{\frac{s-1}{2}}H^{0}},\end{aligned}$$

and therefore, it is enough to estimate $\left\|\partial_t(X^{(n)}-X^{(n-1)})\right\|_{H^{\frac{s-1}{2}}H^0}$. As we did before we obtain

$$\begin{aligned}\left\|\partial_t(X^{(n)}-X^{(n-1)})\right\|_{H^{\frac{s-1}{2}}H^{0}} \le& CT^{\varepsilon}\left\|\partial_t(X^{(n)}-X^{(n-1)})\right\|_{H^{\frac{s-1}{2}+\varepsilon}H^{0}}\\ \le& CT^{\varepsilon}\left\|X^{(n)}-X^{(n-1)}\right\|_{F^{s+1}}.\end{aligned}$$

Finally we estimate d_4 in (71). It is possible to get

$$\begin{aligned}\|d_4\|_{H^{\frac{s+1}{2}}H^{-1}} \le& \left\|(\nabla v^{(n)}-\nabla v^{(n-1)})(A\circ X^{(n)}-A)\right\|_{H^{\frac{s+1}{2}}H^{-1}}\\ \le& \left\|\partial_t(\nabla v^{(n)}-\nabla v^{(n-1)})(A\circ X^{(n)}-A)\right\|_{H^{\frac{s-1}{2}}H^{-1}}+\left\|(\nabla w^{(n)}-\nabla w^{(n-1)})\partial_t A\circ X^{(n)}\right\|_{H^{\frac{s-1}{2}}H^{-1}}.\end{aligned}\tag{75}$$

By applying part 4 of lemma 3.5 and part 3 of lemma 3.7 we have that

$$\begin{aligned}\|d_4\|_{H^{\frac{s+1}{2}}H^{-1}} \le& \left\|\partial_t\nabla w^{(n)}-\partial_t\nabla w^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{-1}}\left\|A\circ X^{(n)}-A\right\|_{H^{\frac{s-1}{2}}H^{1+\delta}}\\ &+C(M)\left\|\nabla w^{(n)}-\nabla w^{(n-1)}\right\|_{H^{\frac{s-1}{2}}H^{-1}},\end{aligned}$$

and we have already seen that then

$$\|d_4\|_{H^{\frac{s+1}{2}}H^{-1}}\le C(M)T^{\varepsilon}\left\|w^{(n)}-w^{(n-1)}\right\|_{H^{ht,s+1}}.$$

P2.3. Estimate for $h^{(n)}-h^{(n-1)}$:

We split using (67) and decompose further $h_v^{(n)}-h_v^{(n-1)}=d_1+d_2+d_3+d_4$ where the differences are given by

$$d_1=\nabla v^{(n)}(\zeta^{(n)}-\zeta^{(n-1)})\nabla_J X^{(n)}n_0,\quad d_2=\nabla v^{(n)}\zeta^{(n-1)}(\nabla_J X^{(n)}-\nabla_J X^{(n-1)})n_0,\tag{76}$$

$$d_3 = \nabla(v^{(n)} - v^{(n-1)})(\zeta^{(n-1)} - \mathbb{I})\nabla_J X^{(n-1)} n_0, \quad d_4 = \nabla(v^{(n)} - v^{(n-1)})(\nabla_J X^{(n-1)} - \mathbb{I}) n_0.$$

We estimate as follows. For d_1 we find:

$$\begin{aligned}
\left|d_1\right|_{L^2 H^{s-\frac{1}{2}}} &\leq C \left|\nabla v^{(n)}\right|_{L^2 H^{s-\frac{1}{2}}} \left|\zeta^{(n)} - \zeta^{(n-1)}\right|_{L^\infty H^{s-\frac{1}{2}}} \left|\nabla X^{(n)}\right|_{L^\infty H^{s-\frac{1}{2}}} \\
&\leq C \left\|v^{(n)}\right\|_{L^2 H^{s+1}} \left\|\zeta^{(n)} - \zeta^{(n-1)}\right\|_{L^\infty H^{s}} \left\|X^{(n)}\right\|_{L^\infty H^{s+1}} \\
&\leq C(M) \left\|X^{(n)} - X^{(n-1)}\right\|_{L^\infty H^{s+1}} \leq C(M) T^{\frac{1}{4}} \left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}},
\end{aligned}$$

and

$$\begin{aligned}
\left|d_1\right|_{H^{\frac{s}{2}-\frac{1}{4}} L^2} &\leq C \Big(\left|\nabla v^{(n)}\right|_{H^{\frac{s}{2}-\frac{1}{4}} L^2} + 1 \Big) \\
&\qquad \times \left|\zeta^{(n)} - \zeta^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}} H^{\frac{1}{2}+\delta}} \Big(\left|\nabla X^{(n)}\right|_{H^{\frac{s}{2}-\frac{1}{4}} H^{\frac{1}{2}+\delta}} + 1 \Big) \\
&\leq C \left\|v^{(n)}\right\|_{H^{ht,s+1}} \left\|\zeta^{(n)} - \zeta^{(n-1)}\right\|_{H^{\frac{s}{2}-\frac{1}{4}} H^{1+\delta}} \left\|X^{(n)}\right\|_{H^{\frac{s}{2}-\frac{1}{4}} H^{2+\delta}} \\
&\leq C(M) \left\|X^{(n)} - X^{(n-1)}\right\|_{H^{\frac{s}{2}-\frac{1}{4}} H^{2+\delta}} \leq C(M) T^{\epsilon} \left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.
\end{aligned}$$

Above we have used (68). Similarly

$$\begin{aligned}
\left|d_2\right|_{L^2 H^{s-\frac{1}{2}}} &\leq C \left|\nabla v^{(n)}\right|_{L^2 H^{s-\frac{1}{2}}} \left|\zeta^{(n-1)}\right|_{L^\infty H^{s-\frac{1}{2}}} \left|\nabla X^{(n)} - \nabla X^{(n-1)}\right|_{L^\infty H^{s-\frac{1}{2}}} \\
&\leq C(M) T^{\frac{1}{4}} \left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}},
\end{aligned}$$

and

$$\begin{aligned}
\left|d_2\right|_{H^{\frac{s}{2}-\frac{1}{4}} L^2} &\leq C \Big(\left|\nabla v^{(n)}\right|_{H^{\frac{s}{2}-\frac{1}{4}} L^2} + 1 \Big) \Big(\left|\zeta^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}} H^{\frac{1}{2}+\delta}} + 1 \Big) \\
&\qquad\qquad\qquad \times \left|\nabla X^{(n)} - \nabla X^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}} H^{\frac{1}{2}+\delta}} \\
&\leq C(M) T^{\epsilon} \left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.
\end{aligned}$$

For d_3 we find:

$$\begin{aligned}
\left|d_3\right|_{L^2 H^{s-\frac{1}{2}}} &\leq C \left|\nabla(v^{(n)} - v^{(n-1)})\right|_{L^2 H^{s-\frac{1}{2}}} \left|\zeta^{(n-1)} - \mathbb{I}\right|_{L^\infty H^{s-\frac{1}{2}}} \left|\nabla X^{(n)}\right|_{L^\infty H^{s-\frac{1}{2}}} \\
&\leq C \left\|v^{(n)} - v^{(n-1)}\right\|_{L^2 H^{s+1}} \left\|\zeta^{(n-1)} - \mathbb{I}\right\|_{L^\infty H^{s}} \left\|X^{(n)}\right\|_{L^\infty H^{s+1}} \\
&\leq C(M) T^{\frac{1}{4}} \left\|v^{(n)} - v^{(n-1)}\right\|_{H^{ht,s+1}},
\end{aligned}$$

and

$$\begin{aligned}
\left|d_3\right|_{H^{\frac{s}{2}-\frac{1}{4}} L^2} &\leq C \left|\nabla(v^{(n)} - v^{(n-1)})\right|_{H^{\frac{s}{2}-\frac{1}{4}} L^2} \left|\zeta^{(n-1)} - \mathbb{I}\right|_{H^{\frac{s}{2}-\frac{1}{4}} H^{\frac{1}{2}+\delta}} \\
&\qquad \times \Big(\left|\nabla X^{(n)}\right|_{H^{\frac{s}{2}-\frac{1}{4}} H^{\frac{1}{2}+\delta}} + 1 \Big) \\
&\leq C \left\|v^{(n)} - v^{(n-1)}\right\|_{H^{ht,s+1}} \left\|\zeta^{(n-1)} - \mathbb{I}\right\|_{H^{\frac{s}{2}-\frac{1}{4}} H^{1+\delta}} \Big(\left\|X^{(n)}\right\|_{H^{\frac{s}{2}-\frac{1}{4}} H^{2+\delta}} + 1 \Big) \\
&\leq C(M) T^{\epsilon} \left\|v^{(n)} - v^{(n-1)}\right\|_{H^{ht,s+1}}.
\end{aligned}$$

Above we have used (68). Similarly

$$\left|d_4\right|_{L^2 H^{s-\frac{1}{2}}} \leq C(M) T^{\frac{1}{4}} \left\|v^{(n)} - v^{(n-1)}\right\|_{H^{ht,s+1}}, \text{ and } \left|d_4\right|_{H^{\frac{s}{2}-\frac{1}{4}} L^2} \leq C(M) T^{\epsilon} \left\|v^{(n)} - v^{(n-1)}\right\|_{H^{ht,s+1}}.$$

We use the splitting $h_{v*}^{(n)} - h_{v*}^{(n-1)} = d_1^* + \ldots + d_8^*$ with

$$d_1^* = (\nabla v^{(n)} (\zeta^{(n)} - \zeta^{(n-1)}) A \circ X^{(n)})^* A^{-1} \circ X^{(n)} \nabla_J X^{(n)} n_0,$$

$$d_2^* = (\nabla v^{(n)} \zeta^{(n-1)} (A \circ X^{(n)} - A \circ X^{(n-1)}))^* A^{-1} \circ X^{(n)} \nabla_J X^{(n)} n_0,$$

$$
\begin{aligned}
d_3^* &= (\nabla v^{(n)}\zeta^{(n-1)}A\circ X^{(n-1)})^*(A^{-1}\circ X^{(n)} - A^{-1}\circ X^{(n-1)}))\nabla_J X^{(n)} n_0,\\
d_4^* &= (\nabla v^{(n)}\zeta^{(n-1)}A\circ X^{(n-1)})^*A^{-1}\circ X^{(n-1)}(\nabla_J X^{(n)} - \nabla_J X^{(n-1)})n_0,\\
d_5^* &= (\nabla(v^{(n)} - v^{(n-1)})(\zeta^{(n-1)} - \mathbb{I})A\circ X^{(n-1)})^*A^{-1}\circ X^{(n-1)}\nabla_J X^{(n-1)} n_0,\\
d_6^* &= (\nabla(v^{(n)} - v^{(n-1)})(A\circ X^{(n-1)} - A))^*A^{-1}\circ X^{(n-1)}\nabla_J X^{(n-1)} n_0,\\
d_7^* &= (\nabla(v^{(n)} - v^{(n-1)})A)^*(A^{-1}\circ X^{(n-1)} - A^{-1})\nabla_J X^{(n-1)} n_0,\\
d_8^* &= (\nabla(v^{(n)} - v^{(n-1)})A)^*A^{-1}(\nabla_J X^{(n)} - \mathbb{I})n_0.
\end{aligned}
\tag{77}
$$

Then

$$
\begin{aligned}
|d_1^*|_{L^2H^{s-\frac12}} &\le C\left|\nabla v^{(n)}\right|_{L^2H^{s-\frac12}}\left|\zeta^{(n)} - \zeta^{(n-1)}\right|_{L^\infty H^{s-\frac12}}\left|X^{(n)}\right|^2_{L^\infty H^{s-\frac12}}\left|\nabla X^{(n)}\right|_{L^\infty H^{s-\frac12}}\\
&\le C\left\|v^{(n)}\right\|_{L^2H^{s+1}}\left\|\zeta^{(n)} - \zeta^{(n-1)}\right\|_{L^\infty H^s}\left\|X^{(n)}\right\|^3_{L^\infty H^{s+1}}\\
&\le C(M)\left\|X^{(n)} - X^{(n-1)}\right\|_{L^\infty H^{s+1}} \le C(M)T^{\frac14}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.
\end{aligned}
$$

and

$$
\begin{aligned}
|d_1^*|_{H^{\frac{s}{2}-\frac14}L^2} \le C\Big(&\left|\nabla v^{(n)}\right|_{H^{\frac{s}{2}-\frac14}L^2} + 1\Big)\left|\zeta^{(n)} - \zeta^{(n-1)}\right|_{H^{\frac{s}{2}-\frac14}H^{\frac12+\delta}}\\
&\times\Big(\left|X^{(n)}\right|^2_{H^{\frac{s}{2}-\frac14}H^{\frac12+\delta}} + 1\Big)\Big(\left|\nabla X^{(n)}\right|_{H^{\frac{s}{2}-\frac14}H^{\frac12+\delta}} + 1\Big)\\
&\le C(M)\left\|X^{(n)} - X^{(n-1)}\right\|_{H^{\frac{s}{2}-\frac14}H^{2+\delta}} \le C(M)T^{\epsilon}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.
\end{aligned}
$$

Above we have used (68). Similarly

$$
\begin{aligned}
|d_2^*|_{L^2H^{s-\frac12}} &\le C(M)\left|\nabla v^{(n)}\right|_{L^2H^{s-\frac12}}\left|\zeta^{(n-1)}\right|_{L^\infty H^{s-\frac12}}\left|X^{(n)} - X^{(n-1)}\right|_{L^\infty H^{s-\frac12}}\left\|X^{(n)}\right\|^2_{L^\infty H^{s+1}}\\
&\le C(M)T^{\frac14}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}},
\end{aligned}
$$

and

$$
\begin{aligned}
|d_2^*|_{H^{\frac{s}{2}-\frac14}L^2} \le C\Big(&\left|\nabla v^{(n)}\right|_{H^{\frac{s}{2}-\frac14}L^2} + 1\Big)\Big(\left|\zeta^{(n-1)}\right|_{H^{\frac{s}{2}-\frac14}H^{\frac12+\delta}} + 1\Big)\\
&\times\left|X^{(n)} - X^{(n-1)}\right|_{H^{\frac{s}{2}-\frac14}H^{\frac12+\delta}}\Big(\left\|X^{(n)}\right\|^2_{H^{\frac{s}{2}-\frac14}H^{2+\delta}} + 1\Big)\\
&\le C(M)T^{\epsilon}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.
\end{aligned}
$$

We proceed for d_3^* and d_4^* as for d_2^* to find

$$
\begin{aligned}
|d_3^*|_{L^2H^{s-\frac12}} + |d_4^*|_{L^2H^{s-\frac12}} &\le C(M)T^{\frac14}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}},\\
|d_3^*|_{H^{\frac{s}{2}-\frac14}L^2} + |d_4^*|_{H^{\frac{s}{2}-\frac14}L^2} &\le C(M)T^{\epsilon}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.
\end{aligned}
$$

A similar approach is used to find

$$
\begin{aligned}
|d_5^*|_{L^2H^{s-\frac12}} \le C(M)&\left|\nabla(v^{(n)} - v^{(n-1)})\right|_{L^2H^{s-\frac12}}\\
&\times\left|\zeta^{(n-1)} - \mathbb{I}\right|_{L^\infty H^{s-\frac12}}\Big(\left\|X^{(n-1)}\right\|^3_{L^\infty H^{s+1}} + 1\Big)\\
&\le C(M)\left\|v^{(n)} - v^{(n-1)}\right\|_{L^2H^{s+1}}\left\|\zeta^{(n-1)} - \mathbb{I}\right\|_{L^\infty H^s}\\
&\le C(M)T^{\frac14}\left\|w^{(n)} - w^{(n-1)}\right\|_{H^{ht,s+1}},
\end{aligned}
$$

and

$$
\begin{aligned}
|d_5^*|_{H^{\frac{s}{2}-\frac14}L^2} \le C&\left|\nabla(v^{(n)} - v^{(n-1)})\right|_{H^{\frac{s}{2}-\frac14}L^2}\\
&\times\left|\zeta^{(n-1)} - \mathbb{I}\right|_{H^{\frac{s}{2}-\frac14}H^{\frac12+\delta}}\Big(\left\|X^{(n-1)}\right\|^3_{H^{\frac{s}{2}-\frac14}H^{2+\delta}} + 1\Big)\\
&\le C(M)\left\|v^{(n)} - v^{(n-1)}\right\|_{H^{ht,s+1}}\left\|\zeta^{(n-1)} - \mathbb{I}\right\|_{H^{\frac{s}{2}-\frac14}H^{1+\delta}}\\
&\le C(M)T^{\epsilon}\left\|w^{(n)} - w^{(n-1)}\right\|_{H^{ht,s+1}}.
\end{aligned}
$$

It is possible to distribute the norms in a similar manner as in d_5 to conclude that

$$\begin{aligned}|d_6^*|_{L^2H^{s-\frac{1}{2}}} + |d_7^*|_{L^2H^{s-\frac{1}{2}}} + |d_8^*|_{L^2H^{s-\frac{1}{2}}} &\le C(M)T^{\frac{1}{4}} \left\|w^{(n)} - w^{(n-1)}\right\|_{H^{ht,s+1}},\\ |d_6^*|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} + |d_7^*|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} + |d_8^*|_{L^2H^{s-\frac{1}{2}}} &\le C(M)T^{\epsilon} \left\|w^{(n)} - w^{(n-1)}\right\|_{H^{ht,s+1}}.\end{aligned}$$

Next, we deal with $h_q^{(n)} - h_q^{(n-1)} = d_1^q + d_2^q + d_3^q + d_4^q$ where

$$\begin{aligned}d_1^q &= q^{(n-1)}(A^{-1}\circ X^{(n-1)} - A^{-1}\circ X^{(n)})\nabla_J X^{(n-1)} n_0,\\ d_2^q &= q^{(n-1)}A^{-1}\circ X^{(n)}(\nabla_J X^{(n-1)} - \nabla_J X^{(n)})n_0,\\ d_3^q &= (q^{(n-1)} - q^{(n)})(A^{-1}\circ X^{(n)} - A^{-1})\nabla_J X^{(n)} n_0,\\ d_4^q &= (q^{(n-1)} - q^{(n)})A^{-1}(\nabla_J X^{(n)} - \mathbb{I})n_0.\end{aligned} \tag{78}$$

We start as follows

$$\begin{aligned}|d_1^q|_{L^2H^{s-\frac{1}{2}}} &\le C\left|q^{(n-1)}\right|_{L^2H^{s-\frac{1}{2}}}\left|X^{(n)} - X^{(n-1)}\right|_{L^\infty H^{s-\frac{1}{2}}}\left|\nabla X^{(n-1)}\right|_{L^\infty H^{s-\frac{1}{2}}}\\ &\le C(M)T^{\frac{1}{4}}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}},\end{aligned}$$

and

$$\begin{aligned}|d_1^q|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} \le C\Big(\left|q^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} + 1\Big)&\left|X^{(n)} - X^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}\\ &\times(\left|\nabla X^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}} + 1)\\ \le C(M)T^{\epsilon}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.&\end{aligned}$$

Next

$$\begin{aligned}|d_2^q|_{L^2H^{s-\frac{1}{2}}} &\le C\left|q^{(n-1)}\right|_{L^2H^{s-\frac{1}{2}}}\left|X^{(n)}\right|_{L^\infty H^{s-\frac{1}{2}}}\left|\nabla X^{(n)} - \nabla X^{(n-1)}\right|_{L^\infty H^{s-\frac{1}{2}}}\\ &\le C(M)T^{\frac{1}{4}}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}},\end{aligned}$$

and

$$\begin{aligned}|d_2^q|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} \le C\Big(\left|q^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} + 1\Big)&\Big(\left|X^{(n)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}} + 1\Big)\\ &\times\left|\nabla X^{(n)} - \nabla X^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}}\\ \le C(M)T^{\epsilon}\left\|X^{(n)} - X^{(n-1)}\right\|_{F^{s+1}}.&\end{aligned}$$

The next term provides

$$\begin{aligned}|d_3^q|_{L^2H^{s-\frac{1}{2}}} &\le C\left|q^{(n)} - q^{(n-1)}\right|_{L^2H^{s-\frac{1}{2}}}\left|X^{(n)} - \alpha\right|_{L^\infty H^{s+\frac{1}{2}}}\left|\nabla X^{(n-1)}\right|_{L^\infty H^{s-\frac{1}{2}}}\\ &\le C(M)T^{\frac{1}{4}}\|q_w^{(n)} - q_w^{(n-1)}\|_{H_{pr}^{ht,s}},\end{aligned}$$

and

$$\begin{aligned}|d_3^q|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} \le C\left|q^{(n)} - q^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}L^2}&\left|X^{(n)} - \alpha\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{3}{2}+\delta}}\\ &\times(\left|\nabla X^{(n-1)}\right|_{H^{\frac{s}{2}-\frac{1}{4}}H^{\frac{1}{2}+\delta}} + 1)\\ \le C(M)T^{\epsilon}\|q_w^{(n)} - q_w^{(n-1)}\|_{H_{pr}^{ht,s}}.&\end{aligned}$$

As for d_3^q we find

$$\begin{aligned}|d_4^q|_{L^2H^{s-\frac{1}{2}}} &\le C(M)T^{\frac{1}{4}}\|q_w^{(n)} - q_w^{(n-1)}\|_{H_{pr}^{ht,s}},\\ |d_4^q|_{H^{\frac{s}{2}-\frac{1}{4}}L^2} &\le C(M)T^{\epsilon}\|q_w^{(n)} - q_w^{(n-1)}\|_{H_{pr}^{ht,s}}.\end{aligned}$$

Corollary 5.5 *By the contraction mapping principle, there is a unique fixed point, which is the solution of (11-14).*

6 Structural stability

Let us assume that (v, q, X) is a solution with initial data v_0 and Ω_0, where Ω_0 is the projection by P of a splash domain. We also assume that $(v'_\varepsilon, q'_\varepsilon, X'_\varepsilon)$ is a solution with initial data $v'_{\varepsilon\,0}$ and Ω_ε where Ω_ε is a translation of Ω_0 of size ε, i.e.,

$$\Omega_\varepsilon = \Omega_0 + \varepsilon b$$

where b is a constant vector, $|b| = 1$ and such that $P^{-1}(\Omega_\varepsilon)$ is a good domain. We define $(v_\varepsilon, q_\varepsilon, X_\varepsilon)$ in the following way

$$v_\varepsilon(\alpha, t) = v'_\varepsilon(\alpha + \varepsilon b, t), \qquad q_\varepsilon(\alpha, t) = q'_\varepsilon(\alpha + \varepsilon b, t), \quad X_\varepsilon(\alpha, t) = X'_\varepsilon(\alpha + \varepsilon b, t),$$

for $\alpha \in \Omega_0$. We choose $v'_{\varepsilon 0}$ to have $v_\varepsilon(\alpha, 0) = v_0(\alpha)$. Since the change of variables from the variables with prime to the variables without prime is just a constant translation, the functions $(v_\varepsilon, q_\varepsilon, X_\varepsilon)$ satisfy (11)-(14) and (15). However instead of (16) we have that $X_\varepsilon(\alpha, 0) = \alpha + \varepsilon b$.

We denote by $Q^2_\varepsilon(\alpha)$ and $A_\varepsilon(\alpha)$ the following functions:

$$Q^2_\varepsilon(\alpha) = Q^2(\alpha + \varepsilon b), \quad A_\varepsilon(\alpha) = A(\alpha + \varepsilon b).$$

Therefore, we have

$$\begin{aligned}
\partial_t v_\varepsilon - Q^2_\varepsilon \Delta v_\varepsilon + A^*_\varepsilon \nabla q_\varepsilon &= f_\varepsilon \text{ in } \Omega_0 \\
Tr(\nabla v_\varepsilon A_\varepsilon) &= g_\varepsilon \text{ in } \Omega_0 \\
(q_\varepsilon \mathbb{I} - ((\nabla v_\varepsilon A_\varepsilon) + (\nabla v_\varepsilon A_\varepsilon)^*)) A^{-1}_\varepsilon n_0 &= h_\varepsilon \text{ in } \partial\Omega_0,
\end{aligned}$$

where

$$\begin{aligned}
f_{\varepsilon,i} &= Q^2 \circ X_\varepsilon (\zeta_\varepsilon)_{kj} \partial_k((\zeta_\varepsilon)_{lj} \partial_l v_i) - Q^2_\varepsilon \Delta v_{\varepsilon,i} + (A_\varepsilon)_{ki} \partial_k q_\varepsilon - A_{ki} \circ X_\varepsilon (\zeta_\varepsilon)_{jk} \partial_j q_\varepsilon \\
g_\varepsilon &= Tr(\nabla v_\varepsilon A_\varepsilon) - Tr(\nabla v_\varepsilon \zeta_i A \circ X_\varepsilon) \\
h_\varepsilon &= q_\varepsilon A^{-1}_\varepsilon n_0 - q_\varepsilon A^{-1} \circ X_\varepsilon \nabla_J X_\varepsilon n_0 + (\nabla v_\varepsilon \zeta_\varepsilon A \circ X_\varepsilon + (\nabla v_\varepsilon \zeta_\varepsilon A \circ X_\varepsilon)^*) A^{-1} \circ X_\varepsilon \nabla_J X_\varepsilon n_0 \\
&\quad - (\nabla v_\varepsilon A_\varepsilon + (\nabla v_\varepsilon A_\varepsilon)^*) A^{-1}_\varepsilon n_0
\end{aligned}$$

We construct ϕ_ε in an analogous way: ensuring that $v_\varepsilon = w_\varepsilon + \phi_\varepsilon$, with $w_\varepsilon = \partial_t w_\varepsilon = 0$. This yields:

$$\psi_\varepsilon = v_0 + t \exp(-t^2)(Q^2_\varepsilon \Delta v_0 - A^*_\varepsilon \nabla q_{\phi,\varepsilon}),$$

where

$$\begin{aligned}
-Q^2_\varepsilon \Delta q_{\phi,\varepsilon} &= Tr(\nabla v_0 A_\varepsilon \nabla v_0 A_\varepsilon) \text{ in } \Omega_0 \\
q_{\phi,\varepsilon}(A^{-1}_\varepsilon n_0 \cdot A^{-1}_\varepsilon n_0) &= A^{-1}_\varepsilon n_0 (\nabla v_0 A_\varepsilon + (\nabla v_0 A_\varepsilon)^*) A^{-1}_\varepsilon n_0 \text{ in } \partial\Omega_0.
\end{aligned}$$

Thus, we have the following system:

$$\begin{aligned}
\partial_t w_\varepsilon - Q^2_\varepsilon \Delta w_\varepsilon + A^*_\varepsilon \nabla q_{w,\varepsilon} &= f_\varepsilon + f^L_{\phi,\varepsilon} \text{ in } \Omega_0, \\
Tr(\nabla w_\varepsilon A_\varepsilon) &= g_\varepsilon + g^L_{\phi,\varepsilon} \text{ in } \Omega_0, \\
(q_{w,\varepsilon} I - ((\nabla w_\varepsilon A_\varepsilon) + (\nabla w_\varepsilon A_\varepsilon)^*)) A^{-1}_\varepsilon n_0 &= h_\varepsilon + h^L_{\phi,\varepsilon} \text{ on } \partial\Omega_0,
\end{aligned}$$

where

$$\begin{aligned}
f^L_{\phi,\varepsilon} &= -\partial_t \phi_\varepsilon + Q^2_\varepsilon \Delta \phi_\varepsilon - A^*_\varepsilon \nabla q_\varepsilon \\
g^L_{\phi,\varepsilon} &= -Tr(\nabla \phi_\varepsilon A_\varepsilon) \\
h^L_{\phi,\varepsilon} &= -q_{\phi,\varepsilon} A^{-1}_\varepsilon n_0 + (\nabla \phi_\varepsilon A_\varepsilon + (\nabla \phi_\varepsilon A_\varepsilon)^*) A^{-1}_\varepsilon n_0.
\end{aligned} \tag{79}$$

The next step will be to compare both solutions (w, q_w, X) and $(w_\varepsilon, q_{w,\varepsilon}, X_\varepsilon)$. Subtracting one equation from the other:

$$\begin{aligned}\partial_t(w-w_\varepsilon) - Q^2\Delta(w-w_\varepsilon) + A^*\nabla(q_w - q_{w,\varepsilon}) &= F_\varepsilon\\ Tr(\nabla(w-w_\varepsilon)A) &= G_\varepsilon\\ ((q_w - q_{w,\varepsilon})I - ((\nabla(w-w_\varepsilon)A) + (\nabla(w-w_\varepsilon)A)^*))A^{-1}n_0 &= H_\varepsilon,\end{aligned}$$

where $F_\varepsilon, G_\varepsilon, H_\varepsilon$ are given by

$$F_\varepsilon = f - f_\varepsilon + f_\phi^L - f_{\phi,\varepsilon}^L - (Q^2 - Q_\varepsilon^2)\Delta w_\varepsilon + (A - A_\varepsilon)^*\nabla q_{w,\varepsilon} \quad (80)$$

$$G_\varepsilon = g - g_\varepsilon + g_\phi^L - g_{\phi,\varepsilon}^L - Tr(\nabla w_\varepsilon(A - A_\varepsilon)) \quad (81)$$

$$\begin{aligned}H_\varepsilon =& h - h_\varepsilon + h_\phi^L - h_{\phi,\varepsilon}^L - q_{w,\varepsilon}(A^{-1} - A_\varepsilon^{-1})n_0 - ((\nabla w_\varepsilon(A - A_\varepsilon)) + (\nabla w_\varepsilon(A - A_\varepsilon))^*)A_\varepsilon^{-1}n_0 \quad (82)\\ &- (\nabla w_\varepsilon A + (\nabla w_\varepsilon A)^*)(A^{-1} - A_\varepsilon^{-1})n_0.\end{aligned}$$

This means

$$(w - w_\varepsilon, q_w - q_{w,\varepsilon}) = L^{-1}(F_\varepsilon, G_\varepsilon, H_\varepsilon, 0).$$

Taking norms as in the previous section, we obtain the following estimates:

$$\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|q_w - q_{w,\varepsilon}\|_{H_{pr}^{ht,s}} \le C(\|F_\varepsilon\|_{H^{ht,s-1}} + \|G_\varepsilon\|_{\overline{H}^{ht,s}} + |H_\varepsilon|_{H^{ht,s-\frac{1}{2}}}).$$

Our goal is to prove the next lemmas:

Lemma 6.1 *We have the following estimate, for $\delta > 0$ small enough:*

$$\|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma} \le C\varepsilon + CT^\delta(\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma})$$

Lemma 6.2 *We have the following estimate, for $\delta > 0$ small enough:*

$$\begin{aligned}\|F_\varepsilon\|_{H^{ht,s-1}} + \|G_\varepsilon\|_{\overline{H}^{ht,s}} + |H_\varepsilon|_{H^{ht,s-\frac{1}{2}}} \le C\varepsilon + CT^\delta(&\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|q_w - q_{w,\varepsilon}\|_{H_{pr}^{ht,s}}\\ &+ \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma}).\end{aligned}$$

We remark that the constants in the previous lemmas are uniformly bounded if T is bounded above. Combining both inequalities, for $0 < T \le \frac{1}{(2C)^{1/\delta}}$:

$$\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|q_w - q_{w,\varepsilon}\|_{H_{pr}^{ht,s}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma} \le 2C\varepsilon$$

Thus, for $0 < T \le \frac{1}{(2C)^{1/\delta}}$:

$$\|X - X_\varepsilon\|_{L^\infty H^{s+1}} \le 2C\varepsilon,$$

and therefore both functions are as close as we want for a time that only depends on the local existence time of the solution, but it does not depend on ε. We therefore have

$$X(\Omega_0, t) \approx X_\varepsilon(\Omega_{0,\varepsilon}, t)$$

and for ε small enough there exists a t_s such that

$$P^{-1}(X_\varepsilon(\Omega_{0,\varepsilon}, t_s)) = P^{-1}(X_\varepsilon(\Omega_0 + \varepsilon b, t_s))$$

is a splash domain. Next, we will show the proofs of the lemmas:

Proof of Lemma 6.1: We recall that

$$\frac{dX_\varepsilon}{dt} = A\circ X_\varepsilon v_\varepsilon, \qquad\qquad X_\varepsilon(\alpha,0) = \alpha+\varepsilon b.$$

We can produce the following bounds:

$$\begin{aligned}\|X-X_\varepsilon\|_{H^{s+1}}(t) &\le C\varepsilon + \int_0^t \|A\circ Xv - A\circ X_\varepsilon v_\varepsilon\|_{H^{s+1}}(\tau)d\tau\\ &\le C\varepsilon + \|A\circ X - A\circ X_\varepsilon\|_{L^\infty H^{s+1}} t^{\frac12}\|v\|_{L^2H^{s+1}} + \|A\circ X_\varepsilon\|_{L^\infty H^{s+1}} t^{\frac12}\|v-v_\varepsilon\|_{L^2H^{s+1}}\\ &\le C\varepsilon + Ct^{\frac12}\|X-X_\varepsilon\|_{L^\infty H^{s+1}} + Ct^{\frac12}\|w-w_\varepsilon\|_{L^2H^{s+1}} + Ct^{\frac12}\|\phi-\phi_\varepsilon\|_{L^2H^{s+1}}\end{aligned}$$

We are only left to prove

$$\|\phi-\phi_\varepsilon\|_{L^2H^{s+1}} \le C\varepsilon.$$

After doing so, (we take $t\le 1$) we would obtain the estimate

$$\|X-X_\varepsilon\|_{L^\infty H^{s+1}} \le C\varepsilon + CT^{\frac12}(\|X-X_\varepsilon\|_{L^\infty H^{s+1}} + \|w-w_\varepsilon\|_{L^2H^{s+1}}). \tag{83}$$

Let's prove the estimate. We have that

$$\|A-A_\varepsilon\|_{H^r} \le C\varepsilon, \quad \|Q^2-Q_\varepsilon^2\|_{H^r} \le C\varepsilon, \text{ for all } r, \tag{84}$$

since Q and A are C^∞ functions. Thus

$$\begin{aligned}-\Delta(q_\phi - q_{\phi,\varepsilon}) &= \frac{1}{Q^2}Tr(\nabla v_0 A\nabla v_0 A) - \frac{1}{Q_\varepsilon^2}Tr(\nabla v_0 A_\varepsilon \nabla v_0 A_\varepsilon) \qquad \text{in } \Omega_0\\ q_\phi - q_{\phi,\varepsilon} &= \frac{A^{-1}n_0(\nabla v_0 A + (\nabla v_0 A)^*)A^{-1}n_0}{|A^{-1}n_0|^2} - \frac{A_\varepsilon^{-1}n_0((\nabla v_0 A_\varepsilon) + (\nabla v_0 A_\varepsilon)^*)A_\varepsilon^{-1}n_0}{|A_\varepsilon^{-1}n_0|^2} \qquad \text{on } \partial\Omega_0.\end{aligned}$$

This implies:

$$\|q_\phi - q_{\phi,\varepsilon}\|_{H^{r+1}} \le \|\Delta(q_\phi - q_{\phi,\varepsilon})\|_{H^{r-1}} + |q_\phi - q_{\phi,\varepsilon}|_{H^{r+\frac12}} \le C\varepsilon, \text{ for all } r\ge 0. \tag{85}$$

Using the definition of ϕ,ϕ_ε:

$$\phi-\phi_\varepsilon = t\exp(-t^2)((Q-Q_\varepsilon)\Delta v_0 - (A^*\nabla q_\phi - A_\varepsilon^*\nabla q_{\phi,\varepsilon}))$$

yields

$$\|\phi-\phi_\varepsilon\|_{L^\infty H^{s+1}} \le C\varepsilon,$$

for sufficiently smooth v_0.
To get the H^2H^γ estimate, we find that for all $t\le 1$:

$$\begin{aligned}\|X-X_\varepsilon\|_{H^2H^\gamma} &\le C\varepsilon + \left\|\int_0^t (A\circ Xv - A\circ X_\varepsilon v_\varepsilon)d\tau\right\|_{H^2H^\gamma}\\ &\le C\varepsilon + \left\|\int_0^t (A\circ Xv - A\circ X_\varepsilon v_\varepsilon) - (A-A_\varepsilon)v_0 d\tau\right\|_{H^2H^\gamma}\end{aligned}$$

At this point we can use Lemma 3.4 to get

$$
\begin{aligned}
\|X - X_\varepsilon\|_{H^2H^\gamma} &\leq C\varepsilon + C\|A \circ Xv - A \circ X_\varepsilon v_\varepsilon - (A - A_\varepsilon)v_0\|_{H^1H^\gamma} \\
&\leq C\varepsilon + C\|(A \circ X - A \circ X_\varepsilon)v\|_{H^1H^\gamma} + C\|A \circ X_\varepsilon(v - v_\varepsilon)\|_{H^1H^\gamma} \\
&\leq C\varepsilon + C\|X - X_\varepsilon\|_{H^1H^\gamma} + C\|v - v_\varepsilon\|_{H^1H^\gamma} \\
&\leq C\varepsilon + C\|X - X_\varepsilon\|_{H^1H^\gamma} + C\|w - w_\varepsilon\|_{H^1H^\gamma} + C\|\phi - \phi_\varepsilon\|_{H^1H^\gamma}.
\end{aligned}
$$

Now it is easy to find

$$
\|X - X_\varepsilon\|_{H^2H^\gamma} \leq C\varepsilon + CT^\delta\|X - X_\varepsilon\|_{H^2H^\gamma} + CT^\delta\|w - w_\varepsilon\|_{H^{ht,s+1}}, \tag{86}
$$

since $\|\phi - \phi_\varepsilon\|_{H^1H^\gamma} \leq C\varepsilon$. That way, the lemma is proved.

□

Proof of Lemma 6.2:

Part I: Estimates for F_ε:

We consider first in (80) the terms

$$
-(Q^2 - Q_\varepsilon^2)\Delta w_\varepsilon + (A - A_\varepsilon)^*\nabla q_{w,\varepsilon}.
$$

Using (84) it is easy to check that

$$
\|(Q^2 - Q_\varepsilon^2\Delta w_\varepsilon\|_{H^{ht,s-1}} + \|(A - A_\varepsilon)^*\nabla q_{w,\varepsilon}\|_{H^{ht,s-1}} \leq C\varepsilon.
$$

Concerning f, we split as in (64) (ignoring the superindices), with $f = f_w + f_\phi + f_q$. In an analogous way, in (80) we consider $f_\varepsilon = f_{w,\varepsilon} + f_{\phi,\varepsilon} + f_{q,\varepsilon}$.

We then split as:

$$
\begin{aligned}
f_w - f_{w,\varepsilon} &= Q^2 \circ X\zeta\partial(\zeta\partial w) - Q^2\Delta w - (Q^2 \circ X_\varepsilon\zeta_\varepsilon\partial(\zeta_\varepsilon\partial w_\varepsilon) - Q_\varepsilon^2\Delta w_\varepsilon) \\
&= d_1^e + d_2^e + d_3^e + d_4^e + d_5^e + d_6^e + d_{1,\varepsilon}^e,
\end{aligned}
$$

with

$$
\begin{aligned}
d_1^e &= (Q^2 \circ X - Q^2 \circ X_\varepsilon)\zeta\partial(\zeta\partial w), & d_2^e &= Q^2 \circ X_\varepsilon(\zeta - \zeta_\varepsilon)\partial(\zeta\partial w) \\
d_3^e &= Q^2 \circ X_\varepsilon\zeta_\varepsilon\partial((\zeta - \zeta_\varepsilon)\partial w), & d_4^e &= (Q^2 \circ X_\varepsilon - Q_\varepsilon^2)\zeta_\varepsilon\partial(\zeta_\varepsilon\partial(w - w_\varepsilon)) \\
d_5^e &= Q_\varepsilon^2(\zeta_\varepsilon - I)\partial(\zeta_\varepsilon\partial(w - w_\varepsilon)), & d_6^e &= Q_\varepsilon^2\partial((\zeta_\varepsilon - I)\partial(w - w_\varepsilon)) \\
d_{1,\varepsilon}^e &= (Q_\varepsilon^2 - Q^2)\Delta w
\end{aligned}
$$

As before, (84) yields

$$
\|d_{1,\varepsilon}^e\|_{H^{ht,s-1}} \leq C\varepsilon.
$$

To estimate d_j^e, $1 \leq j \leq 6$, we will compare the procedure with the one for d_j (69). It is easy to see that they are similar, by identifying X^n, ζ^n, w^n with X, ζ, w and $X^{n-1}, \zeta^{n-1}, w^{n-1}$ with $X_\varepsilon, \zeta_\varepsilon, w_\varepsilon$. As an illustration, we will discuss d_1^e with detail. As in the d_1 case, we find

$$
\|d_1^e\|_{L^2H^{s-1}} \leq C(M)\|X - X_\varepsilon\|_{L^\infty H^{s-1}} \leq C(M)\|X - X_\varepsilon\|_{L^\infty H^{s+1}}.
$$

Using (83), we obtain

$$
\|d_1^e\|_{L^2H^{s-1}} \leq C\varepsilon + CT^{\frac{1}{2}}(\|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|w - w_\varepsilon\|_{L^2H^{s+1}})
$$

Concerning the $H^{\frac{s-1}{2}}L^2$ estimate, we have for $\eta > 0$ small enough:

$$\begin{aligned}\|d_1^e\|_{H^{\frac{s-1}{2}}L^2} &\leq C(M)\|X - X_\varepsilon\|_{H^{\frac{s-1}{2}}H^{1+\eta}} \\ &\leq C(M)(\|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma}) \\ &\leq C\varepsilon + CT^\delta(\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma}),\end{aligned}$$

by using (83) and (86). Analogous estimates can be deduced for $d_j^e, 2 \leq j \leq 6$, achieving

$$\sum_{j=1}^{6}\|d_j^e\|_{H^{ht,s-1}} \leq C\varepsilon + CT^\delta(\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma})$$

We next consider

$$\begin{aligned}f_\phi - f_{\phi,\varepsilon} &= Q^2 \circ X\zeta\partial(\zeta\partial\phi) - Q^2\Delta\phi - (Q_\varepsilon^2 \circ X_\varepsilon\zeta_\varepsilon\partial(\zeta_\varepsilon\partial\phi_\varepsilon) - Q_\varepsilon^2\Delta\phi_\varepsilon) \\ &= d_7^e + d_8^e + d_9^e + d_{2,\varepsilon}^e + d_{3,\varepsilon}^e + d_{4,\varepsilon}^e,\end{aligned}$$

where

$$\begin{aligned}d_7^e &= (Q^2 \circ X - Q^2 \circ X_\varepsilon)\zeta\partial(\zeta\partial\phi), \qquad & d_8^e &= Q^2 \circ X_\varepsilon(\zeta - \zeta_\varepsilon)\partial(\zeta\partial\phi), \\ d_9^e &= Q^2 \circ X_\varepsilon\zeta_\varepsilon\partial((\zeta - \zeta_\varepsilon)\partial\phi), \qquad & d_{2,\varepsilon}^e &= Q^2 \circ X_\varepsilon\zeta_\varepsilon\partial(\zeta_\varepsilon\partial(\phi - \phi_\varepsilon)), \\ d_{3,\varepsilon}^e &= -(Q^2 - Q_\varepsilon^2)\Delta\phi, \qquad & d_{4,\varepsilon}^e &= -Q_\varepsilon^2\Delta(\phi - \phi_\varepsilon).\end{aligned}$$

We proceed as for d_1^e, d_2^e, d_3^e to find

$$\sum_{j=7}^{9}\|d_j\|_{H^{ht,s-1}} \leq C\varepsilon + CT^\delta(\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma})$$

Using the fact that $\phi - \phi_\varepsilon$ is small we obtain

$$\|d_{2,\varepsilon}^e\|_{H^{ht,s-1}} + \|d_{4,\varepsilon}^e\|_{H^{ht,s-1}} \leq C\varepsilon.$$

As we saw before, identity (84) provides

$$\|d_{3,\varepsilon}^e\|_{H^{ht,s-1}} \leq C\varepsilon.$$

We are therefore done with the terms $f_\phi - f_{\phi,\varepsilon}$. We now estimate $f_q - f_{q,\varepsilon}$. We have the following splitting:

$$f_q - f_{q,\varepsilon} = d_1^{q,e} + d_2^{q,e} + d_3^{q,e} + d_4^{q,e} + d_{5,\varepsilon}^{q,e},$$

with

$$\begin{aligned}d_1^{q,e} &= ((A \circ X_\varepsilon - A \circ X)\zeta_\varepsilon)^*\nabla q_\varepsilon, \qquad & d_2^{q,e} &= (A \circ X(\zeta_\varepsilon - \zeta))^*\nabla q_\varepsilon, \\ d_3^{q,e} &= (A \circ X - A\zeta) * (\nabla q_\varepsilon - \nabla q), \qquad & d_4^{q,e} &= (A(\zeta - I))^*(\nabla q_\varepsilon - \nabla q), \\ d_{5,\varepsilon}^{q,e} &= (A - A_\varepsilon)^*\nabla q_\varepsilon.\end{aligned}$$

We already know that

$$\|d_{5,\varepsilon}^{q,e}\|_{H^{ht,s-1}} \leq C\varepsilon$$

due to (84).

The proximity of $A^* \circ X_\varepsilon - A^* \circ X$ and $\zeta_\varepsilon - \zeta$ allows us to conclude in an analogous way that

$$\|d_1^{q,e}\|_{H^{ht,s-1}} + \|d_2^{q,e}\|_{H^{ht,s-1}} \leq C(M)\varepsilon + CT^\delta(\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2H^\gamma}).$$

Similarly to the case of d_3^q and d_4^q in (70), we can write

$$\begin{aligned}\|d_3^q\|_{H^{ht,s-1}} + \|d_4^q\|_{H^{ht,s-1}} &\leq CT^\delta\|\nabla q_\varepsilon - \nabla q\|_{H^{ht,s-1}}\\ &\leq CT^\delta(\|\nabla q_{w,\varepsilon} - \nabla q_w\|_{H^{ht,s-1}} + \|\nabla q_{\phi,\varepsilon} - \nabla q_\phi\|_{H^{ht,s-1}}).\end{aligned}$$

Via the estimate (85) we can control $\|\nabla q_{\phi,\varepsilon} - \nabla q_\phi\|_{H^{ht,s-1}}$ by $C\varepsilon$ which implies that

$$\|d_3^q\|_{H^{ht,s-1}} + \|d_4^q\|_{H^{ht,s-1}} \leq C\varepsilon + CT^\delta\|q_{w,\varepsilon} - q_w\|_{H^{ht,s}_{pr}}.$$

With this estimate we finish the control of $f - f_\varepsilon$. We are left to estimage $f_\phi^L - f_{\phi,\varepsilon}^L$ given in (60) and (79). Proceeding as before, the control for $\phi - \phi_\varepsilon, Q - Q_\varepsilon, A - A_\varepsilon$ makes us conclude that

$$\|f_\phi^L - f_{\phi,\varepsilon}^L\|_{H^{ht,s-1}} \leq C\varepsilon,$$

and we are done with F_ε.

Part II: Estimates for G_ε:

We consider first in (81) $Tr(\nabla w_\varepsilon(A - A_\varepsilon))$. This term can be estimated using (84) in a way such that

$$\|Tr(\nabla w_\varepsilon(A - A_\varepsilon))\|_{\overline{H}^{ht,s}} \leq C\varepsilon.$$

Next we consider $g - g_\varepsilon$. We first estimate the L^2H^s norm, and then we will estimate the $H^{\frac{s+1}{2}}H^{-1}$ one. We have that

$$g - g_\varepsilon = \sum_{j=1}^4 d_j^e + d_{1,\varepsilon}^e,$$

where

$$\begin{aligned} d_1^e &= -Tr(\nabla(v - v_\varepsilon)(\zeta - I)A \circ X), \qquad & d_2^e &= -Tr(\nabla v_\varepsilon(\zeta - \zeta_\varepsilon)A \circ X),\\ d_3^e &= -Tr(\nabla v_\varepsilon \zeta_\varepsilon(A \circ X - A \circ X_\varepsilon)), \qquad & d_4^e &= -Tr(\nabla(v - v_\varepsilon)(A \circ X - A),\\ d_{1,\varepsilon}^e &= -Tr(\nabla v_\varepsilon(A_\varepsilon - A)). \end{aligned}$$

It is easy to check using (84) that $\|d_{1,\varepsilon}^e\|_{L^2H^s} \leq C\varepsilon$. To estimate d_j^e we compare it with (71). In an analogous way we get

$$\begin{aligned}\|d_1^e\|_{L^2H^s} &\leq \|\nabla(w - w_\varepsilon)(\zeta - I)A \circ X\|_{L^2H^s} + \|\nabla(\phi - \phi_\varepsilon)(\zeta - I)A \circ X\|_{L^2H^s}\\ &\leq C\varepsilon + CT^\delta\|w - w_\varepsilon\|_{H^{ht,s+1}}\end{aligned}$$

For d_2^e, we have:

$$\begin{aligned}\|d_2^e\|_{L^2H^s} &\leq C\|\zeta - \zeta_\varepsilon\|_{L^\infty H^s} \leq C\|X - X_\varepsilon\|_{L^\infty H^{s+1}}\\ &\leq C\varepsilon + CT^{\frac{1}{2}}(\|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|w - w_\varepsilon\|_{L^2H^{s+1}}),\end{aligned}$$

using (83). The same procedure applied to d_3^e yields:

$$\|d_3^e\|_{L^2H^s} \leq C\|X - X_\varepsilon\|_{L^\infty H^s} \leq C\varepsilon + CT^{\frac{1}{2}}(\|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|w - w_\varepsilon\|_{L^2H^{s+1}})$$

We are only left with d_4^e, but in an analogous way we get

$$\begin{aligned}\|d_4^e\|_{L^2H^s} &\leq C\|v-v_\varepsilon\|_{H^{ht,s+1}}\|X-X_\varepsilon\|_{L^\infty H^s}\\ &\leq C\varepsilon + CT^{\frac{1}{4}}\|w-w_\varepsilon\|_{H^{ht,s+1}}.\end{aligned}$$

We now move on to the $H^{\frac{s+1}{2}}H^{-1}$ norm. We consider first

$$\|d_{1,\varepsilon}^e\|_{H^{\frac{s+1}{2}}H^{-1}} \leq C\varepsilon,$$

using (84). We now split $d_1^e = d_{11}^e + d_{12}^e + d_{13}^e$, where

$$\begin{aligned}d_{11}^e &= -\int_0^t Tr(\nabla(\partial_t(v-v_\varepsilon))(\zeta - I)A\circ X)d\tau\\ d_{12}^e &= -\int_0^t Tr(\nabla(v-v_\varepsilon)\partial_t(\zeta - I)A\circ X)d\tau\\ d_{13}^e &= -\int_0^t Tr(\nabla(v-v_\varepsilon)(\zeta - I)\partial_t(A\circ X)d\tau\end{aligned}$$

One obtains

$$\begin{aligned}\|d_{11}^e\|_{H^{\frac{s+1}{2}}L^2} &\leq \|\nabla(\partial_t(v-v_\varepsilon))(\zeta-I)A\circ X\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\leq C\|\nabla(\partial_t(v-v_\varepsilon))\|_{H^{\frac{s-1}{2}}H^{-1}}\|\zeta - I\|_{H^{\frac{s-1}{2}}H^{1+\delta}}\\ &\leq CT^\delta\|v-v_\varepsilon\|_{H^{ht,s+1}} \leq C\varepsilon + CT^\delta\|w-w_\varepsilon\|_{H^{ht,s+1}}\end{aligned}$$

Analogously to d_{12} in (72):

$$\begin{aligned}\|d_{12}^e\|_{H^{\frac{s+1}{2}}H^{-1}} &\leq C\|\nabla(v-v_\varepsilon)\partial_t\zeta A\circ X\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\leq C\|\nabla(v-v_\varepsilon)\|_{H^{\frac{s-1}{2}}H^1}\|\partial_t\zeta - \partial_t\zeta|_{t=0}\|_{H^{\frac{s-1}{2}}L^2} + C\|\nabla(v-v_\varepsilon)\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\leq CT^\delta\|v-v_\varepsilon\|_{H^{ht,s+1}}\\ &\leq C\varepsilon + CT^\delta\|w-w_\varepsilon\|_{H^{ht,s+1}}\end{aligned}$$

The term d_{13}^e is easier to estimate, yielding:

$$\|d_{13}^e\|_{H^{\frac{s+1}{2}}H^{-1}} \leq C\varepsilon + CT^\delta\|w-w_\varepsilon\|_{H^{ht,s+1}}$$

In d_2^e we split further: $d_2^e = d_{21}^e + d_{22}^e + d_{23}^e + d_{2,\varepsilon}^e$, where

$$\begin{aligned}d_{21}^e &= -Tr\left(\int_0^t \nabla\partial_t v_\varepsilon\left(\zeta - \zeta_\varepsilon - \tau e^{-\tau^2}\nabla((A-A_\varepsilon)v_0)\right)A\circ X d\tau\right)\\ d_{22}^e &= -Tr\left(\int_0^t \nabla v_\varepsilon\partial_t\left(\zeta - \zeta_\varepsilon - \tau e^{-\tau^2}\nabla((A-A_\varepsilon)v_0)\right)A\circ X d\tau\right)\\ d_{23}^e &= -Tr\left(\int_0^t \nabla v_\varepsilon(\zeta - \zeta_\varepsilon - \tau e^{-\tau^2})\nabla((A-A_\varepsilon)v_0)\partial_t(A\circ X)d\tau\right)\\ d_{2,\varepsilon}^e &= -Tr\left(\nabla v_\varepsilon t e^{-t^2}\nabla((A-A_\varepsilon)v_0)A\circ X\right)\end{aligned}$$

Using (84) it is easy to conclude that

$$\|d_{2,\varepsilon}^e\|_{H^{\frac{s+1}{2}}H^{-1}} \leq C\varepsilon.$$

As for d_2 in (73), we have that

$$\begin{aligned}\sum_{j=1}^{3}\|d^e_{2,j}\|_{H^{\frac{s+1}{2}}H^{-1}} \leq{}& \|\nabla\partial_t v_\varepsilon(\zeta-\zeta_\varepsilon - te^{-t^2}\nabla((A-A_\varepsilon)v_0))A\circ X\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &+\|\nabla v_\varepsilon\partial_t(\zeta-\zeta_\varepsilon - te^{-t^2}\nabla((A-A_\varepsilon)v_0))A\circ X\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &+\|\nabla v_\varepsilon(\zeta-\zeta_\varepsilon - te^{-t^2}\nabla((A-A_\varepsilon)v_0))\partial_t A\circ X\|_{H^{\frac{s-1}{2}}H^{-1}},\end{aligned}$$

and therefore

$$\begin{aligned}\sum_{j=1}^{3}\|d^e_{2,j}\|_{H^{\frac{s+1}{2}}H^{-1}} \leq{}& C(1+\|\nabla\partial_t v_\varepsilon\|_{H^{\frac{s-1}{2}}H^{-1}})\|\zeta-\zeta_\varepsilon - te^{-t^2}\nabla((A-A_\varepsilon)v_0)\|_{H^{\frac{s-1}{2}}H^{1+\eta}}\\ &+C(1+\|\nabla v_\varepsilon\|_{H^{\frac{s-1}{2}}H^{-1}})\|\partial_t\zeta-\partial_t\zeta_\varepsilon-\partial_t(te^{-t^2}\nabla((A-A_\varepsilon)v_0))\|_{H^{\frac{s-1}{2}}L^2}\\ &+C(1+\|\nabla v_\varepsilon\|_{H^{\frac{s-1}{2}}H^{-1}})\|\zeta-\zeta_\varepsilon - te^{-t^2}\nabla((A-A_\varepsilon)v_0)\|_{H^{\frac{s-1}{2}}L^2}\\ &\leq C\varepsilon + CT^\delta(\|w-w_\varepsilon\|_{H^{ht,s+1}}+\|X-X_\varepsilon\|_{L^\infty H^{s+1}}+\|X-X_\varepsilon\|_{H^2H^\gamma}),\end{aligned}$$

with $\eta>0$ sufficiently small. In d^e_3 we split it in the following way:

$$d^e_3=\sum_{j=1}^{3}d^e_{3,j}+d^e_{3,\varepsilon},$$

where

$$\begin{aligned}d^e_{31}&=-\int_0^t Tr(\nabla\partial_t v_\varepsilon\zeta_\varepsilon(A\circ X-A\circ X_\varepsilon-\chi))d\tau\\ d^e_{32}&=-\int_0^t Tr(\nabla v_\varepsilon\partial_t\zeta_\varepsilon(A\circ X-A\circ X_\varepsilon-\chi))d\tau\\ d^e_{33}&=-\int_0^t Tr(\nabla v_\varepsilon\zeta_\varepsilon\partial_t(A\circ X-A\circ X_\varepsilon-\chi))d\tau\\ d^e_{3,\varepsilon}&=-Tr(\nabla v_\varepsilon\zeta_\varepsilon\chi)\\ \chi&=A-A_\varepsilon-\frac{d}{dt}\left.(A\circ X-A\circ X_\varepsilon)\right|_{t=0}te^{-t^2}.\end{aligned}$$

We compare here $d^e_{3,j}$ with $d_{3,j}$ in (74) and in an analogous way we can bound

$$\begin{aligned}\sum_{j=1}^{3}\|d^e_{3,j}\|_{H^{\frac{s+1}{2}}H^{-1}} \leq{}& \|\nabla\partial_t v_\varepsilon\zeta_\varepsilon(A\circ X-A\circ X_\varepsilon-\chi(t))\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &+\|\nabla v_\varepsilon\partial_t\zeta_\varepsilon(A\circ X-A\circ X_\varepsilon-\chi(t))\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &+\|\nabla v_\varepsilon\zeta_\varepsilon\partial_t(A\circ X-A\circ X_\varepsilon-\chi(t))\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\leq C\varepsilon + CT^\delta(\|w-w_\varepsilon\|_{H^{ht,s+1}}+\|X-X_\varepsilon\|_{L^\infty H^{s+1}}+\|X-X_\varepsilon\|_{H^2H^\gamma}).\end{aligned}$$

We further split for $d^e_{3,\varepsilon}$ into $d^e_{3,\varepsilon}=d^{e,1}_{3,\varepsilon}+d^{e,2}_{3,\varepsilon}$, where

$$\begin{aligned}d^{e,1}_{3,\varepsilon}&=-Tr(\nabla v_\varepsilon(\zeta_\varepsilon-\mathbb{I}-\partial_t\zeta_\varepsilon|_{t=0}te^{-t^2})\chi(t))\\ d^{e,2}_{3,\varepsilon}&=-Tr(\nabla v_\varepsilon(\mathbb{I}+\partial_t\zeta_\varepsilon|_{t=0}te^{-t^2})\chi(t))\end{aligned}$$

Using that decomposition, $d^{e,1}_{3,\varepsilon}$ is estimated as before to get

$$\|d^{e,1}_{3,\varepsilon}\|_{H^{\frac{s+1}{2}}H^{-1}} \leq C\varepsilon.$$

The term $d^{e,2}_{3,\varepsilon}$ is estimated easily using (84) directly. We have finished with d^e_3. Finally, to deal with d^e_4 we proceed as for d_4 in (75) which would yield

$$\begin{aligned}\|d^e_4\|_{H^{\frac{s+1}{2}}H^{-1}} &\leq \|\partial_t(\nabla(v-v_\varepsilon))(A\circ X - A)\|_{H^{\frac{s-1}{2}}H^{-1}} + \|\nabla(v-v_\varepsilon)\partial_t(A\circ X)\|_{H^{\frac{s-1}{2}}H^{-1}}\\ &\leq C\varepsilon + CT^\delta(\|w-w_\varepsilon\|_{H^{ht,s+1}} + \|X-X_\varepsilon\|_{L^\infty H^{s+1}} + \|X-X_\varepsilon\|_{H^2H^\gamma}),\end{aligned}$$

by using the splitting $v-v_\varepsilon = (\phi-\phi_\varepsilon)+(w-w_\varepsilon)$. This finishes with the control over d^e_4 and therefore over $g-g_\varepsilon$. We are only left with the estimate of $g^L - g^L_{\phi,\varepsilon}$ given in (60) and (79), but the proximity between ϕ and ϕ_ε as well as between A and A_ε gives us

$$\|g-g_\varepsilon\|_{H^{\frac{s+1}{2}}H^{-1}} \leq C\varepsilon.$$

Part III: Estimates for H_ε:

We first consider in (82) the term

$$\begin{aligned}\overline{H}_\varepsilon &= -q_{w,\varepsilon}(A^{-1}-A^{-1}_\varepsilon)n_0 - ((\nabla w_\varepsilon(A-A_\varepsilon)) + (\nabla w_\varepsilon(A-A_\varepsilon))^*)A^{-1}_\varepsilon n_0\\ &\quad - ((\nabla w_\varepsilon A) + (\nabla w_\varepsilon A)^*)(A^{-1}-A^{-1}_\varepsilon)n_0\end{aligned}$$

First estimate in (84) yields

$$|\overline{H}_\varepsilon|_{H^{ht,s-\frac{1}{2}}} \leq C\varepsilon.$$

Next, the use of (84), (85) and the smallness of $\phi-\phi_\varepsilon$ allows us to obtain

$$|h^L_\phi - h^L_{\phi,\varepsilon}|_{H^{ht,s-\frac{1}{2}}} \leq C\varepsilon.$$

It remains to deal with $h-h_\varepsilon$. As we did in (67), we split $h = h_v + h_{v^*} + h_q$, and similarly $h_\varepsilon = h_{v,\varepsilon} + h_{v^*,\varepsilon} + h_{q,\varepsilon}$. We estimate first $h_v - h_{v,\varepsilon}$ using the splitting

$$h_v - h_{v,\varepsilon} = d^e_1 + d^e_2 + d^e_3 + d^e_4,$$

where

$$\begin{aligned} d^e_1 &= \nabla v(\zeta-\zeta_\varepsilon)\nabla_J X n_0, & d^e_2 &= \nabla v\zeta_\varepsilon(\nabla_J X - \nabla_J X_\varepsilon)n_0,\\ d^e_3 &= \nabla(v-v_\varepsilon)(\zeta_\varepsilon - I)\nabla_J X_\varepsilon n_0, & d^e_4 &= \nabla(v-v_\varepsilon)(\nabla_J X_\varepsilon - I)n_0,\end{aligned}$$

in a similar way to (76). As before, we are able to bound as follows:

$$\begin{aligned}|d^e_1|_{H^{ht,s-\frac{1}{2}}} + |d^e_2|_{H^{ht,s-\frac{1}{2}}} &\leq C(\|X-X_\varepsilon\|_{L^\infty H^{s+1}} + \|X-X_\varepsilon\|_{H^2H^\gamma})\\ &\leq C\varepsilon + CT^\delta(\|q_{w,\varepsilon}-q_w\|_{H^{ht,s+1}} + \|X-X_\varepsilon\|_{L^\infty H^{s+1}} + \|X-X_\varepsilon\|_{H^2H^\gamma}),\end{aligned}$$

where we have used Lemma 6.1. Thus, we have obtained the appropriate estimate. Repeating the procedure in the splitting (76) to d_3 and d_4, we find

$$|d^e_3|_{H^{ht,s-\frac{1}{2}}} + |d^e_4|_{H^{ht,s-\frac{1}{2}}} \leq CT^\delta\|v-v_\varepsilon\|_{H^{ht,s+1}} \leq C\varepsilon + CT^\delta\|w-w_\varepsilon\|_{H^{ht,s+1}}.$$

We split further for $h_{v^*} - h_{v^*,\varepsilon} = \sum_{j=1}^{8} d^{*,e}_j + \sum_{j=1}^{2} d^{*,e}_{j,\varepsilon}$, where

$$
\begin{aligned}
d_1^{*,e} &= (\nabla v(\zeta - \zeta_\varepsilon) A \circ X)^* A^{-1} \circ X \nabla_J X n_0 \\
d_2^{*,e} &= (\nabla v \zeta_\varepsilon (A \circ X - A \circ X_\varepsilon))^* A^{-1} \circ X \nabla_J X n_0 \\
d_3^{*,e} &= (\nabla v \zeta_\varepsilon A \circ X_\varepsilon)^* (A^{-1} \circ X - A^{-1} \circ X_\varepsilon) \nabla_J X n_0 \\
d_4^{*,e} &= (\nabla v \zeta_\varepsilon A \circ X_\varepsilon)^* A^{-1} \circ X_\varepsilon (\nabla_J X - \nabla_J X_\varepsilon) n_0 \\
d_5^{*,e} &= (\nabla (v - v_\varepsilon)(\zeta_\varepsilon - I) A \circ X_\varepsilon)^* A^{-1} \circ X_\varepsilon \nabla_J X_\varepsilon n_0 \\
d_6^{*,e} &= (\nabla (v - v_\varepsilon)(A \circ X_\varepsilon - A_\varepsilon))^* A^{-1} \circ X_\varepsilon \nabla_J X_\varepsilon n_0 \\
d_7^{*,e} &= (\nabla (v - v_\varepsilon) A_\varepsilon)^* (A^{-1} \circ X_\varepsilon - A_\varepsilon^{-1}) \nabla_J X_\varepsilon n_0 \\
d_8^{*,e} &= (\nabla (v - v_\varepsilon) A_\varepsilon)^* A_\varepsilon^{-1} (\nabla_J X_\varepsilon - I) n_0 \\
d_{1,\varepsilon}^{*,e} &= (\nabla v (A_\varepsilon - A))^* A_\varepsilon^{-1} n_0 \\
d_{2,\varepsilon}^{*,e} &= (\nabla v A)^* (A_\varepsilon^{-1} - A^{-1}) n_0
\end{aligned}
$$

The first estimate in (84) gives us

$$|d_{1,\varepsilon}^{*,e}|_{H^{ht,s-\frac{1}{2}}} + |d_{2,\varepsilon}^{*,e}|_{H^{ht,s-\frac{1}{2}}} \leq C\varepsilon,$$

and comparing $d_j^{*,e}$ with (77), using the previous techniques

$$\sum_{j=1}^{8} |d_j^{*,e}|_{H^{ht,s-\frac{1}{2}}} \leq C\varepsilon + CT^\delta (\|w - w_\varepsilon\|_{H^{ht,s+1}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2 H^\gamma}).$$

To finish, we consider $h_q - h_{q,\varepsilon} = \sum_{j=1}^4 d_j^{q,e} + d_{3,\varepsilon}^{q,e}$, where

$$
\begin{aligned}
d_1^{q,e} &= q_\varepsilon (A^{-1} \circ X_\varepsilon - A^{-1} \circ X) \nabla_J X_\varepsilon n_0 \\
d_2^{q,e} &= q_\varepsilon A^{-1} \circ X (\nabla_J X_\varepsilon - \nabla_J X) n_0 \\
d_3^{q,e} &= (q_\varepsilon - q)(A^{-1} \circ X - A^{-1}) \nabla_J X n_0 \\
d_4^{q,e} &= (q_\varepsilon - q) A^{-1} (\nabla_J - I) X n_0 \\
d_{3,\varepsilon}^{q,e} &= q_\varepsilon (A^{-1} - A_\varepsilon^{-1}) n_0.
\end{aligned}
$$

We can identify this decomposition with the one in (78), in such a way that the same estimates yield

$$\sum_{j=1}^{4} |d_j^{q,e}|_{H^{ht,s-\frac{1}{2}}} + |d_{3,\varepsilon}^{q,e}|_{H^{ht,s-\frac{1}{2}}} \leq C\varepsilon + CT^\delta (\|q_w - q_{w,\varepsilon}\|_{H^{ht,s-1}} + \|X - X_\varepsilon\|_{L^\infty H^{s+1}} + \|X - X_\varepsilon\|_{H^2 H^\gamma})$$

□

7 Setting the right initial normal velocity

We consider the following parametrization of the boundary of Ω:

$$z(s), \quad |z_s(s)| = 1,$$

we also consider a small enough neighbourhood of the boundary, U. In U one can use the coordinates (s, λ) given by

$$x(s, \lambda) = z(s) + \lambda z_s^\perp(s).$$

The stream function ψ, in U,will be given by

$$
\begin{aligned}
\overline{\psi}(s, \lambda) =& \psi_0(s) + \psi_1(s)\lambda + \frac{1}{2}\psi_2(s)\lambda^2 \\
\psi(x(s, \lambda)) =& \overline{\psi}(s, \lambda).
\end{aligned}
$$

Then we can extend in a smooth way ψ to the rest of the domain Ω and take $v_0 = \nabla^\perp \psi$. v_0 is clearly divergence free.

The initial velocity $v_0(x)$ must satisfy

$$t\left(\nabla v_0 + \nabla v_0^*\right)\big|_{\partial\Omega} n = 0,$$

where t and n are the tangential and normal vectors to the boundary of Ω respectively.

If T and N are an extension of t and n to U respectively, we can write

$$\left(T\left(\nabla v_0 + \nabla v_0^*\right) N\right)|_{\partial\Omega} = 0.$$

We will take

$$\begin{aligned} T(s,\lambda) =& x_s(s,\lambda) = z_s(s) + \lambda z_{ss}^\perp(s) = (1-\lambda\kappa(s))z_s(s) \\ N(s,\lambda) =& x_\lambda(s,\lambda) = z_s(s)^\perp, \end{aligned}$$

where

$$\kappa(s) = z_{ss}(s)\cdot z_s^\perp,$$

and we notice that

$$\nabla\psi\circ x(s,\lambda) = \frac{1}{1-\kappa\lambda}\overline{\psi}_s\, z_s + \overline{\psi}_\lambda\, z_s^\perp.$$

By defining

$$\overline{v_0}(s,\lambda) = v_0\circ x(s,\lambda)$$

we have that

$$\overline{v_0}(s,\lambda) = \frac{1}{1-\kappa\lambda}\overline{\psi}_s\, z_s^\perp - \overline{\psi}_\lambda\, z_s.$$

Now we can compute that

$$\begin{aligned} T^i\partial_i v_0^j \circ x(s,\lambda) =& \partial_s \overline{v_0}^j(s,\lambda) \\ N^j\partial_j v_0^i \circ x(s,\lambda) =& \partial_\lambda \overline{v_0}^i(s,\lambda). \end{aligned}$$

And then

$$\begin{aligned} T^i\partial_i v_0^j \circ x(s,\lambda) N^j &= \partial_s\left(\overline{v_0}\cdot N\right) - \overline{v_0}\cdot N_s \\ N^j\partial_j v_0^i \circ x(s,\lambda) T^i &= \partial_\lambda\left(\overline{v_0}\cdot T\right) - \overline{v_0}\cdot T_\lambda. \end{aligned}$$

But

$$\begin{aligned} T_\lambda =& z_{ss}^\perp(s) \\ N_s =& z_{ss}^\perp(s) = -\kappa(s) z_s(s). \end{aligned}$$

In addition

$$\overline{v_0}\cdot N = \frac{1}{1-\lambda\kappa}\overline{\psi}_s.$$

Then

$$\partial_s(\overline{v_0}\cdot N)|_{\lambda=0} = \partial_s^2\overline{\psi}(s,0).$$

Also

$$\overline{v_0}\cdot T = -(1-\lambda\kappa)\overline{\psi}_\lambda.$$

Therefore

$$\partial_\lambda(\overline{v_0}\cdot T)|_{\lambda=0} = \kappa(s)\partial_\lambda\overline{\psi}(s,0) - \partial_\lambda^2\overline{\psi}(s,0).$$

Finally

$$(\overline{v_0}\cdot T_\lambda)|_{\lambda=0} = \overline{v_0}\cdot N_s|_{\lambda=0} = \kappa\overline{\psi}_\lambda(s,0).$$

Thus, we have that

$$\partial_s^2\overline{\psi}(s,0) - \kappa\partial_\lambda\overline{\psi}(s,0) - \partial_\lambda^2\overline{\psi}(s,0) = 0.$$

Taking $\psi_1 = 0$ yields

$$\partial_s^2\psi_0 = \psi_2(s).$$

Just to conclude we notice that $v_0\cdot n|_{\partial\Omega} = \partial_s\psi_0(s)$. We first pick up ψ_0 in order to choose the normal component of the velocity at the interface and then ψ_2 to satisfy the continuity of the stress tensor.

Acknowledgements

AC, DC, CF, FG and JGS were partially supported by the grant MTM2011-26696 (Spain) and ICMAT Severo Ochoa project SEV-2011-0087. AC was partially supported by the Ramón y Cajal program RyC-2013-14317 and ERC grant 307179-GFTIPFD. CF was partially supported by NSF grant DMS-1265524. FG was partially supported by the Ramón y Cajal program RyC-2010-07094 and by P12-FQM-2466 grant from Junta de Andalucía, Spain.

References

[1] H. Abels. The initial-value problem for the Navier-Stokes equations with a free surface in L^q-Sobolev spaces. *Adv. Differential Equations*, 10(1):45–64, 2005.

[2] S. Agmon, A. Douglis, and L. Nirenberg. Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II. *Comm. Pure Appl. Math.*, 17:35–92, 1964.

[3] D. M. Ambrose and N. Masmoudi. The zero surface tension limit of two-dimensional water waves. *Comm. Pure Appl. Math.*, 58(10):1287–1315, 2005.

[4] H. Bae. Solvability of the free boundary value problem of the Navier-Stokes equations. *Discrete Contin. Dyn. Syst.*, 29(3):769–801, 2011.

[5] J. T. Beale. The initial value problem for the Navier-Stokes equations with a free surface. *Comm. Pure Appl. Math.*, 34(3):359–392, 1981.

[6] J. T. Beale. Large-time regularity of viscous surface waves. *Arch. Rational Mech. Anal.*, 84(4):307–352, 1983/84.

[7] J. T. Beale and T. Nishida. Large-time behavior of viscous surface waves. In *Recent topics in nonlinear PDE, II (Sendai, 1984)*, volume 128 of *North-Holland Math. Stud.*, pages 1–14. North-Holland, Amsterdam, 1985.

[8] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and J. Gómez-Serrano. Finite time singularities for water waves with surface tension. *Journal of Mathematical Physics*, 53(11):115622–115622, 2012.

[9] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and J. Gómez-Serrano. Finite time singularities for the free boundary incompressible Euler equations. *Ann. of Math. (2)*, 178(3):1061–1134, 2013.

[10] A. Córdoba, D. Córdoba, and F. Gancedo. Interface evolution: water waves in 2-D. *Adv. Math.*, 223(1):120–173, 2010.

[11] D. Cordoba, A. Enciso, and N. Grubic. Splash and almost-splash stationary solutions to the Euler equations. *Arxiv preprint arXiv:1412.7382*, 2014.

[12] D. Coutand and S. Shkoller. Unique solvability of the free-boundary Navier-Stokes equations with surface tension. *arXiv preprint math/0212116*, 2002.

[13] D. Coutand and S. Shkoller. On the Finite-Time Splash and Splat Singularities for the 3-D Free-Surface Euler Equations. *Comm. Math. Phys.*, 325(1):143–183, 2014.

[14] D. Coutand and S. Shkoller. On the impossibility of finite-time splash singularities for vortex sheets. *Arxiv preprint arXiv:1407.1479*, 2014.

[15] D. Coutand and S. Shkoller. 2015. Personal Communication.

[16] R. Danchin and P. B. Mucha. A Lagrangian approach for the incompressible Navier-Stokes equations with variable density. *Comm. Pure Appl. Math.*, 65(10):1458–1480, 2012.

[17] R. Danchin and P. B. Mucha. Incompressible flows with piecewise constant density. *Arch. Ration. Mech. Anal.*, 207(3):991–1023, 2013.

[18] T. Elgindi and D. Lee. Uniform regularity for free-boundary Navier-Stokes equations with surface tension. *Arxiv preprint arXiv:1403.0980*, 2014.

[19] C. Fefferman, A. D. Ionescu, and V. Lie. On the absence of "splash" singularities in the case of two-fluid interfaces. *arXiv preprint arXiv:1312.2917*, 2013.

[20] Y. Guo and I. Tice. Almost exponential decay of periodic viscous surface waves without surface tension. *Arch. Ration. Mech. Anal.*, 207(2):459–531, 2013.

[21] Y. Guo and I. Tice. Decay of viscous surface waves without surface tension in horizontally infinite domains. *Anal. PDE*, 6(6):1429–1533, 2013.

[22] Y. Guo and I. Tice. Local well-posedness of the viscous surface wave problem without surface tension. *Anal. PDE*, 6(2):287–369, 2013.

[23] Y. Hataya. Decaying solution of a Navier-Stokes flow without surface tension. *J. Math. Kyoto Univ.*, 49(4):691–717, 2009.

[24] J. Huang, M. Paicu, and P. Zhang. Global well-posedness of incompressible inhomogeneous fluid systems with bounded density or non-Lipschitz velocity. *Arch. Ration. Mech. Anal.*, 209(2):631–682, 2013.

[25] D. Lannes. *The Water Waves Problem: Mathematical Analysis and Asymptotics.* Mathematical Surveys and Monographs. Amer Mathematical Society, 2013.

[26] X. Liao and P. Zhang. On the global regularity of 2-d density patch for inhomogeneous incompressible viscous flow. *Arxiv preprint arXiv:1503.04898*, 2015.

[27] J.-L. Lions and E. Magenes. *Non-homogeneous boundary value problems and applications. Vol. I.* Springer-Verlag, New York-Heidelberg, 1972. Translated from the French by P. Kenneth, Die Grundlehren der mathematischen Wissenschaften, Band 181.

[28] N. Masmoudi and F. Rousset. Uniform regularity and vanishing viscosity limit for the free surface Navier-Stokes equations. *Arxiv preprint arXiv:1202.0657*, 2012.

[29] P. Neff. On Korn's first inequality with non-constant coefficients. *Proceedings of the Royal Society of Edinburgh, Section: A Mathematics*, 132:221–243, 2 2002.

[30] T. Nishida, Y. Teramoto, and H. Yoshihara. Global in time behavior of viscous surface waves: horizontally periodic motion. *J. Math. Kyoto Univ.*, 44(2):271–323, 2004.

[31] M. Paicu, P. Zhang, and Z. Zhang. Global unique solvability of inhomogeneous Navier-Stokes equations with bounded density. *Comm. Partial Differential Equations*, 38(7):1208–1234, 2013.

[32] V. A. Solonnikov. Solvability of the problem of the motion of a viscous incompressible fluid that is bounded by a free surface. *Izv. Akad. Nauk SSSR Ser. Mat.*, 41(6):1388–1424, 1448, 1977.

[33] V. A. Solonnikov. An initial-boundary value problem for a Stokes system that arises in the study of a problem with a free boundary. *Trudy Mat. Inst. Steklov.*, 188:150–188, 192, 1990. Translated in Proc. Steklov Inst. Math. **1**991, no. 3, 191–239, Boundary value problems of mathematical physics, 14 (Russian).

[34] V. A. Solonnikov. Solvability of a problem on the evolution of a viscous incompressible fluid, bounded by a free surface, on a finite time interval. *Algebra i Analiz*, 3(1):222–257, 1991.

[35] D. L. G. Sylvester. Large time existence of small viscous surface waves without surface tension. *Comm. Partial Differential Equations*, 15(6):823–903, 1990.

[36] A. Tani. Small-time existence for the three-dimensional Navier-Stokes equations for an incompressible fluid with a free surface. *Arch. Rational Mech. Anal.*, 133(4):299–331, 1996.

[37] A. Tani and N. Tanaka. Large-time existence of surface waves in incompressible viscous fluids with or without surface tension. *Arch. Rational Mech. Anal.*, 130(4):303–314, 1995.

[38] S. Wu. Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.*, 130(1):39–72, 1997.

Angel Castro
Departamento de Matemáticas
Facultad de Ciencias
Universidad Autónoma de Madrid
Campus Cantoblanco UAM, 28049 Madrid

Instituto de Ciencias Matemáticas, CSIC-UAM-UC3M-UCM
C/ Nicolás Cabrera 13-15
Campus Cantoblanco UAM, 28049 Madrid
Email: angel_castro@icmat.es

Diego Córdoba
Instituto de Ciencias Matemáticas
Consejo Superior de Investigaciones Científicas
C/ Nicolás Cabrera, 13-15
Campus Cantoblanco UAM, 28049 Madrid
Email: dcg@icmat.es

Charles Fefferman
Department of Mathematics
Princeton University
1102 Fine Hall, Washington Rd,
Princeton, NJ 08544, USA
Email: cf@math.princeton.edu

Francisco Gancedo
Departamento de Análisis Matemático & IMUS
Universidad de Sevilla
C/ Tarfia, s/n
Campus Reina Mercedes, 41012 Sevilla
Email: fgancedo@us.es

Javier Gómez-Serrano
Department of Mathematics
Princeton University
610 Fine Hall, Washington Rd,
Princeton, NJ 08544, USA
Email: jg27@math.princeton.edu